

## 最終氷期以降における西南極氷床の融解史：批判的レビュー

Retreat history of West Antarctic Ice Sheet after the last glacial maximum: A critical review

\*大河内 直彦<sup>1</sup>

\*Naohiko Ohkouchi<sup>1</sup>

## 1. 海洋研究開発機構

1. Japan Agency for Marine-Earth Science and Technology

In the community of Antarctic paleoceanography, it has been long believed that the melting of West Antarctic Ice Sheet (WAIS) started just after the last glacial maximum (LGM), and occurred mainly in the deglaciation period (19-10 kyr BP), a timing of melting of Northern Hemisphere ice sheets such as Laurentide Ice Sheet. In contrast, some geological/geochemical evidences especially from Antarctic Continent have suggested that the melting occurred mainly in the Holocene and even in the latest Holocene. These two contrasting views can be at least partly ascribed to the lack of robust tools for the reconstruction of the ice sheet melting in the sedimentary record. Particularly, chronological framework in the sediments has been a key issue. Our recent evidence with compound-specific radiocarbon dating of the marine sediments suggested that the ice shelf edge in the Ross Sea retreated as large as 400 km during the last 5000 years, confirming above view. In this presentation, I critically review this problem, the timing of melting of WAIS after the LGM. I stress that the importance of precise sediment chronologies for the Antarctic paleoceanography.

南極氷床辺縁部のアイス・ライズとアイス・ランブル：分布と特徴、氷床と棚氷の動力学に与える影響、気候と氷床の変動復元に果たす役割

Antarctic ice rises and rumples: Their properties and significance for ice-sheet dynamics and evolution

\*松岡 健一<sup>1</sup>

\*Kenichi Matsuoka<sup>1</sup>


1.ノルウェー極地研究所

1.Norwegian Polar Institute

棚氷に囲まれ局地的に設置している地域（アイス・ライズまたはアイス・ランブルと呼ぶ）は、南極氷床からの氷の流出、すなわち海水準への寄与を抑制する大きな役割を果たしている。本発表では、昨年末に出版されたレビュー論文（Matsuoka and 19 others, 2015, Earth-Science Reviews）のなかから、アイス・ライズとアイス・ランブルの分布と特徴、氷床と棚氷の動力学に与える影響、気候と氷床の変動復元に果たす役割について概要を述べる。添付の図は、本論文で発表したインヴェントリに含まれる、194のアイス・ライズ（青、位置と大きさを示す）と510のアイス・ランブル（赤、位置のみを示す）の分布である。

キーワード：南極氷床、棚氷、アイス・ライズ

Keywords: Antarctic Ice Sheet, ice shelf, ice rise



## 氷床棚氷モデル IcIES の開発と MISMIP(+) 実験

Development of a numerical ice-sheet/ice-shelf model IcIES  
and its performance on the MISMIP(+) experiments

\*齋藤 冬樹<sup>1</sup>、阿部 彩子<sup>2</sup>、高橋 邦生<sup>1</sup>

\*Fuyuki SAITO<sup>1</sup>, Ayako Abe-Ouchi<sup>2</sup>, Kunio Takahashi<sup>1</sup>

1.国立研究開発法人海洋研究開発機構、2.東京大学大気海洋研究所

1.Japan Agency for Marine-Earth Science and Technology, 2.Atmosphere and Ocean Research Institute,  
The University of Tokyo

Ice sheet model for Integrated Earth-system Studies (IcIES) has been developed to simulate Greenland and Antarctic ice sheets as well as paleo-climate studies of past Northern Hemisphere ice sheets.

Experimental design for Marine Ice-Sheet and Ocean Model Intercomparison Projects (MISOMIP) has been launched (Asay-Davis 2015, GMDD).

Marine Ice-Sheet Intercomparison Projects third phase (MISMIP+) is one of the three subprojects of MISOMIP, which focuses on the comparison among 'stand-alone' ice-sheet/ice-shelf models.

Seven sensitivity experiments are proposed: each of them is a 100-year (optionally 900-year) transient simulation under prescribed basal melting below ice shelf.

This study reports preliminary tests of MISMIP+ experiments using IcIES.

キーワード：棚氷、数値モデリング

Keywords: Ice shelf, Numerical modeling

## 東南極における海洋 - 海氷 - 氷床システムの相互作用と変動の解明

## Research of Ocean-ice BOundary INteraction and Change around Antarctica (ROBOTICA)

青木 茂<sup>2</sup>、\*田村 岳史<sup>1</sup>Shigeru Aoki<sup>2</sup>, \*Takeshi Tamura<sup>1</sup>

1.国立極地研究所、2.低温科学研究所

1.National Institute of Polar Research, 2.Institute of Low Temperature Science

Antarctica and surrounding Southern Ocean are changing. Acceleration of ice mass loss and warming of the coastal ocean in the West Antarctica are the problems of substantial impacts on the global climate system. In the East Antarctica, which has been considered to be stable and attracted relatively less attention, regional characteristics of interactions among climate subsystems have been recently revealed and evidences of variations on various time scales from decades to millennium have been accumulating. Off Wilks Land underneath the Totten Glacier Ice Shelf, whose ice discharge is accelerating, a potential pathway of warm water access has been discovered (Greenbaum et al., 2015). Along the East Antarctic coast, at the same time, sea ice formation and subsequent brine rejection in polynyas, including Cape Darnley Polynya as the head of the list, result in production of Dense Shelf Water and lead to the export of bottom water (Ohshima et al., 2013; Kitade et al., 2014). In the Lutzow-holm Bay off Enderby Land, oceanic temperature variability on decadal time scale was observed, and disintegration/stabilization of the landfast ice and Shirase Glacier Tongue seem to have a quasi-periodicity of one to two decades. In the deep past during the Pliocene when the surface temperature was higher by several degrees than that of the present climate, geological evidence was found for the substantial disintegration of Ice Sheet for the George V Land (Cook et al., 2013). Despite the growing awareness on the importance of ice-ocean interaction and long-term variabilities off the East Antarctic Coast, quantitative descriptions and understandings of the mechanisms are still insufficient. Given the global impact of the coastal variability through the bottom water export, investigations of the mechanisms and variabilities in the East Antarctica are indispensable.

As for the oceans and ice sheets, importance of repeated observations to describe their variabilities and changes, even at an interval of two to three decades, was stressed at least as early as IGY period. From 1976, year-round hydrographic observations have been conducted occasionally in Lutzow-holm Bay by Japanese parties, which provides one of the longest observational records. Together with the asset of long record of tide gauge near Syowa station, Lutzow-holm Bay is the important monitoring site for the description of temporal variability. However, a constant and sustained observation system is not yet established due to the logistic difficulties of sea ice. Even the bathymetric information, which is essential in any discipline of oceanography, is still insufficient. However, recent rapid progress in the techniques of remote autonomous observation and satellite communication are beginning to change this situation. Hence, under the project called ROBOTICA for the coming 9th six-year plan (2016-2023), we plan to utilize state-of-the-art unmanned observations such as under-ice oceanographic, seafloor and cryospheric observations using ROV/AUVs, geodetic network observations of ice/ocean motion and deformation using GPS/ GNSS, and oceanographic observations using tethered and moored profiling observation systems. Combinations with the conventional and robust observational techniques will enable us to acquire the detailed environmental information both in time and space. Implementation of this project can provide us a big step forward for realization of the dream of the sustained observation system around Antarctica. Application of the remote observation techniques to the new horizons such as Totten Glacier and Cape Darnley regions will enhance the understandings of the mechanisms of

different ice-ocean interaction regimes.

## 氷期の大気中二酸化炭素変動に果たす南大洋の役割

Role of Southern Ocean in glacial atmospheric CO<sub>2</sub> reduction\*小林 英貴<sup>1</sup>、岡 顕<sup>1</sup>、阿部 彩子<sup>1</sup>\*Hidetaka Kobayashi<sup>1</sup>, Akira Oka<sup>1</sup>, Ayako Abe-Ouchi<sup>1</sup>

1. 東京大学大気海洋研究所

1. Atmosphere and Ocean Research Institute, The University of Tokyo

過去約 80 万年間の気候は、約 10 万年の周期で温暖な気候と寒冷な気候が繰り返し起こる氷期-間氷期サイクルと呼ばれる気候変動で特徴づけられる。氷期の大気中二酸化炭素濃度(pCO<sub>2</sub><sup>atm</sup>)は、間氷期と比べて約 100 ppmv 低かったことが知られている。これまでにその変動メカニズムに迫るため、海洋大循環モデル(Ocean General Circulation Model: OGCM) を用いた数値実験が多く行われてきたが、モデルでは pCO<sub>2</sub><sup>atm</sup> の変動を再現することができず、メカニズムの詳細には未解明な点が多く残っている。

古気候プロキシを用いた古気候復元から、約 2 万年前の最終氷期には、南大洋底層に高塩分かつ水塊年齢が非常に古い水が存在していたことが示唆されている。これは、氷期の南大洋は塩分による成層が強く、表層から隔離された深層に多量の炭素が保持されていた可能性を示す。そのため、南大洋は氷期-間氷期の pCO<sub>2</sub><sup>atm</sup> の変動メカニズムを議論する上で、重要な海域であることが認識されつつある。本研究では、古気候復元から推測された氷期の南大洋における深層の塩分や水塊年齢の分布を再現したうえで、氷期-間氷期の pCO<sub>2</sub><sup>atm</sup> の変動に対する南大洋の物理過程の寄与を、OGCM を用いた数値実験で定量的に評価した。数値実験の結果から、氷期の南大西洋深層における高塩分データを説明するためには、東南極における深層水形成が重要であったが、そのような塩分場は、海洋の子午面循環の南極底層水起源の北向き流量を増加させるため、海洋中の炭素の滞留時間が小さくなり、pCO<sub>2</sub><sup>atm</sup> を増加させる方向にはたらくことがわかった。一方、成層の強化に伴う鉛直混合の弱さを考慮すると、溶存無機炭素の鉛直勾配が大きくなるため、pCO<sub>2</sub><sup>atm</sup> を減少させる。しかしながら、これら南大洋物理場の変化に加え、海面水温の低下、海洋循環場の変化、鉄肥沃化の寄与のすべてを合わせても、モデルでは氷期の pCO<sub>2</sub><sup>atm</sup> の低下の半分程度しか説明できないことがわかった (Kobayashi et al., 2015)。

氷期-間氷期の pCO<sub>2</sub><sup>atm</sup> 変動のような千年スケールの海洋炭素循環変動を議論する上では、炭酸塩の堆積・溶解によるフィードバック作用である炭酸塩補償過程を考慮する必要がある。氷期には、深層での溶存無機炭素濃度が増加するため、炭酸塩は溶解しやすくなる。炭酸塩の溶解は二酸化炭素分圧を減少させる方向にはたらく。氷期のpCO<sub>2</sub><sup>atm</sup> の低下にも寄与していた可能性が高いと考えられる。これまでの我々の数値実験では、海底に到達した粒子状有機物や炭酸塩は、海底でただちに溶解すると仮定していたため、炭酸塩補償過程を考慮することができていなかった。炭酸塩補償過程の影響を定量的に評価するためには、これまでの海洋内部での再循環のみを考慮した“クローズドシステム”から、海洋と海洋堆積層との物質のやり取りを陽に考慮する“オープンシステム”へとモデルを切り替える必要がある。現在、そのために、新たに海底堆積物モデルを開発し、それをOGCMと結合して数値実験を行うことで、炭酸塩補償過程を含む“オープンシステム”において、氷期のpCO<sub>2</sub><sup>atm</sup> の低下がどのように説明されるのか、改めて定量的に議論することを目指して研究を進めている。講演では、新たに開発した堆積モデルによる数値実験の結果についても報告したいと考えている。

キーワード：炭素循環、氷期-間氷期サイクル、南大洋、子午面循環、堆積モデル

Keywords: carbon cycle, glacial/interglacial, Southern Ocean, meridional overturning circulation, sediment model

## 近年の地球温暖化の停滞における南大洋の役割

## Role of the Southern Ocean in the post-2002 global warming hiatus

\*岡 顕<sup>1</sup>、渡部 雅浩<sup>1</sup>\*Akira Oka<sup>1</sup>, Masahiro Watanabe<sup>1</sup>

1. 東京大学大気海洋研究所

1. Atmosphere and Ocean Research Institute, The University of Tokyo

In spite of increasing atmospheric CO<sub>2</sub> concentration, the warming rate of the global-mean surface temperature slows down in the twenty-first century. Almost all of Coupled Model Intercomparison Project phase5 (CMIP5) climate model simulations fail to reproduce this global warming hiatus. Here, we discuss how the excess energy from top-of-atmosphere is stored in the ocean and quantify the influence of the ocean heat uptake on the surface temperature anomaly. Our numerical simulation, where wind stress anomaly in the equatorial Pacific Ocean is prescribed from reanalysis data, suggests that subsurface warming in the Pacific Ocean takes place at the beginning phase of global warming hiatus (1998-2002) as reported in a previous study. We newly clarify that this subsurface anomaly is transported into the Southern Ocean at the latter phase of hiatus (after 2002), which leads to Southern Ocean heating acceleration below subsurface. The historical observed data of ocean temperature also supports this scenario; warming trend of the Southern Ocean after 2002 is detected in data and its spatial pattern is consistent with our simulation. This result provides us with a clear evidence that the deeper parts of the Southern Ocean has a critical role in the post-2002 global warming hiatus.

キーワード：地球温暖化の停滞、海洋熱吸収

Keywords: global warming hiatus, ocean heat uptake



Temporal and spatial variability of the ACC, fronts and eddies in the Indian Sector of the Southern Ocean derived from satellite radar altimetry

Temporal and spatial variability of the ACC, fronts and eddies in the Indian Sector of the Southern Ocean derived from satellite radar altimetry

\*Mizobata Kohei<sup>1</sup>

\*Kohei Mizobata<sup>1</sup>

1.東京海洋大学

1.Tokyo University of Marine Science and Technology

In the Southern Ocean, the Antarctic Circumpolar Current (ACC) and fronts (e.g., Polar front) have been detected by using in-situ data, satellite data and numerical simulation (e.g., Gille, 1994; Belkin and Gordon, 1996; Feng et al. 2013). Except for numerical simulation, it is difficult to obtain temporal evolution of the ACC and fronts. In this study, the temporal and spatial variability of large-scale features such as the ACC and fronts are extracted from merged absolute dynamic topography (MADT) distributed by CNES/AVISO. To decompose MADT into those features, 2-D FFT and inverse FFT were applied. Results clearly show temporal variability of meandering of the ACC and fronts, and geostrophic velocity field in relation to the ACC variability. Using our method, mesoscale features without long-term trend was also derived, while the AVISO Sea level anomaly contains long-term and nonlinear trend.

キーワード : satellite radar altimetry、南極周極流、中規模渦

Keywords: satellite radar altimetry, Antarctic circumpolar current, mesoscale eddy

## アフリカ南部の気候におけるウェッデル海水変動の役割

## Role of Weddell Sea ice variability in southern African climate

\*森岡 優志<sup>1</sup>、Engelbrecht Francois<sup>2</sup>、Behera Swadhin<sup>1</sup>

\*Yushi Morioka<sup>1</sup>, Francois Engelbrecht<sup>2</sup>, Swadhin Behera<sup>1</sup>

1.海洋研究開発機構、2.Council for Scientific and Industrial Research

1.Japan Agency for Marine-Earth Science and Technology, 2.Council for Scientific and Industrial Research

A potential role of interannual sea ice variability in the Weddell Sea in the southern African climate is investigated through data analysis and coupled general circulation model (CGCM) experiments. The Weddell Sea ice undergoes a distinct interannual variability during the austral summer when the southern African rainfall experiences most of its annual rainfall. The sea ice concentration in the Weddell Sea shows a significantly negative correlation with the southern African rainfall during the season. It is found that the low sea ice concentration anomalies in the Weddell Sea are associated with anticyclonic circulation anomalies to the northeast in the South Atlantic which facilitate more moisture advection from the southern Indian Ocean toward the southern Africa. The composite analysis reveals that the low sea ice concentration anomalies in the Weddell Sea may be attributed to the atmospheric forcing such as the increased solar radiation related to the decreased albedo and the northwesterly wind anomalies. This low sea ice concentration anomalies, in turn, act to cause the skin temperature warmer than normal and reduce the meridional temperature gradient to the north. This is a favoring condition for sustaining the anticyclonic circulation anomalies there by increasing the atmospheric stability. The observed association between the sea ice concentration and atmospheric circulation anomalies is simulated in the CGCM experiment, and the local sea-ice impact on the atmospheric circulation anomalies is also demonstrated in a sensitivity experiment where the interannual sea surface temperature variability in the tropical Pacific is suppressed by the model climatology. These model experiments suggest that besides the remote influence by the tropical Pacific climate variability such as El Niño-Southern Oscillation, the sea ice variability in the Weddell Sea may contribute to the atmospheric variability in the South Atlantic, which may be important for the southern African climate.

キーワード：ウェッデル海氷、経年変動、大気海洋海氷相互作用

Keywords: Weddell Sea ice, Interannual variability, Atmosphere ocean sea-ice interaction

## 南極海の沿岸ポリニヤ域におけるAMSR2薄氷厚アルゴリズムの開発

## Development of a thin ice thickness algorithm of AMSR2 for Antarctic coastal polynyas

\*Nihashi Sohey<sup>1</sup>、大島 慶一郎<sup>2</sup>

\*Sohey Nihashi<sup>1</sup>, Kay I. Ohshima<sup>2</sup>

1.苫小牧高専機械科、2.北大低温研

1.Department of Mechanical Engineering, National Institute of Technology, Tomakomai College,

2.Institute of Low Temperature Science, Hokkaido University

A coastal polynya is newly-forming thin sea-ice areas formed by divergent ice motion driven by prevailing winds and/or ocean currents. In coastal polynyas, huge amounts of heat flux from the ocean to the atmosphere occur because the heat insulation effect of sea-ice is greatly reduced in the case of thin ice, and accordingly sea ice is formed actively. Dense water formed in Antarctic coastal polynyas with the intense sea-ice production is a major source of Antarctic Bottom Water, which is a key player in the global climate system.

In Antarctic coastal polynya areas, algorithms that detect the polynya areas and estimate the thin ice thickness from passive microwave satellite (SSM/I or AMSR-E) data have been developed to estimate the ice and dense water production. The spatial (grid) resolution of AMSR-E, which was launched in May 2002 onboard NASA's Aqua satellite, is four times higher than that of SSM/I in the pixel density. This advantage of AMSR-E is critical for the monitoring of the coastal polynyas because of their fairly small areal extent (i.e., from 10 to 100km at most). Although AMSR-E failed in October 2011, AMSR2 (Advanced Microwave Scanning Radiometer 2), the successor to AMSR-E, was launched in May 2012 onboard the GCOM-W (Shizuku) satellite. The spatial resolution of AMSR2 is improved about 17% from AMSR-E (about 5 km at 89 GHz). In this study, we present a thin ice thickness algorithm for AMSR2 data.

The thin ice algorithm has been developed based on a relationship between polarization ratios (PR) of AMSR2 brightness temperatures (TBs) and thermal ice thickness, as in previous algorithms of AMSR-E. We used AMSR2 TBs at 89 GHz and 36.5 GHz. The thermal ice thickness is based on heat flux calculation using sea-ice surface temperatures derived from satellite thermal infrared images. As the first step of the AMSR2 algorithm development, we used 14 clear-sky MODIS images acquired in the Ross Ice Shelf polynya area. The AMSR2 PR is negatively correlated with the thermal ice thickness. The AMSR2 PR vs. ice thickness relationship is similar to that of AMSR-E. We plan to develop the AMSR2 algorithm also using clear-sky MODIS images acquired in other Antarctic coastal polynya areas, such as the Ronne Ice Shelf and Cape Darnley polynyas.

キーワード：南極海、沿岸ポリニヤ、AMSR2

Keywords: Antarctic Ocean, Coastal polynya, AMSR2

## ポリニヤタイプを考慮した薄氷厚アルゴリズムの開発

Estimation of thin ice thickness discriminating polynya type from AMSR-E passive microwave data

\*中田 和輝<sup>1</sup>、大島 慶一郎<sup>2</sup>、二橋 創平<sup>3</sup>

\*Kazuki Nakata<sup>1</sup>, Kay I. Ohshima<sup>2</sup>, Sohey Nihashi<sup>3</sup>

1.北海道大学大学院環境科学院、2.北海道大学低温科学研究所、3.苫小牧工業高等専門学校

1.Graduate School of Environmental Science, Hokkaido University, 2.Institute of Low Temperature Science, Hokkaido University, 3.National Institute of Technology, Tomakomai College

In Antarctic coastal polynyas, high production of sea ice occurs due to huge heat loss to atmosphere, resulting in the formation of dense water, precursor of Antarctic Bottom Water. Detection of thin ice area and estimation of ice thickness are inevitable for the calculation of sea-ice production. Several studies have developed algorithms for estimation of the thin ice thickness from brightness temperature (TB) of satellite passive microwave sensor [e.g., Martin et al., 2004; Nihashi and Ohshima, 2015]. In these algorithms, ice thickness of less than 20 cm is empirically estimated by utilizing negative correlation between the ice thickness and a ratio of the horizontally to vertically polarized TBs (PR).

Thin ice (polynya) areas are classified roughly into two types. One is frazil ice type: a mixture of open water and grease ice/pancake areas. This type is formed under strong wind condition. The other is nilas or thin consolidated ice type: nearly uniform thin ice covered area. This type is formed under relatively calm condition. It has been speculated that difference in polynya type causes reduced accuracy of the ice thickness algorithm because the two type polynyas likely show quite different microwave characteristics. In this study, we examined the PR-thickness relationship on these two types separately, and developed the thin ice algorithm discriminating the polynya type.

We used 36 GHz and 89 GHz TBs data of Advanced Microwave Scanning Radiometer for EOS (AMSR-E, the resolution of 12.5 and 6.25 km, respectively), MODerate resolution imaging Spectroradiometer (MODIS, the resolution of 1 km) data and the Advanced Synthetic Aperture Radar (ASAR, the resolution of 150 m) data, obtained from the major three polynya around the Antarctica (Ross Ice Shelf polynya (RISP), Ronne Ice Shelf polynya (RONP) and Cape Darnley polynya (CDP)). We estimated sea ice thickness for the AMSR-E footprint using MODIS and ERA-interim atmospheric data. After thin ice areas are divided into the two types from the ASAR data, we examined the AMSR PR-MODIS ice thickness relationship for each polynya type. The result shows the clear difference of PR-thickness relationship between the two polynya types. The exponential fitted curves for frazil and nilas types are similar to those of Martin et al. [2004] and Nihashi and Ohshima [2015], respectively. We considered that the difference of PR-thickness relationship is caused by the presence or absence of open water fraction. Because the ratio of open water in sea-ice region is largely reflected in gradient ratio (GR) of 36 and 89 GHz vertically polarized TBs, the two types can be clearly discriminated by the PR-GR plane. Based on these, we have developed a new thin ice algorithm in which polynyas are classified into the two types by a quadratic discriminant method. Sea ice thickness is estimated from the fitted exponential curve of PR-thickness relationship for each polynya type. Using this algorithm, daily polynya type and ice thickness for the three polynyas have been estimated during April-November in 2003-2010. Sea-ice production has been also estimated by heat flux calculation using the ice thickness and ERA-interim atmospheric data.

It is found that the occurrence frequencies of the two polynya types are largely different among the three polynyas. In the CDP, frazil type is more predominant, compared with other two polynyas.

In previous algorithms, ice thickness was overestimated because the PR-thickness relationship is similar to that of the nilas type. Therefore, sea-ice production in the CDP with high occurrence frequency of frazil type is calculated to be about twice as that of previous studies.

キーワード：沿岸ポリニヤ、氷厚、海氷生産量、AMSR-E

Keywords: coastal polynya, ice thickness, sea-ice production, AMSR-E

## 南大洋インド洋区季節海氷域における溶存炭酸物質の空間分布と時間変化

Temporal and spatial variation of dissolved inorganic carbonates in the summer of Antarctic seasonal sea ice zone

\*橋田 元<sup>1</sup>、高村 友海<sup>1</sup>、野村 大樹<sup>2</sup>、吉川 久幸<sup>3</sup>

\*Gen Hashida<sup>1</sup>, Tomomi Takamura<sup>1</sup>, Daiki Nomura<sup>2</sup>, Hisayuki Yoshikawa<sup>3</sup>

1.大学共同利用機関法人 情報システム研究機構 国立極地研究所、2.北海道大学大学水産学部、3.北海道大学大学院環境科学院

1.Inter-university Research Institute Corporation Research Organization of Information and Systems / National Institute of Polar Research, 2.Hokkaido University, Faculty of Fisheries Science, 3.Hokkaido University, Graduate School of Environmental Science

To clarify the variations of carbonate system in the Southern Ocean, multi-ship observations and line observation which closes ice edge were performed since JARE-51st (2009/10 austral summer) on-board Icebreaker "SHIRASE" and T/V Umitaka-Maru of TUMSAT (Tokyo University of Marine Science and Technology). Although seasonal changes of pCO<sub>2</sub> are attributed to thermodynamics and biological activity, Analysis of DIC vertical profiles reveal that biological uptake and entrainment of sub-surface water played major role and air-sea CO<sub>2</sub> exchange can contribute a little.

キーワード：溶存無機炭酸物質、南大洋、季節海氷域

Keywords: Dissolved inorganic carbonates, Southern Ocean, Seasonal sea ice zone