

## マイクロフォーカスX線CTの紹介と海底資源研究への応用

## Introduction of microfocus X-ray CT scanner and application of marine resources

\*村山 雅史<sup>1</sup>、松崎 琢也<sup>1</sup>、臼井 朗<sup>2</sup>

\*Masafumi MURAYAMA<sup>1</sup>, Takuya Matsuzaki<sup>1</sup>, Akira Usui<sup>2</sup>

1.高知大学海洋コア総合研究センター、2.高知大学自然科学系理学部門

1.Center for Advanced Marine Core Research, Kochi University, 2. Department of Natural Science, Kochi University

マイクロフォーカスX線CT (Zeiss社製) が、高知大学海洋コア総合研究センターに新たに導入された。マイクロフォーカスX線CTは、対象物の内部構造をマイクロ～ナノレベルで非破壊観察が可能であり、地質試料にも応用されている。本発表では、海底資源であるマンガンクラストやマンガンノジュールの内部構造の観察例や他の分析機器と組み合わせた、今後の応用展開についても紹介する。

キーワード：マイクロフォーカスX線CT、海底資源、マンガンノジュール

Keywords: microfocus X-ray CT, marine resources, manganese nodule

## 北西太平洋のマンガングラスト表面に生息する微生物群集の比較解析

Comparative analysis of microbial communities on hydrogenetic ferromanganese crusts in the northwest Pacific

\*加藤 真悟<sup>1</sup>、仁田原 翔太<sup>2</sup>、山岸 明彦<sup>2</sup>

\*Shingo Kato<sup>1</sup>, Shota Nitahara<sup>2</sup>, Akihiko Yamagishi<sup>2</sup>

1.国立研究開発法人海洋研究開発機構、2.東京薬科大学生命科学部

1.Japan Agency for Marine-Earth Science and Technology, 2.Tokyo University of Pharmacy and Life Science

Ferromanganese crusts are commonly found on outcrops of slopes of seamounts in the Pacific, and contain economically valuable elements, such as Co, Ni, Cu, Pt and REE, therefore the crusts are great interest of deep-sea mining. Microorganisms are thought to contribute to formation of the crusts and play a significant role in accumulation of the elements. Actually, the presence of abundant and diverse microorganisms on a ferromanganese crust collected at 3000 m water depth in the Takuyo-Daigo Seamount has been reported (Nitahara et al., 2011). However, our knowledge of commonality and difference in the abundance, diversity and distribution of microorganisms of the ferromanganese crusts is still limited. To assess the commonality and difference, in the present study, we collected ferromanganese crusts from three regions (Takuyo-Daigo Seamount, Ryusei Seamount, and Daito Ridge) at several water depths (1200 m to 2200 m) during the cruises NT09-02, KY11-02 and NT12-25, and analyzed microbial communities of the crusts. In addition, we collected and analyzed surrounding sediments and bottom seawater as references to assess the uniqueness of the crust communities. Cell densities estimated by quantitative PCR were significantly higher in the crusts than the seawater, but comparable to or lower than the sediments. Both of bacterial and archaeal 16S rRNA genes were detected in all samples. Phylogenetic diversities were higher in the crusts than the seawater, but comparable to or slightly lower than the sediments. Comparative analyses of the community compositions showed 1) the presence of unique microorganisms to the crusts, which were not detected in the sediments and seawater, and 2) the presence of common microorganisms among the crusts at every region and almost every depth, which are likely key members for ecosystem functioning on the crusts. Based on the results, microbial contribution to the formation of the crusts will be discussed.

キーワード：コバルトリッチマンガングラスト、微生物群集、北西太平洋

Keywords: Co-rich ferromanganese crust, Microbial community, Northwest Pacific

## 南太平洋環流域の遠洋性粘土に存在する大量の微小マンガングラン粒について

## Abundant manganese microparticles in oxic pelagic clay of the South Pacific Gyre

\*浦本 豪一郎<sup>1</sup>、諸野 祐樹<sup>1</sup>、富岡 尚敬<sup>1</sup>、和穎 朗太<sup>2</sup>、上杉 健太郎<sup>3</sup>、竹内 晃久<sup>3</sup>、星野 真人<sup>3</sup>、鈴木 芳生<sup>3,4</sup>、光延 聖<sup>5</sup>、菅 大暉<sup>6</sup>、宮本 千尋<sup>4</sup>、高橋 嘉夫<sup>4</sup>、稲垣 史生<sup>1</sup>

\*Uramoto Go-Ichiro<sup>1</sup>, Yuki Morono<sup>1</sup>, Naotaka Tomioka<sup>1</sup>, Rota Wagai<sup>2</sup>, Kentaro Uesugi<sup>3</sup>, Akihisa Takeuchi<sup>3</sup>, Masato Hoshino<sup>3</sup>, Yoshio Suzuki<sup>3,4</sup>, Satoshi Mitsunobu<sup>5</sup>, Hiroki Suga<sup>6</sup>, Chihiro Miyamoto<sup>4</sup>, Yoshio Takahashi<sup>4</sup>, Fumio Inagaki<sup>1</sup>

1.国立研究開発法人海洋研究開発機構、2.国立研究開発法人農業環境技術研究所、3.公益財団法人高輝度光科学研究センター、4.東京大学、5.愛媛大学、6.広島大学

1.Japan Agency for Marine-Earth Science and Technology, 2.National Institute for Agro-Environmental Sciences, 3.Japan Synchrotron Radiation Research Institute, 4.University of Tokyo, 5.Ehime University, 6.Hiroshima University

Manganese nodules ubiquitously occur in the abyssal plain, with varying concentrations of metals, including manganese, iron, nickel, cobalt, and copper. Typical manganese nodules observed in pelagic sediments are in centimeter to sub-millimeter size of spherical crusts and often concreted to the pavement over the seafloor. During the Integrated Ocean Drilling Program (IODP) Expedition 329 in 2010, we drilled the entire sedimentary sequence at 6 sites in the ultra-oligotrophic region of the South Pacific Gyre (SPG), where dissolved O<sub>2</sub> and aerobic microbial communities are present from the seafloor to the sediment-basement interface [1]. Massive manganese nodules often occur at the seafloor of these drilling sites, and the subseafloor sediments generally consist of zeolitic metalliferous clay. Using a newly developed sediment observation technique [2], we observed numerous micrometer-scale particles of (ferro-)manganese minerals in the oxic zeolitic clay from present-day to several tens of millions of years. Using a synchrotron-based X-ray microtomography and FIB-SEM-EDS, high-resolution three-dimensional micro-textures of manganese microparticles, as well as elemental compositions, were visualized, suggesting that those particles were deposited from the water column, and well preserved in the widespread area of deep-sea oxic sediments over geologic time.

[1] D'Hondt, S., Inagaki, F., Zarikian, C. A., Abrams, L. J., Dubois, N., Engelhardt, T., Evans, H., Ferdelman, T., Gribsholt, B., Harris, R. N., Hoppie, B. W., Hyun, J.-H., Kallmeyer, J., Kim, J., Lynch, J. E., McKinley, C. C., Mitsunobu, S., Morono, Y., Murray, R. W., Pockalny, R., Sauvage, J., Shimono, T., Shiraishi, F., Smith, D. C., Smith-Duque, C., Spivack, A. J., Steinsbu, B. O., Suzuki, Y., Szpak, M., Toffin, L., Uramoto, G., Yamaguchi, T. Y., Zhang, G., Zhang, X.-H., and Ziebis, W. Presence of oxygen and aerobic communities from seafloor to basement in deep-sea sediment. *Nature Geosciences*, 8(4), 299-304, 2015.

[2] Uramoto, G.-I., Morono, Y., Uematsu, K., and Inagaki, F. An improved sample preparation method for imaging microstructures of fine-grained marine sediment using microfocuss X-ray computed tomography and scanning electron microscopy. *Limnology and Oceanography: Methods*, 12, 469-483, 2014.

キーワード：マンガングラン、微小パーティクル、遠洋性粘土、深海平原、南太平洋環流域

Keywords: manganese, microparticles, pelagic clay, abyssal plain, South Pacific Gyre

## Distributions of Rare Earth Elements between Seawater and Fe-Mn Crust Surface at the Takyo Daigo Seamount, Northwestern Pacific

\*Yohey Suzuki<sup>1</sup>, Ayaka Tokumaru<sup>1</sup>, Yanbei Zhu<sup>2</sup>, Akira Usui<sup>3</sup>

1.Graduate School of Science, The University of Tokyo, 2.National Metrology Institute of Japan (NMIJ), National Institute of Advanced Industrial Science and Technology (AIST), 3.Geology Department, Kochi University

Since the Late Paleocene, ferromanganese crusts (Fe-Mn crusts) have been deposited extensively on the surface of seamounts or plateaus at 400 to 6000 meters below sea level (mbsl) with a thickness range of 1 to 20 cm. Owing to the exceedingly slow growth rates (1 to 10 mm/Myr) and the nm-scale constituents of Fe and Mn oxides/oxyhydroxides, rare earth elements (REEs) are highly concentrated in Fe-Mn crusts from ambient seawater. Over the past two decades, the speciation across the solid-water interface has been intensively studied for many REEs. In terms of the distributions of REEs between seawater and Fe-Mn crusts, apparent distribution coefficients of REEs have been estimated between reference seawater and Fe-Mn crusts with variable locations, depths and ages. Although this compilation provides fundamental insights into the selective enrichment pattern of REEs, it might be deviated from the intrinsic values by taking the heterogeneity of vertical profiles of REE concentrations in seawater adjacent to Fe-Mn crusts and the progressive transformation from metastable Fe and Mn oxides/oxyhydroxides to their stable states in Fe-Mn crusts. For accurate estimation of the apparent distribution coefficients, it is necessary to clarify equilibrium solute and sorbate concentrations in seawater and Fe-Mn crusts, because *in situ* physicochemical conditions represented by high hydraulic pressure, partial pressures of CO<sub>2</sub> and O<sub>2</sub> and constant fluxes of low solute concentrations are difficult to be reproduced experimentally. In addition, the complex nm-scale mineral assemblages of Fe-Mn crusts are impossible to be synthesized. As for sorbate concentrations in Fe-Mn crusts, it is critical to sample from the surface layer in Fe-Mn crusts, which is equilibrated with ambient seawater. To collect ambient seawater adjacent to the outcrop of Fe-Mn crusts, seawater sampling by a submersible is of technical advance in comparison to the deployment of a water sampler such as the rosette from a research vessel. Furthermore, seawater sampling needs to be optimized by developing a sampling device to minimize the levels of contamination and loss of trace elements. In this study, it is aimed to obtain the apparent distribution coefficients of REEs between seawater and Fe-Mn crusts by using vertical profiles of REE concentrations of the surface layer of Fe-Mn crusts and the corresponding ambient seawater from one of the largest deposits of Fe-Mn crusts in the northwestern Pacific. As for Ce, anomalously low concentrations in ambient seawater and high apparent coefficients were evident at depths. We will discuss the potential causes of the Ce anomaly as well as the distribution patterns of the other REEs.

## 鉄マンガングラスト中のジルコニウム-ハフニウム分別挙動

## Fractionation of Zirconium-Hafnium in ferromanganese crusts

\*坂口 綾<sup>1</sup>、稲垣 純平<sup>1</sup>、井上 美南<sup>2</sup>、白井 朗<sup>3</sup>、羽場 宏光<sup>4</sup>、柏原 輝彦<sup>5</sup>、山崎 信哉<sup>1</sup>、末木 啓介<sup>1</sup>、高橋 嘉夫<sup>6</sup>

\*Aya Sakaguchi<sup>1</sup>, Junpei Inagaki<sup>1</sup>, Minami Inoue<sup>2</sup>, Akira Usui<sup>3</sup>, Hiromitsu Haba<sup>4</sup>, Teruhiko Kashiwabara<sup>5</sup>, Shinya Yamasaki<sup>1</sup>, Keisuke Sueki<sup>1</sup>, Yoshio Takahashi<sup>6</sup>

1.筑波大学、2.広島大学、3.高知大学、4.理化学研究所、5.海洋研究開発機構、6.東京大学

1.Univ. of Tsukuba, 2.Hiroshima Univ., 3.Kochi Univ., 4.RIKEN, 5.JAMSTEC, 6.Univ. of Tokyo

Pair of Zirconium (Zr) and Hafnium (Hf) is called "Geochemical twins" due to the consistencies of their ionic valences and radii. As a consequence, this elemental pair shows same behaviour and uniform Zr/Hf ratio, which is theoretically same as that of chondrite meteorite, in the environment. However, the significant fractionation between these elements in natural samples has been found, and it has been proposed as enigma of Zr-Hf fractionation (Niu et al. 2012). The large fractionation of these elemental couple has also been found in the ferromanganese crust (FMC) (Bau 1996). In this study, we attempted to clarify the accumulation mechanism of Zr and Hf in FMCs with some methods including X-ray absorption fine structure (XAFS) technique for synthesised and natural samples.

Six FMC samples were collected from the Takuyo-Daigo and Ryusei seamounts, from 950 m to 3000 m water depth, with hyper-dolphin (remotely operated vehicle) equipped with live video camera and manipulators. Near surface layers (less than 1 mm) of all FMCs were analysed with XRD, and measured elemental concentrations by ICP-MS after the total decomposition of samples. Sequential extraction was conducted by following Koschinsky et al. (1995) to confirm the existence phase of Zr and Hf in FMC. XAFS analysis was also conducted to confirm the mineral composition and speciation of Zr together with chemical reagents, Zr minerals and rock samples as standard materials. Furthermore, distribution coefficients (Kd) and chemical states were determined through the adsorption experiments of Zr and Hf DFO complexes with ferrihydrite and d-MnO<sub>2</sub>.

The major mineral composition of Fe and Mn had no significant variation with the water depth of these seamounts. The concentrations of Zr and Hf were increased with depth, and their ratios were varied without showing any trends. However, these ratios were totally fractionated from those of the seawater (Firdaus et al., 2011), and Hf was more enriched in all FMCs than Zr. The observed Zr chemical states in FMCs were 1) coprecipitation with ferrihydrite, 2) coprecipitation with d-MnO<sub>2</sub>, and 3) basalt-like composition, although the results of sequential extraction showed that Zr and Hf dominantly exist in Fe fraction. From the results of adsorption experiments of Zr and Hf with ferrihydrite and d-MnO<sub>2</sub>, it was found that the Hf-DFO was more adsorbed compared with Zr-DFO. In this case, the observed bond length of Hf-O was significantly shorter than that of Zr-O in the synthesised minerals.

キーワード：鉄マンガングラスト、ジルコニウム、ハフニウム

Keywords: ferromanganese crusts, Zirconium, Hafnium

## 東北日本弧新第三系に産する層状マンガン鉱床についての最近の研究

## A Review: Recent Studies of Neogene Strata-bound Manganese Oxide Deposits From the Northeastern Japan Arc

\*伊藤 孝<sup>1,2</sup>\*Takashi Ito<sup>1,2</sup>

1.茨城大学教育学部、2.海洋研究開発機構海底資源開発資源センター

1.Faculty of Education, Ibaraki University, 2.JAMSTEC, Research and Development (R&amp;D) Center for Submarine Resources

In 1950s and 1960s, Neogene strata-bound manganese oxide deposits distributed in the Northeastern Japan Arc had studied vigorously in the viewpoint of the mineral exploration (e.g. Yoshimura. 1952, 1967). Based on the mineralogy, the major elements chemistry and the geological setting, it has been concluded that these deposits formed by submarine hydrothermal activities. Miura et al. (1992) carried out the geochemical studies including trace elements and rare earth elements (REE) compositions for the Neogene strata-bound manganese oxide deposits from southwestern Hokkaido and also concluded them to having hydrothermal origin. On the other hand, they also pointed out that these deposits showed the positive Ce anomalies, which are different characters from the hydrothermal manganese deposits occurred around the modern spreading centers. Recently, Sakai et al. (in prep.) showed radiolarian stratigraphy in the Fukaura district, one of the main occurrences of the Neogene strata-bound manganese oxide deposit in the Northeastern Japan Arc, and found the long term (ca. 6 to 7 m.y.) hiatus just above the manganese deposit horizon. The long-term exposure of the manganese oxide deposit to oxic seawater may be the cause of the REE pattern showing the positive Ce anomaly.

キーワード：マンガン鉱床、東北日本弧、放射虫

Keywords: Manganese deposits, Northeastern Japan Arc, Radiolarians