

Observations of a viable process for the loss to space of terrestrial atmospheric matter

*Qing-He Zhang¹, Michael Lockwood², John C. Foster³, Qiu-Gang Zong⁴, Malcolm W. Dunlop⁵, Shun-Rong Zhang³, Brian Walsh⁶, Jøran Moen⁷, Bei-Chen Zhang⁸

1.Institute of Space Sciences, Shandong University, Weihai, China, 2.Department of Meteorology, University of Reading, Earley Gate, Post Office Box 243, RG6 6BB, UK., 3.MIT Haystack Observatory, Westford, MA 01886, USA, 4.School of Earth and Space Sciences, Peking University, Beijing, China, 5.Space Sciences Division, SSTD, Rutherford Appleton Laboratory, Didcot, UK, 6.Mechanical Engineering Department and Center for Space Physics, Boston University, Boston, MA, USA., 7.Department of Physics, University of Oslo, Blindern, Oslo, Norway, 8.SOA Key Laboratory for Polar Science, Polar Research Institute of China, Shanghai, China

Earth's atmosphere is protected from direct solar wind erosion by the geomagnetic field. Escape of atmospheric atoms to interplanetary space requires photoionization and subsequent upward ion acceleration to overcome Earth's gravity. However, these ionospheric ions have not yet escaped the terrestrial environment because the Dungey circulation (convection) keeps most within the inner magnetosphere where they either remain or precipitate back into the atmosphere. Key problems in estimating isotopic loss rates on geological scales are quantifying the fluxes of source upflows and evaluating the fraction that subsequently returns to Earth. We present direct evidence of ionospheric ions making their way, via detached regions from the dusk plasmasphere, into the accelerated flows along the magnetopause generated by magnetic reconnection. Using recently developed techniques to find the transit time of open flux tubes across the polar cap, we show that they gain enough energy to escape from the terrestrial environment into interplanetary space.

Keywords: Plasmaspheric plume, Dayside magnetic reconnection, Particle acceleration

位相理論と4衛星観測をもちいた渦中心と極値線同定法

Identifying Vortex Core and Extremum Lines using four Satellites based on Field Topology

*CAI DongSheng¹*DongSheng CAI¹

1.University of Tsukuba

1.University of Tsukuba

Identifying vortices are the key to understanding the turbulence in plasma shear layers. Here, the term 'vortex' or 'vortex core' is associated with a region of Galilean invariance [Jeong and Hussain, 1995]. Unfortunately, no single precise definition of a vortex is currently universally accepted, despite the fact that many space plasma authors claim that many observations have detected "vortices" (as Kelvin-Helmholtz vortices at/around the magnetopause). By using the four satellite velocity data, and Taylor series, we expand the velocity data around the satellites, calculate its first order tensor, and linearly approximate the field. We can identify the vortex structures by using various vortex identification criteria as follows: (i) The first criterion is Q-criterion that defines vortices as regions in which the vorticity energy prevails other energies; (ii) the second criterion is the lambda2-criterion that is related to the minus of the Hessian matrix of the pressure related term; and (iii) the third criterion requires the existence of vortex-core-lines that is the Galilean invariance inside the four satellite tetrahedral region. Using these methods, we can identify and analyze more precisely the 3D vortex using tetrahedral satellite configuration. In the field topology theory, the extremum lines (X-lines) that are other manifolds corresponds to different eigenvalues also can be identified using the same method.

キーワード：渦、xライン

Keywords: vortex, x-line

Global MHD simulation study of the vortex at the magnetopause boundary for the southward IMF and steady solar wind conditions

*Kyung Sun Park¹, Tatsuki Ogino², Dae-Young Lee¹, Raymond J Walker³, Khan-Hyuk Kim⁴, Dong-Hun Lee⁴

1.Chungbuk National University, 2.ISEE, Nagoya University, 3.IGPP, UCLA, 4.Kyung Hee University

We have used a high-resolution and three-dimensional global magnetohydrodynamic (MHD) simulation to study the interaction between the steady solar wind and Earth's magnetosphere during the weak southward IMF. The simulation results show that the vortex like is generated at about $= 11.7R_E$ region (1600LT-1700LT) with a vortex size of about $2.9 R_E$ at the inner boundary of magnetosphere. The vortices are propagating tailward with a velocity of 55 km/s up to 86 km/s. Moreover, the quasi-periodic fluctuations of magnetic field and plasma properties clearly show 8-10 min variations across the vortex. The total magnetic field and density are enhanced in center of the vortex with a bipolar magnetic field perturbation in the field component normal to the magnetopause. Also the velocity is low in the center of the vortex and V_x and V_y components have an opposite polarity across the vortex. Magnetic reconnection favorably occurs in anti-parallel field region with slower shear velocity in the magnetosheath. The magnetic field lines are highly bent by parallel vorticity in the flanks of the magnetopause boundary. We suggest that the reconnection is a mechanism of generating vortex with a periodicity in the dayside during the southward IMF.

Keywords: MHD simulation, vortex, reconnection, magnetopause

Particle dynamics in the electron current layer in collisionless magnetic reconnection
Particle dynamics in the electron current layer in collisionless magnetic reconnection

*銭谷 誠司¹、篠原 育²、長井 嗣信³

*Seiji Zenitani¹, Iku Shinohara², Tsugunobu Nagai³

1.国立天文台、2.宇宙科学研究所、3.東京工業大学

1.National Astronomical Observatory of Japan, 2.Institute of Space and Astronautical Science,

3.Tokyo Institute of Technology

Particle dynamics in the electron current layer in collisionless magnetic reconnection is investigated by using particle-in-cell simulations. Electron motions and velocity distribution functions are studied by directly tracking self-consistent particle trajectories. New classes of electron orbits are discovered: a figure-eight-shaped regular orbit inside the electron jet, another regular orbit on the jet flank boundaries, a Speiser-like noncrossing orbit, and nongyrotropic electrons in the downstream of the jet termination region (a remagnetization front). Based on these discoveries, we will discuss the composition of electron velocity distribution functions, fluid properties of a super-Alfvénic outflow jet, and implications for upcoming MMS observation in the magnetotail.

キーワード：磁気リコネクション、PICシミュレーション、粒子軌道

Keywords: Magnetic reconnection, PIC simulation, Particle dynamics

サブストーム開始に伴う磁気圏近尾部とオーロラアークの発展：新しいサブストーム発生機構のモデル

Near-Earth magnetotail and auroral arc development associated with substorm onset: A new interpretation of substorm triggering

*宮下 幸長¹、平木 康隆²、Angelopoulos Vassilis³、家田 章正¹、町田 忍¹

*Yukinaga Miyashita¹, Yasutaka Hiraki², Vassilis Angelopoulos³, Akimasa Ieda¹, Shinobu Machida¹

1.名古屋大学宇宙地球環境研究所、2.電気通信大学情報理工学部、3.Department of Earth, Planetary, and Space Sciences, University of California, Los Angeles

1.Institute for Space-Earth Environmental Research, Nagoya University, 2.Department of Communication Engineering and Informatics, University of Electro-Communications, 3.Department of Earth, Planetary, and Space Sciences, University of California, Los Angeles

Using data from Time History of Events and Macroscale Interactions during Substorms (THEMIS) spacecraft and ground-based observatories at high temporal and spatial resolutions, we studied the time sequence of near-Earth magnetotail and auroral arc development associated with a substorm onset. We discuss four steps of auroral development, auroral fading, initial brightening of an auroral onset arc, enhancement of the arc's wave-like structure, and poleward expansion, and link them to magnetotail changes. A case study shows that near-Earth magnetic reconnection began at $X \sim -17 R_E$ at least ~ 1 min before auroral fading and ~ 3 min before initial auroral brightening. Large-scale ionospheric convection was also enhanced just before auroral fading and before initial auroral brightening. Then low-frequency waves were amplified in the plasma sheet at $X \sim -10 R_E$, with the pressure increase likely due to arrival of an earthward flow from the near-Earth reconnection site ~ 4 min after initial auroral brightening and ~ 50 s before enhancement of the wave-like auroral structure. Dipolarization began ~ 7 min after initial auroral brightening and ~ 30 s before auroral poleward expansion. On the basis of these observations, we suggest that near-Earth magnetic reconnection plays two roles in substorm triggering. First, it generates a fast earthward flow and Alfvén waves. When the Alfvén waves, which propagate much faster than the fast flow, reach the ionosphere, large-scale ionospheric convection is enhanced, leading to auroral fading, initial brightening, and gradual growth of the wave-like auroral structure. Second, when the reconnection-initiated fast flow reaches the near-Earth magnetotail, it promotes rapid growth of an instability, such as a ballooning instability, and the wave-like auroral structure is further enhanced. When the instability has grown sufficiently, dipolarization and auroral poleward expansion are initiated.

キーワード：サブストーム、オーロラアーク、オーロラ爆発、磁気圏尾部、磁気リコネクション、サブストームトリガー

Keywords: substorm, auroral arc, auroral breakup, magnetotail, magnetic reconnection, substorm triggering

Response of the incompressible ionosphere to the compression of the magnetosphere during the geomagnetic sudden commencements

*Takashi Kikuchi¹, Kumiko Hashimoto², Ichiro Tomizawa³, Yusuke Ebihara⁴, Yukitoshi Nishimura⁵, Tohru Araki⁶, Atsuki Shinbori⁴, B. Veenadhari⁷, Takashi Tanaka⁸, Tsutomu Nagatsuma⁹

1.Institute for Space-Earth Environmental Research, Nagoya University, 2.Kibi International University, 3.Center for Space Science and Radio Engineering, University of Electro-Communications, 4.Research Institute for Sustainable Humanosphere, Kyoto University, 5.Department of Atmospheric and Oceanic Sciences, University of California, 6.Geophysical Institute, Kyoto University, 7.Indian Institute of Geomagnetism, 8.International Center for Space Weather Science and Education, Kyushu University, 9.National Institute of Information and Communications Technology

The ionospheric plasma in midlatitude moves upward/downward during the geomagnetic sudden commencement causing the HF Doppler frequency changes; SCF (+ -) and (- +) on the day- and night-sides, respectively, except for the SCF (+ -) in the evening as found by Kikuchi et al.[1985]. Although the preliminary and main frequency deviations (PFD, MFD) of the SCF have been attributed to the dusk-to-dawn and dawn-to-dusk potential electric fields, there still remain questions if the positive PFD can be caused by the compressional magnetohydrodynamic (MHD) wave and what causes the evening anomaly of the SCF. With the HF Doppler sounder, we show that the dayside ionosphere moves upward toward the sun during the main impulse (MI) of the SC, when the compressional wave is supposed to push the ionosphere downward. The motion of the ionosphere is shown to be correlated with the equatorial electrojet (EEJ), matching the potential electric field transmitted with the ionospheric currents from the polar ionosphere. We confirmed that the electric field of the compressional wave is severely suppressed by the conducting ionosphere and reproduced the SC electric fields using the global MHD simulation in which the potential solver is employed. The model calculations well reproduced the PI and MI electric fields and their evening anomaly. It is suggested that the electric potential is transmitted from the polar ionosphere to the equator by the TM_0 mode waves in the Earth-ionosphere waveguide. The near-instantaneous transmission of the electric potential leads to instantaneous global response of the incompressible ionosphere.

Keywords: Incompressible ionosphere, geomagnetic sudden commencement, TM_0 mode wave, ionospheric current, potential electric field

Magnetosphere-Ionosphere Coupling by Alfvén waves during Geomagnetic Storms

*Christopher Carew Chaston^{1,2}, Van Allen Probes Team

1.University of California Berkeley, 2.University of Sydney

We present observations from the Van Allen probes which show the coincident occurrence of Alfvénic fluctuations in the equatorial plane, field-aligned electrons and outflowing energized ionospheric ions during geomagnetic storms. Based on these observations we build a model for the observed wave-fields and the particle acceleration that occurs within them. It is shown how Alfvén waves extract ions from the topside ionosphere and through the action of trapping and stochastic acceleration drive the acceleration of these ions to energies which may exceed 100 keV in the equatorial plane. The agreement in the form and evolution of the observed and simulated ion distributions provides confidence in the veracity of the modelling and simulation results. It is estimated for observed wave amplitudes and ion densities that this process may make a significant contribution to magnetospheric ion energy density. This is supported by statistical results which reveal an inverse correlation between the time rate of change of Dst and Alfvén wave spectral energy density.

Keywords: Alfvén waves, ionosphere, ring current

磁気圏-電離圏フィードバック結合をともなうバルーニング不安定性

Ballooning instability with the magnetosphere-ionosphere feedback coupling

*渡邊 智彦¹*Tomo-Hiko Watanabe¹

1.名古屋大学大学院理学研究科

1.Graduate School of Science, Nagoya University

The ballooning instability has been investigated as a possible mechanism for triggering the substorm in the magnetosphere. The ballooning mode is destabilized, when the interchange term provided with the pressure gradient and the magnetic curvature overcomes the line bending term causing the shear Alfvén waves. In the magnetosphere-ionosphere (M-I) coupling system in polar regions, on the other hand, the shear Alfvén waves (or the kinetic Alfvén waves) can also be destabilized by the magnetospheric convection, if the ionospheric density change is taken into account with the feedback mechanism [1-3].

In the present study, we have investigated the ballooning and the feedback instabilities in the same theoretical framework. Our linear analysis demonstrates that, as the interchange term increases, the "unstable" shear Alfvén waves with the opposite sign of the real eigenvalues in their lowest harmonic branch collide with each other and transit to the ballooning mode. It implies that competitions and/or interactions between the two instabilities may provide a plausible explanation of auroral breakup triggered through the M-I coupling.

[1] T. Sato, J. Geophys. Res., 83, doi:10.1029/JA083iA03p01042.

[2] T.-H. Watanabe, Phys. Plasmas, 17, 022904 (2010).

[3] T.-H. Watanabe, Geophys. Res. Lett., 41, doi:10.1002/2014GL061166 (2014).

キーワード：オーロラ、磁気圏-電離圏結合、不安定性

Keywords: aurora, magnetosphere-ionosphere coupling, instability

What if the evolution of auroral forms does not reflect magnetospheric processes?

*Shinichi Ohtani¹

1.The Johns Hopkins University Applied Physics Laboratory

We often find auroral images very helpful for diagnosing magnetospheric processes, especially from a global point of view, supplementing spatially sparse satellite observations. The assumption that is very often made, if not explicitly, is that the temporal and spatial development of auroral forms reflects that of the corresponding magnetospheric processes. Although this assumption may be reasonable in many cases, caution needs to be exercised since the aurora is a manifestation of complex coupling between the magnetosphere and ionosphere, and the ionosphere does not always respond passively. In this presentation I shall discuss, with an actual example, how our perspective would change if the foundation of this assumption is not as solid as we generally consider.

Keywords: Magnetosphere-ionosphere coupling, Aurora, Field-aligned currents

オーロラの残像に見られる複雑な高速流

Complex and fast motion of the afterglow of aurora

*内田 ヘルベルト陽仁^{1,2}、片岡 龍峰^{2,1}、福田 陽子³、三好 由純⁴、海老原 祐輔⁵、Hampton Donald⁶、Nicolls Michael⁷

*Herbert Akihito Uchida^{1,2}, Ryuho Kataoka^{2,1}, Yoko Fukuda³, Yoshizumi Miyoshi⁴, Yusuke Ebihara⁵, Donald Hampton⁶, Michael Nicolls⁷

1.総合研究大学院大学、2.国立極地研究所、3.東京大学大学院、4.名古屋大学宇宙地球環境研究所、5.京都大学生存圏研究所、6.Geophysical Institute, University of Alaska Fairbanks、7.Center for Geospace Studies, SRI International

1.SOKENDAI (The Graduate University for Advanced Studies), 2.National Institute of Polar Research, 3.University of Tokyo, 4.Institute for Space-Earth Environmental Research, Nagoya University, 5.Research Institute for Sustainable Humanosphere, Kyoto University, 6.Geophysical Institute, University of Alaska Fairbanks, 7.Center for Geospace Studies, SRI International

Afterglow of aurora at 732 nm can be used to track the plasma flow in the topside ionosphere at 300 km altitude (Dahlgren et al., 2009). From ground-based high-speed (50 fps and 200 fps) imaging observations made by Kataoka et al. (2015) at Poker Flat Research Range in Alaska, we found two interesting afterglow events on January 7 and January 28 2015, in which the speed of the brightest arc is suddenly accelerated within the field-of-view (15 deg by 15 deg). Both events occurred near the substorm onset. The existence of the afterglow is confirmed at wavelength range of >665 nm, and any contributions from major forbidden lines at 557.7 and 630.0 nm are rejected. The lifetime of the afterglow is an order of 5 s, which is consistent with the expected emissions from metastable O⁺ ions at 300 km altitude. The motion of afterglow is complicated and contains counterclockwise rotation. The maximum speed of the motion is more than an order of magnitude faster than those previously reported by Dahlgren et al. (2009). We report the results of the analysis of the optical flow to investigate the motion of afterglow, combined with the ionospheric conditions obtained by Poker Flat Incoherent Scatter Radar, and discuss the possibility of such a fast and complex plasma flow in the topside ionosphere. In addition, we have started our high-speed observation for this season in January. The accurate timing measurement of the imaging system has been realized by installing a newly developed GNSS timing board. That makes it possible to perform a synchronized observation with multiple cameras in different places. We also report our newly developed system in this presentation.

キーワード：オーロラ、残像、高速撮像

Keywords: Aurora, Afterglow, High-speed imaging

フリッカリングオーロラの微細構造

Fine scale morphology of flickering aurora

*福田 陽子¹、片岡 龍峰²、三好 由純³、内田 ヘルベルト陽仁²、加藤 雄人⁴、西山 尚典²、塩川 和夫³、海老原 祐輔⁵、Hampton Donald⁶、岩上 直幹¹、関 華奈子¹

*Yoko Fukuda¹, Ryuho Kataoka², Yoshizumi Miyoshi³, Herbert Akihito Uchida², Yuto Katoh⁴, Takanori Nishiyama², Kazuo Shiokawa³, Yusuke Ebihara⁵, Donald Hampton⁶, Naomoto Iwagami¹, Kanako Seki¹

1.東大、2.極地研/総研大、3.名大ISEE、4.東北大、5.京大生存圏、6.アラスカ大学フェアバンクス校

1.UTokyo, 2.NIPR/SOKENDAI, 3.ISEE, Nagoya Univ., 4.Tohoku Univ., 5.RISH, Kyoto Univ., 6.Univ. of Alaska Fairbanks

The generation mechanism of flickering auroras has been considered as Landau resonance between electrons and electromagnetic ion cyclotron (EMIC) waves. The typical frequencies of the flickering aurora are 3-15Hz which correspond to oxygen ion cyclotron frequency at altitudes of 3000-10000 km. The necessary condition of the appearance of the flickering aurora remains unsolved. Recent cameras enable us to see the extreme situation and the moments when the flickering aurora appears or disappears. An sCMOS camera we used in this study has two advantages: a high spatial resolution and a high sampling rate. Observations taken the former advantage were already conducted during two winter seasons. The camera captures fine images at 50 frames per second (fps). The field of view and the spatial resolution at the 100 km altitude is 26.6 x 26.6 km and 52 m, respectively. We found that the modulation of small flickering patches less than 1 km occur in narrow frequency variations while the modulation of large flickering patches with a few kilometers scale occur in broad frequencies. Such a tendency is consistent with the dispersion relation of O⁺ EMIC waves. On the subject of the auroral intensity, the camera is capable of detecting extremely bright auroras without causing a saturation. The flickering aurora occur in the bright and extremely bright auroras. The ranges of the auroral intensity are roughly estimated approximately 10-100 kR at 557.7 nm by comparison with a keogram obtained from a meridian spectrograph at PFRR. We also found that the frequency ranges of the flickering aurora are constant, but the ratio of the flickering amplitude to the steady auroral intensity systematically increases with the decreasing of the steady auroral intensity. The result may suggest the slightly energetic electrons tend not to fulfill the resonance condition of EMIC waves. We also found some flickering auroras coincidentally occur with an isolated magnetic impulse in the Pc5 ULF range observed by magnetometer nearby PFRR. Such a tendency may indicate that mesoscale (of the order of 100 km) traveling current vortices are one of the necessary conditions to generate the flickering aurora. We will discuss on the relation between the excitation of EMIC waves and the traveling auroral arcs with the current centralization. To verify the possibility of EMIC waves as the generation mechanisms, we are now challenging a high-speed imaging at 320 fps which is taken the latter advantage. It has the potential to capture faster modulations by H⁺ and He⁺ EMIC waves. We will collect the data in April 2016, and will report initial results in this talk.

2015年3月14日に観測されたディスクリートアークのオーロラトモグラフィ解析初期結果

Preliminary results of auroral tomography analysis of discrete arcs observed on March 14, 2015

*田中 良昌¹、小川 泰信¹、門倉 昭¹、Gustavsson Bjorn²、Partamies Noora³、Kauristie Kirsti⁴、Whiter Daniel⁵、Enell Carl-fredrik⁶、Braendstroem Urban⁷、Sergienko Tima⁷、Kozlovsky Alexander⁸、宮岡 宏¹、Kosch Mike⁹

*Yoshimasa Tanaka¹, Yasunobu Ogawa¹, Akira Kadokura¹, Bjorn Gustavsson², Noora Partamies³, Kirsti Kauristie⁴, Daniel Whiter⁵, Carl-fredrik Enell⁶, Urban Braendstroem⁷, Tima Sergienko⁷, Alexander Kozlovsky⁸, Hiroshi Miyaoka¹, Mike Kosch⁹

1.国立極地研究所、2.UiT The Arctic University of Norway, Norway、3.The University Centre in Svalbard, Norway、4.Finnish Meteorological Institute, Finland、5.University of Southampton, UK、6.EISCAT Scientific Association、7.Swedish Institute of Space Physics, Sweden、8.Sodankyla Geophysical Observatory, Finland、9.South African National Space Agency, South Africa

1.National Institute of Polar Research, 2.UiT The Arctic University of Norway, Norway, 3.The University Centre in Svalbard, Norway, 4.Finnish Meteorological Institute, Finland, 5.University of Southampton, UK, 6.EISCAT Scientific Association, 7.Swedish Institute of Space Physics, Sweden, 8.Sodankyla Geophysical Observatory, Finland, 9.South African National Space Agency, South Africa

We conducted a campaign of auroral tomography observation using multi-point imager network and the EISCAT UHF radar in Northern Europe in March, 2015. During this campaign, an auroral breakup was observed at 23:15 UT on March 14 by three all-sky EMCCD imagers and three wide-view CCD imagers. Wavy structure of discrete arcs was often observed around the magnetic zenith at Tromso from 22 to 23 UT and pulsating aurora was observed after the breakup. The monochromatic (427.8 nm) images were taken at a sampling interval of 2 seconds by the three EMCCD imagers and at an interval of 10 seconds by all the six imagers. The EISCAT UHF radar at Tromso measured the ionospheric parameters along the field line at the magnetic zenith from 20 to 24 UT.

We apply the auroral tomography method to these data set to reconstruct reliable three-dimensional distribution of the 427.8 nm emission, that will allow us to investigate quantitatively the following subjects; (1) relation between the 427.8 nm emission and electron density enhancement along the field line, (2) spatial distribution of energy of precipitating electrons in the wavy structure, in particular, relation between the energy of precipitating electrons and thickness of discrete arc, (3) relation between motion of the wavy structure and spatial distribution of the ionospheric conductivity, and (4) spatial and temporal variations of energy distribution of precipitating electrons at the auroral breakup. We present preliminary results from the auroral tomography analysis of the discrete arcs.

キーワード：オーロラ、トモグラフィ解析、3次元分布、電離圏、EISCATレーダー

Keywords: aurora, tomography analysis, 3D distribution, ionosphere, EISCAT radar

脈動オーロラに伴う電離圏変動

Ionospheric variation during pulsating aurora

*細川 敬祐¹、小川 泰信²*Keisuke Hosokawa¹, Yasunobu Ogawa²

1.電気通信大学大学院情報理工学研究科、2.国立極地研究所

1.Department of Communication Engineering and Informatics, University of Electro-Communications,

2.National Institute of Polar Research

We have statistically analyzed data from the European Incoherent SCATter (EISCAT) UHF/VHF radars in Tromsø (69.60N, 19.20E), Norway to reveal how the occurrence of pulsating aurora (PsA) modifies the electron density profile in the ionosphere. By checking 5 winter seasons (2007-2012) observations of all-sky aurora cameras of National Institute of Polar Research (NIPR) in Tromsø, we have extracted 21 cases of PsA. During these PsA events, either UHF or VHF radar of EISCAT was operative and the electron density profiles were obtained along the field-aligned or vertical direction near the zenith. From these electron density measurements, we calculated hmE (E region peak height) and NmE (E region peak density), which are proxies for the energy and flux of the precipitating PsA electrons, respectively. Then, we examined how these two parameters changed during the evolution of 21 PsA events in a statistical fashion. The results can be summarized as follows: (1) hmE is lower (the energy of precipitation electrons is higher) during the periods of PsA than that in the surrounding interval, (2) When NmE is higher (flux of PsA electrons is larger), hmE tends to be lower (precipitation is harder), (3) hmE is lower and NmE is larger in the later magnetic local time, (4) When the AE index during the preceding substorm is larger, hmE is lower and NmE is larger. These tendencies are discussed in terms of the characteristics of particles and plasma waves in the source of PsA in the magnetosphere. In addition to the statistics of the EISCAT data, we carried out several detailed case studies, in which the altitude profiles of the electron density were derived by separating the ON and OFF phases of PsA. This allows us to estimate the true altitude profiles of the PsA ionization, which can be used for estimating the characteristic energy of the PsA electrons and better understanding the wave-particle interaction process in the magnetosphere.

キーワード：脈動オーロラ、電離圏、磁気圏

Keywords: Pulsating Aurora, Ionosphere, Magnetosphere

A quantification method for the properties of diffuse and pulsating aurorae based on auroral image data

A quantification method for the properties of diffuse and pulsating aurorae based on auroral image data

*中野 慎也¹、小川 泰信²

*Shin'ya Nakano¹, Yasunobu Ogawa²

1.情報・システム研究機構 統計数理研究所、2.情報・システム研究機構 国立極地研究所

1.The Institute of Statistical Mathematics, 2.National Institute of Polar Research

The auroral images provide much information on the state of the magnetosphere-ionosphere system. However, since the variations of aurorae are highly complicated, it was difficult to quantify the dynamical properties of the aurorae. We are developing a technique for obtaining quantitative metrics about various auroral properties by analyzing a sequence of auroral images taken with high temporal resolution. We introduce a state space model to describe the translational motion of diffuse aurorae, and estimate the translational velocity by using an algorithm which approximates the Kalman filter. We also extract the features of pulsating aurora, patches of pulsating aurora by using a technique based on a sparse modelling. We will report some results of our analyses.

キーワード : pulsating aurora、kalman filter、sparse modelling

Keywords: pulsating aurora, kalman filter, sparse modelling

オメガバンドオーロラの統計的特性

Statistical feature of Omega band aurora observed by THEMIS all-sky imager network

*佐藤 夏雄^{1,2}、門倉 昭^{1,2}、田中 良昌^{1,2}、堀 智昭³、行松 彰^{1,3}*Natsuo Sato^{1,2}, Akira Kadokura^{1,2}, Yoshimasa Tanaka^{1,2}, Tomoaki Hori³, Akira Sessai Yukimatu^{1,3}

1.国立極地研究所、2.総研大、3.名古屋大学宇宙地球環境研究所

1.National Institute of Polar Research, 2.SOKENDAI, 3.Institute for Space-Earth Environment Research, Nagoya University

We examine morphological and statistical characteristics of Omega band auroras. In order to study statistical feature of Omega band aurora we used the THEMIS ground-based all-sky imager network data for 8 years from January 2007 to December 2014. Omega band auroras show their own distinctive behavior on shape and movement, so we could easily pick up an Omega band auroral event from the THEMIS summary plot database on keogram and all-sky image. Consequently, we could identify ~330 events of Omega band type auroras.

In this study we examine statistical features for the following parameters when Omega aurora observed; 1) magnetic local time (MLT), 2) seasonal variations, 3) annual variations, 4) life time, 5) drift speed, 6) scale size, 7) Kp and AE index dependence, 8) solar wind speed and pressure, 9) IMF By and Bz dependence. Then we will discuss their signature.

キーワード：オーロラ、オメガバンドオーロラ、磁気圏、電離圏、磁気圏－電離圏結合、極域

Keywords: aurora, omega band aurora, magnetosphere, ionosphere, magnetosphere-ionosphere coupling, polar region

Precipitation of high-energy particles at high latitudes and impact on middle atmosphere

*Esa Turunen¹

1.Sodankyla Geophysical Observatory, University of Oulu

When trying to understand the role of precipitating high-energy particles in variations of atmospheric properties, we are still facing the fact that accurate measurements throughout the chain from the origin of the particles, their acceleration and interactions in the magnetosphere-ionosphere system down to details of final atmospheric effects are limited. Similarly models mostly cover only specific regions and a consistent holistic model is not available. However, recent individual studies have shown for example evidence of energetic electron precipitation causing statistically significant decrease of upper stratospheric and mesospheric ozone during extended periods of time. Indeed, we need to include energetic electron precipitation as a process in general atmospheric circulation models, if we want to understand our atmosphere as a whole. Here we first review shortly the impact of energetic particles in atmosphere in general, and present the current status of knowledge in chemical variations of atmosphere caused by these particles, including galactic cosmic rays, solar protons and electrons of magnetospheric origin. The effects are both direct and indirect by first generating chemically active minor constituents of the atmosphere, such as odd nitrogen and odd hydrogen, which in turn can affect atmospheric ozone via catalytic reactions either directly in-situ, or after transport in atmosphere to lower altitudes and lower latitudes. Then we discuss recent advance in studying the effects of high-energy electron precipitation in atmosphere. In order to assess the role of precipitating particles in atmospheric variations one needs new measurements which characterize more accurately the energy and flux, as well as spatial and temporal variations of the energetic electron precipitation, both at high and subauroral latitudes, such as given for example by the Japanese ERG satellite mission, so that combined studies using advanced ground-based and satellite measurements together with theoretical modeling would be possible.

Keywords: high-energy particle precipitation, coupling of atmospheric regions, chemical variations

Plasmaspheric modeling using the new Ionosphere-Plasmasphere-Electrodynamics (IPE) model

*Naomi Maruyama^{1,2}, Phil Richards³, Mariangel Fedrizzi^{1,2}, Tzu-Wei Fang^{1,2}, Tim Fuller-Rowell^{1,2}, Mihail Codrescu²

1.Univ. of Colorado at Boulder, 2.NOAA Space Weather Prediction Center, 3.George Mason Univ.

Cold plasma in the plasmasphere and plumes plays an important role in wave-particle interactions in the inner magnetosphere. Plumes regulate the excitation of electromagnetic ion cyclotron (EMIC) waves. Plumes modulate the resonant EMIC wave-particle interactions by energization of cold ions and the pitch angle scattering of the ring current hot ions and the radiation belt relativistic electrons. Furthermore, a remarkable correlation has been observed between the inner edge of the outer radiation belt electrons and the innermost plasmopause location. Very recent observations show that plasmaspheric plumes can influence the dayside magnetic reconnection rate.

A newly developed global three-dimensional ionosphere-plasmasphere-electrodynamics (IPE) model is used to understand the dynamical redistribution of the cold plasma in the plasmasphere and its coupling to the ionosphere-thermosphere system. The IPE model reproduces the ionospheric Storm Enhanced Density (SED) plumes as frequently observed in TEC during geomagnetically active conditions. The SED plumes are transported into the cusp and over the pole due to the high latitude convection, which is characterized as Tongues of Ionization (TOIs). The model captures the corresponding formation of the plasmaspheric drainage plume-like structure in the magnetospheric equatorial plane as reported in previous studies. The plumes gradually start to rotate around the earth, and the plasmasphere gradually refills from the ionosphere as the storm time convection weakens. In this presentation, the temporal and spatial evolution of the redistribution of cold plasma between the plasmasphere and ionosphere is examined depending on the types of the solar wind driving conditions. Furthermore, the role of the Sub-Auroral Polarization Streams (SAPS) electric field is evaluated in draining the plasmaspheric plasma and plumes, as SAPS field penetrates into the plasmasphere due to enhanced convection. An example of how the ERG spacecraft will measure the dynamical evolution of the plasmasphere and plumes is demonstrated.

Keywords: plasmasphere, Modeling and forecasting, coupling to ionosphere-thermosphere, ERG mission

Data assimilation of low-altitude magnetic perturbations into a global magnetosphere model

*Dmitri Kondrashov¹, Slava Merkin²

1.University of California, Los Angeles, 2.Johns Hopkins University Applied Physics Laboratory

The ionosphere is the only region of the terrestrial magnetosphere-ionosphere system where in situ observations with high temporal resolution and approaching global spatial scale are possible. Ionospheric measurements of magnetic fields with such spatio-temporal coverage have become available from the Active Magnetosphere and Planetary Electrodynamics Response Experiment (AMPERE), combining data from the Iridium satellites. Motivated by the emergence of this dataset, we report here on the first results of assimilation of low-altitude ionospheric magnetic perturbations into the Lyon-Fedder-Mobarry (LFM) global magnetospheric model coupled with the Rice Convection Model (RCM). Our assimilation approach relies on the assumption of a quasi-steady, linear approximate relation between equatorial magnetospheric pressure and ionospheric field-aligned currents. This approximation is implemented numerically by perturbing the coupled LFM-RCM model and considering only large-scale modes from the Fourier decomposition of the ionospheric magnetic field and equatorial magnetospheric pressure. This methodology was validated by using model-based assimilation tests of the so-called "fraternal-twins" type. In this approach, the LFM-RCM model with one set of parameters is used to generate synthetic observations, while a model version with a different parameters is used to assimilate the ionospheric observations and calculate the magnetospheric pressure corrections which are then applied to reproduce the synthetic observations. The model with assimilated synthetic data responded correctly by modifying ionospheric currents and magnetic perturbations in the expected way. We thus found the approach proposed herein to be promising for future assimilation of real data.