

MUレーダー実時間アダプティブクラッター抑圧システムの開発

Development of MU radar real-time processing system with adaptive clutter rejection

*橋口 浩之¹、万城 孝弘¹、山本 衛¹、佐藤 亨²、西村 耕司³、橋本 大志²

*Hiroyuki Hashiguchi¹, Takahiro Manjou¹, Mamoru Yamamoto¹, Toru Sato², Koji Nishimura³, Taishi Hashimoto²

1.京都大学生存圏研究所、2.京都大学情報学研究科、3.国立極地研究所

1.Research Institute for Sustainable Humanosphere, Kyoto University, 2.Graduate School of Informatics, Kyoto University, 3.National Institute of Polar Research (NIPR)

大気レーダー観測において、しばしば強いクラッターエコー(山や建物からのエコー)が問題になることがある。そのクラッター抑圧法としてNC-DCMP (Norm Constrained-Directionally Constrained Minimum Power)法が提案され、MUレーダーによる実観測データに適用し、効果があることが実証されている[Nishimura et al., JTech., 2012]。この処理は、これまではオフラインで実施されていた。そこで本研究では、NC-DCMP法によるクラッター抑圧処理をMUレーダーのオンライン処理システムとして実装することを目的とする。これにより、観測データの記録容量を数百分の1に削減でき、外部記憶装置などの制約の少ない標準観測を行うことが可能になる。NC-DCMP法では、所望信号方向を固定した上で、ウエイトベクトルのノルムをある値以下に制約して、信号電力を最小化するように制約条件付最適化問題を解く。

MUレーダーでは30年以上に渡って、毎月100時間程度の対流圏・成層圏標準観測モードによる観測を継続している。まず、この標準観測モードにNC-DCMP処理を実装した。このモードでの観測データは8秒に1回取得される。そのため、実時間でクラッター抑圧を行うためには全ての信号処理を8秒以内に行う必要があるが、処理方法の工夫により、NC-DCMP法の処理時間を平均1.0秒にまで高速化した。山や建物からのエコーは時間的に大きく変化しないため、インコヒーレント積分7回分(約1分間)の受信信号を用いて最適ウエイトベクトルを求めるようにしたところ、良好な結果を得た。2015年11月の標準観測からNC-DCMP処理を適用しているが、安定運用できている。完全ではないが、航空機からのエコーについても、抑圧する効果が得られている。

インドネシア共和国の西スマトラに建設が計画されている赤道MUレーダーは、八木アンテナ19本を1群とする55群構成で、各群からの受信信号を独立に取得可能なシステムが提案されている。本研究の成果は、この赤道MUレーダーにも適用可能である。

キーワード：大気レーダー、クラッター抑圧、ウエイトノルム拘束付DCMP法、MUレーダー

Keywords: Atmospheric radar, Clutter rejection, NC-DCMP method, MU radar

Variability of Rain Microstructure over Sumatra from Micro Rain Radar Observation

*Marzuki Marzuki¹, Indah Rahayu¹, Hiroyuki Hashiguchi², Toyoshi Shimomai³

1.Department of Physics, Andalas University, 2.Research Institute for Sustainable Humanosphere (RISH), Kyoto University, 3.Shimane University

The vertical profile of rain microstructure such as raindrop size distribution (DSD) is important for understanding rain formation processes and to improve the accuracy of radar-derived rainfall estimates. The variability in the DSD substantially limits the accuracy of the DSD application. The aim of the present study is to investigate the natural variability of vertical profile of DSDs as a function of large-scale atmospheric conditions and storm characteristics using measurements from the Equatorial Atmospheric Radar (EAR) facility at Kototabang, west Sumatra, Indonesia. This study is a follow-up of our previous work on the DSD variability over Sumatra [1,2]. The vertical profile of DSD is obtained from a Micro Rain Radar (MRR) with a near continuous record of operation over four consecutive years (2011-2014). The DSDs from the radar are compared with those from an optical disdrometer. The data are partitioned according to diurnal, convective and stratiform precipitation classifications, monsoonal regime, and MJO phase. Findings support previous Sumatra studies suggesting a significant diurnal and DSD parameter signal associated with both convective-stratiform and MJO phase regime classification. Negligible monsoonal phase influence is determined for the DSD over the Kototabang location. The natural variability of vertical profile of DSDs leads to variability of Z-R relationships in rain column.

[1] Marzuki, Hashiguchi, H., Yamamoto, M. K., Mori, S., and Yamanaka, M. D.: Regional variability of raindrop size distribution over Indonesia, *Ann. Geophys.*, 31, 1941-1948, doi:10.5194/angeo-31-1941-2013, 2013.

[2] Marzuki, Hiroyuki Hashiguchi, Toshiaki Kozu, Toyoshi Shimomai, Yoshiaki Shibagaki, Yukihiro Takahashi, *Precipitation microstructure in different Madden-Julian Oscillation phases over Sumatra*, *Atmospheric Research*, Volume 168, 1 February 2016, Pages 121-138.

Keywords: Raindrop size distribution, Micro Rain Radar, Sumatra

DIURNAL VARIABILITIES ANALYSIS OF METEOROLOGICAL PARAMETER OF CCAM-NWP OUTPUTS AND OBSERVATION AT KOTOTABANG

*Sofiati Iis Soeratman¹, Lely Qodrita Avia¹, Nurzaman Adikusumah¹

1. Indonesian National Institute of Aeronautics and Space

The diurnal variabilities of some meteorological parameter (rainfall, temperature, relative humidity, and pressure) study, needed a model that can predict the behavior of the weather in the future by following the data history in the past. As a validation stage, outcomes such dynamic model of spatial and temporal very necessary compared with surface observational data from Automatic Weather Station (AWS) and u-v wind data from Equatorial Atmosphere Radar (EAR). The purpose of this study is analyzing the diurnal variabilities of meteorological parameter of dynamic model outputs Conformal Cubic Atmospheric Model-Numerical Weather Prediction (CCAM-NWP), and the surface observational data. The data used is the hourly output CCAM-NWP models in (12-15) February, (12-15) June 2014, and (2-14) January 2015, and surface observational data in (2-4) January 2015 as validation. The method on analyzing the output CCAM-NWP from the input models Global Forecast System (GFS-NOAA) with a resolution of the model $0.25^\circ \times 0.25^\circ$, and statistical analysis with method of T-test and F-test. Based on the results shown the diurnal variabilities and dynamic process of meteorological parameter for the model output and surface observation. Furthermore, we obtained a weak correlation between the temporal model output with surface observations from statistical analysis which mean that there are significant differences of the results found from the CCAM-NWP output and surface observations. Additionally, we compared spatial distribution of hourly output CCAM-NWP with satellite observation data such as MTSAT.

Keywords: rainfall, temperature, relative humidity, pressure, CCAM-NWP, T-test, MTSAT.

Solar-Terrestrial Interaction as Studied from Total Solar Eclipse of 9 March 2016

*Rhorom Priyatikanto¹, Tiar Dani¹, Emanuel Sungging Mumpuni¹, Mamat Ruhimat¹, Jiyo .¹

1. Indonesian National Institute of Aeronautics and Space

Total Solar Eclipse of 9 March 2016 provides a great opportunity to study the interaction between the Sun and Earth as a complex system. Several team of observers and scientists from National Institute of Aeronautics and Space will be dispatched to some observing points along the track of Lunar shadow and its proximity. The main missions are: (1) to study the physical nature of lower part of Solar corona, (2) to understand the ionospheric disturbance caused by a sudden drop of Solar radiation to the Earth, and (3) to measure the dynamics of local magnetic field that is influenced by the ionospheric disturbance. Preliminary result of those observations/experiments will be discussed in this paper.

Keywords: Solar Eclipse, Corona, Ionosphere, Geomagnet

RELATIONSHIP OF F LAYER HEIGHT RISE, ESF ONSET TIME, AND AMPLITUDE OF LARGE SCALE WAVE STRUCTURE

*Dyah Rahayu Martiningrum¹, Mamoru Yamamoto², Tatsuhiro Yokoyama³, Prayitno Abadi⁴, Pornchai Supnithi⁵, Minh Le Huy⁶

1.Graduate School of Informatics, Kyoto University, 2.Research Institute for Sustainable Humanosphere, Kyoto University, 3.National Institute of Informatics and Communications Technology (NICT), Tokyo , 4.Solar-Terrestrial Environment Laboratory (STEL), Nagoya University, 5.Faculty of Engineering King Mongkut's Institute of Technology, Ladkrabang, Thailand , 6.Institute of Geophysics Vietnamese Academy of Science and Technology, Vietnam

Under favourable conditions ionospheric plasma at the bottomside of the F-region can become unstable. Plasma irregularities in several scales produced by Rayleigh-Taylor Instability mechanism. The recombination process at the E-region after sunset causing the bottom side F-region electron density gradient become steep. At the same time the eastward electric field intensified just before it reverses to the westward electric field causing the peak of the F region rise known as evening prereversal enhancement. The relationship between onset time of field aligned irregularities related to ESF observed by Equatorial Atmosphere Radar during 2010-2012 with F layer height rise observed by ionosondes near geomagnetic equator, Chumphon (10.7 N; 99.4 E; 0.86 N) and Bac Lieu (9.3 N; 105.7 E; 0.62 N) is discussed in correspondence with growth rate of Rayleigh Taylor instability development. The field aligned irregularities onset time associated with equatorial spread F observed into two different cases, first case where the EPB/ESF onset time concurrent with the peak of virtual height of F region - ($h'F$) time. Second case where the EPB/ESF onset time happened with delay of more than 30 minutes, with reference to the peak of virtual height of F region - ($h'F$) time. The possibility that the amplitude of Large Scale Wave Structure obtained from GRBR data effects the height of plasma instability development also discussed.

Keywords: F layer height rise, Large Scale Wave Structure, onset time of Equatorial Spread F

EIA structure and its relation to plasma bubble occurrence along longitude 100° E in 2012

*Kornyanat Watthanasangmechai¹, Mamoru Yamamoto², Tatsuhiro Yokoyama¹, Michi Nishioka¹, Mamoru Ishii¹

1.National Institute of Information and Communications Technology, 2.RISH, Kyoto Univ.

To reveal a relationship between EIA structure and plasma bubble occurrence along longitude 100° E in 2012, the ground-based observations i.e., GNU Radio Beacon Receivers (GRBR), GPS, ionosonde, and Equatorial Atmospheric Radar (EAR), are employed. GRBR is used to capture the precise structure of EIA. The Rate Of TEC change Index (ROTI) of GPS is used to monitor the plasma bubble occurrence. Temporal variation of the bottom side of the F layer is measured by the ionosonde. The EAR is for detecting Field Aligned Irregularities (FAI). In this paper, we will answer the following questions:

- (1) What does the EIA structure look like on the plasma bubble on and off day?
- (2) Is it necessary that plasma bubble will occur only when EIA is symmetric?
- (3) How does the plasma bubble structure look like on the GRBR TEC?

Keywords: EIA, Plasma bubble

TBExおよびFORMOSAT-7/COSMIC-3衛星を用いた赤道電離圏研究用の新デジタルビーコン受信機
New digital beacon receiver for the study of equatorial ionosphere with satellites TBEx
and FORMOSAT-7/COSMIC-2

*山本 衛¹、岩田 桂一¹、松永 真由美²、Tsunoda Roland³、Doe Richard³、Hsiao Tung-Yuan⁴

*Mamoru Yamamoto¹、Keiichi Iwata¹、Mayumi Matsunaga²、Roland Tsunoda³、Richard Doe³、Tung-Yuan Hsiao⁴

1.京都大学生存圏研究所、2.愛媛大学理工学研究科、3.SRI International、4.Hsing Wu University

1.Research Institute for Sustainable Humanosphere, Kyoto University, 2.Graduate School of Science
and Engineering, Ehime University, 3.SRI International, 4.Hsing Wu University

We have successfully conducted observations of total-electron content (TEC) of the ionosphere using a satellite-to-ground beacon experiment. A unique dual-band (150/400MHz) digital receiver GRBR (GNU Radio Beacon Receiver) was developed for this purpose, which is based on the recent digital-signal processing technologies. The GRBR network was deployed into the southeast Asian and Pacific regions. By using beacon signals from the low-inclination satellite C/NOFS, we studied longitudinal "large-scale wave structures (LSWS)" in detail as a possible source of equatorial Spread-F (ESF) events. Now there are 2 new beacon-satellite plans. One is TBEx (Tandem Beacon Explorer), a project by SRI International, to fly a constellation of two 3U cubesats with tri-band beacon transmitters. Another one is a constellation of FORMOSAT-7/COSMIC-2 satellites, also with tri-band (or quad-band) beacon transmitters. All of these satellites will be placed into low-inclination orbits by the same launch vehicle in late 2016. This launch will provide great opportunities to enhance studies of the low-latitude ionosphere. Kyoto University, Ehime University and Hsing Wu University are now developing the new GRBR system that is expected to be used for the TBEx and FORMOSAT-7/COSMIC-2 beacon experiments. In the presentation we will show development of the new beacon receiver together with plan of observations.

キーワード：衛星—地上ビーコン観測、電離圏全電子数、COSMIC-2、cubesat、赤道電離圏

Keywords: Satellite-ground beacon experiment, Ionospheric TEC, COSMIC-2, cubesat, equatorial ionosphere

太陽地球結合系研究のためのIUGONETデータ解析システム

IUGONET data analysis system for a study of coupling processes in the solar-terrestrial system

*新堀 淳樹¹、田中 良昌²、梅村 宜生³、小山 幸伸⁴、阿部 修司⁶、能勢 正仁⁷、上野 悟⁵

*Atsuki Shinbori¹, Yoshimasa Tanaka², Norio Umemura³, Yukinobu Koyama⁴, Shuji Abe⁶, Masahito Nose⁷, Satoru UeNo⁵

1.京都大学生存圏研究所、2.国立極地研究所、3.名古屋大学宇宙地球環境研究所、4.新領域融合センター、5.京都大学大学院理学研究科天文台、6.九州大学 国際宇宙天気科学・教育センター、7.京都大学大学院理学研究科地磁気センター


1.Research Institute for Sustainable Humanosphere (RISH), Kyoto University, 2.National Institute of Polar Research, 3.Institute for Space-Earth Environment research (ISEE), Nagoya University, 4.Transdisciplinary Research Integration Center, 5.Hida Observatory, Kyoto University, 6.International Center for Space Weather Science and Education (ICSWSE), Kyushu University, 7.WDC for Geomagnetism, Kyoto University

Various kinds of disturbance phenomenon and long-term variation in the upper atmosphere above 100 km are caused by energy input from solar radiation, solar wind, momenta and energies from the lower atmosphere via atmospheric waves, and chemical reaction. Such a disturbance phenomenon and long-term variation observed by various kinds of ground-based and satellite instruments are the result of such complicated processes. Therefore, in order to understand the mechanisms of these phenomena detected in this region, researchers need to conduct comprehensive analyses with various kinds of long-term observation data that have been continued by means of a global network of radars, magnetometers, optical sensors, helioscopes, etc. The IUGONET (Inter-university Upper atmosphere Global Observation NETwork) project initiated in 2009 is proposed to establish a cross-reference system for various kinds of ground-based observation data obtained from different techniques. The IUGONET participants consist of five universities/institutes: the National Institute of Polar Research (NIPR), Tohoku University, Nagoya University, Kyoto University, and Kyushu University. We have developed a metadata database (MDDb) and IUGONET data analysis software (UDAS) of ground-based observation data managed by these IUGONET universities/institutes with an international collaboration in order to promote a comprehensive study on coupling processes in the Sun-Earth system. MDDb provides researchers in a wide range of disciplines with a seamless data environment to link observation databases spread across the IUGONET universities/institutes. Researchers can find basic information of observation data (for example, observation period, data format, data use policy, instrument, observed region, observation site etc.) from the metadata for each observation data they are interested in. Recently, the metadata for several satellite observation data (for example, Akebono, GPS) have been registered to MDDb, and researcher can easily find the satellite database. UDAS will also be of great help in conducting integrated analyses and visualization of various kinds of solar-terrestrial observation data to investigate the long-term variation in the upper atmosphere throughout the Sun-Earth system. UDAS deals with 29 kinds of ground-based observation data provided by the IUGONET universities/institutes. Moreover, the executable file of SPEDAS/UDAS working on Interactive Data Language (IDL)-VM without IDL license has been released on the SPEDAS website. Owing to the executable file, anybody can use UDAS and conduct a comprehensive analysis of various kinds of solar and Earth's atmospheric observation data. Moreover, in the IUGONET project, in order to enhance an international use of these IUGONET products, we held a mini-training workshop of how to use the IUGONET MDDb and UDAS at several developing countries (India, Indonesia, Nigeria, Peru etc). Now, the IUGONET products became an

essential tool to study a mechanism of the solar-terrestrial environmental change. In this talk, we introduce a brief overview of the IUGONET project, and an application of the IUGONET products to typical examples of upper atmospheric researches.

キーワード：大学間連携プロジェクト(IUGONET)、超高層大気、メタデータデータベース、解析ソフト、長期変動、地上観測

Keywords: Inter-university Upper atmosphere Global Observation NETWORK, Upper atmosphere, Metadata database, IUGONET data analysis software (UDAS), Long-term variation, Ground-based observation



Ionospheric weather in the dayside polar cap region

*Hitoshi Fujiwara¹, Satonori Nozawa², Yasunobu Ogawa³, Ryuho Kataoka³, Yasunobu Miyoshi⁴, Hidekatsu Jin⁵, Hiroyuki Shinagawa⁵, Huixin Liu⁴

1.Faculty of Science and Technology, Seikei University, 2.Institute for Space-Earth Environment Research, Nagoya University, 3.National Institute of Polar Research, 4.Department of Earth and Planetary Sciences, Faculty of Sciences, Kyushu University, 5.National Institute of Information and Communications Technology

Using European incoherent scatter radar (EISCAT) and EISCAT Svalbard radar (ESR), we have made simultaneous observations of the dayside polar cap ionosphere at Longyearbyen and Tromsø. Five year observation data show significant disturbances in the polar cap region northward the ESR site even during geomagnetically quiet periods. This suggests that the energy input from the magnetosphere would play a fundamental role for dynamics and energetics in the region all the time. In addition, some researchers suggest effects from the lower atmosphere on the thermospheric variations in the polar region. In order to understand variations of the polar ionosphere/thermosphere from hour to hour, we should understand energy flows from the above (magnetosphere) and below (lower atmosphere).

Numerical simulations with a GCM are also essential to do so. In particular, GCM simulations would reveal physical and chemical processes/mechanisms to produce the thermospheric variations while the thermospheric observations are very few in the dayside.

In the present study, we overview the EISCAT and ESR observations during 2011-2016. The recent progress of our understandings from comprehensive observations and GCM simulations are also shown.

Keywords: ionosphere, thermosphere, EISCAT, GCM, simulation

Using Vondrak smoothing method to improve the prediction of Sunspot Maximum

*ZQ Yin¹, YB Han¹

1.NAO. CAS. CHINA

About 11-year cycle of solar activity is the most significant quasi-periodic component, which largely represents the strength of solar activity and is the basis for dividing solar cycle. Many studies have proven that solar activity can affect the developing process of many phenomena on the Earth, such as climate, geophysical and marine changes etc. In order to meet the demands of heliophysics and solar activity variation characteristics studies, especially, for the demand of Solar Terrestrial Relations studies and space science development, studies of solar activity characteristics and prediction are getting more attention. Some authors summerized a variety of methods for the maximum predictions of 22nd, 23rd, 24th solar cycles, the incomplete statistics are 63, 54 and 75 cases respectively, results of the methods, which the difference between forecasting and observed values within the range of $\pm 15\%$, are 27.0%, 25.9% and 24.3% respectively. Using the 13 points smoothed value of monthly sunspot numbers, we studied correlation between sunspot number rising rate of the first 24 months of the solar cycle and the coming cycle maximum, published forecasting result that the maximum value was 139.2 ± 18.8 for 23rd solar cycle (Han et al., 2000), and the observed value is 120.8, the error is about 15.2%. The present paper introduces our improved forecasting methods. Vondrak smoothing method is used to deal with the monthly sunspot numbers. It is studied that the relationship between the rise rate of earlier months of sunspot numbers of this smoothed sequence and the coming maximum value in each solar cycles. The results show that the first 22, 23, 24 months rise rate of sunspot numbers are highly related with the coming maximum values, and simulated prediction of maximum for 22~24 cycles show that using the 22-month rise rate of three solar cycles, the maximum forecasting error is about 13.2%, using 23-month rise rate, the maximum error is about 11.2%, while using 24-month rise rate, the maximum error is only about 9.3%. The new method can make the forecasting time in advance at least half a year than the common method using 13 points monthly smoothed value, which the difference between prediction and observed values within the range of $\pm 15\%$. The results also show that the maximum error is about 20% if we use 18-month rise rate, and it can make the forecasting time in advance at least one year than the common method using 13 points monthly smoothed value.

Keywords: Sunspot Maximum, Prediction

D- and E-region ISR spectra measured with the EISCAT radars

D- and E-region ISR spectra measured with EISCAT radars

*小川 泰信¹、野澤 悟徳²、Häggström Ingemar³

*Yasunobu Ogawa¹, Satonori Nozawa², Ingemar Häggström³

1.国立極地研究所、2.名古屋大学宇宙地球環境研究所、3.EISCAT本部

1.National Institute of Polar Research, 2.Institute for Space-Earth Environmental Research, Nagoya University, 3.EISCAT Headquarters

The energy from the solar wind is mainly transported to the polar upper atmosphere and causes various phenomena such as auroras characterized by their rapid variability in time and space. Incoherent scatter radar (ISR) located in high latitude is one of the most powerful tools to investigate generation mechanisms of such phenomena and their effects on the atmosphere. The ISR basically gives information of plasma parameters between the bottom-side and topside ionospheres. However, the ISR has several unavoidable limitations to derive ionospheric parameters in the E- and D-region ionospheres, due to limited information of the ISR spectra.

We have investigated ISR spectra in the D- and E-region using the latest techniques of the EISCAT UHF and VHF radars located in Tromsø. Previous studies indicate that the EISCAT UHF and VHF radars have limitations of plasma parameter derivation below 90 and 100 km altitudes, respectively. We show lower limits of reliable plasma parameter derivation using the latest pulse coding techniques of the EISCAT radars. Based on the knowledge and experience of the current EISCAT radar observations, we will discuss possibility and limitation of future E- and D-region observations of the EISCAT_3D radar.

キーワード: ionosphere、Incoherent scatter radar、plasma

Keywords: ionosphere, Incoherent scatter radar, plasma

トロムソOMTIを用いた極域超高層大気の国内・国際共同研究

National/International joint studies with the Tromsø OMTI for studying the polar upper atmosphere

*大山 伸一郎¹、塩川 和夫¹、野澤 悟徳¹、大塚 雄一¹、三好 由純¹、Turunen Esa²、Aikio Anita³、Cai Lei³
*Shin-ichiro Oyama¹, Kazuo Shiokawa¹, Satonori Nozawa¹, Yuichi Otsuka¹, Yoshizumi Miyoshi¹, Esa Turunen², Anita Aikio³, Lei Cai³

1.名古屋大学宇宙地球環境研究所、2.オウル大学ソダンキラ地球物理学観測所、3.オウル大学

1.Institute for Space-Earth Environmental Research, Nagoya University, 2.Sodankyla Geophysical Observatory, University of Oulu, 3.University of Oulu

Institute for Space-Earth Environmental Research (ISEE) has been operating various kinds of optical instruments for more than 15 years at the Tromsø European Incoherent Scatter (EISCAT) radar site in Norway (Geographic: 69.6N, 19.2E, Geomagnetic: 66.7N, 102.2E, L=6.4), which is one of the state-of-art observatories at high latitudes. In January 2009, we began operation of a Fabry-Perot interferometer (FPI) and an all-sky type cooled CCD camera (ASC) at dark nights of the winter months (from middle of September to early April in most years). The FPI (#01 of the Optical Mesosphere Thermosphere Imagers (OMTIs)) measures airglow and aurora by rotationally changing optical wavelengths of 557.7 and 630.0 nm for most of the nights in order to derive the thermospheric wind and temperature. Four cardinal points with zenith angle of 45 or 15 and the geographic vertical (i.e. five directions) are measured sequentially. The combination of data from the four cardinal positions provides the horizontal components of the neutral wind velocity. The all-sky camera (camera #12 of OMTIs) includes a filter wheel used to select one of the six optical filters (channel 1: 557.7 nm, channel 2: 630.0 nm, channel 3: OH band (720 ~1000 nm), channel 4: 589.3 nm, channel 5: 572.5 nm, and channel 6: 732.0 nm). While the exposure time for FPI and ASC measurements is tuned as good for individual scientific objectives, in the case of FPI typical values are 1 minute or 15 seconds for airglow and auroral measurements, respectively, and in the case of ASC those are about 30 and 10 seconds for the individual targets, respectively. However, shorter exposure times have been applied for measuring fine structures of aurora. Other optical instruments such as photometer, digital camera, and all-sky camera (486.1 nm) are collocated with the OMTI FPI and ASC. Data from these instruments have been provided to national and international researching activities. For example, the FPI was operated together with the EISCAT radars as a key diagnostic instrument for providing thermospheric parameters during periods of geomagnetic disturbances. Integrated studies of the FPI-measured wind and the EISCAT-measured ionospheric parameters (including full vector of the electric field) improve our understanding of the energy dissipation and momentum transfer processes and of polar ionospheric/thermospheric responses to the solar-wind change. Effects of energetic electron precipitation (EEP) on the atmosphere are also analyzed using data from our optical instruments and EISCAT radars. This topic is proceeded under collaboration with Chemical Aeronomy in the Mesosphere and Ozone in the Stratosphere (CHAMOS) group. Sodankylä Ion Chemistry (SIC) model, which is used in the group, is a powerful tool for estimating atmospheric responses to EEP, in particular variations in the atmospheric minor components such as NO_x, HO_x, and O₃. Ionospheric and atmospheric variations associated with EEP will be one of the important scientific objectives in the ERG project. This presentation will make a brief introduction of a few results from the national/international join studies.

キーワード：オーロラ、大気光、光学装置、電離圏、熱圏、高緯度

Keywords: Aurora, Airglow, Optical instrument, Ionosphere, Thermosphere, High latitude

脈動オーロラが起こすナトリウム密度変動の研究：高速ナトリウムライダー観測

Study on pulsating aurora-induced sodium density variation: high-speed sodium lidar observation

*高橋 透¹、津田 卓雄²、細川 敬祐²、野澤 悟徳³、小川 泰信⁴、川原 琢也⁵、斎藤 徳人⁶、和田 智之⁶、川端 哲也³、Hall Chris⁷

*Toru Takahashi¹, Takuo T. Tsuda², Keisuke Hosokawa², Satonori Nozawa³, Yasunobu Ogawa⁴, Takuya Kawahara⁵, Norihito Saito⁶, Satoshi Wada⁶, Tetsuya Kawabata³, Chris Hall⁷

1.電気通信大学宇宙・電磁環境研究センター、2.電気通信大学大学院情報理工学研究所、3.名古屋大学宇宙地球環境研究所、4.国立極地研究所、5.信州大学工学部、6.理化学研究所、7.トロムソ大学

1.Center for Space Science and Radio Engineering, The University of Electro-communications, 2.Department of Communication Engineering and Informatics, University of Electro-Communications, 3.Institute for Space-Earth Environment Research, 4.National Institute of Polar Research, 5.Faculty of Engineering, Shinshu University, 6.RIKEN, 7.The Arctic University of Norway

トロムソナトリウムライダーは2010年10月に観測を開始し、2016年2月現在までの冬季に計6シーズンの観測を行っている。これまで、EISCAT観測所に設置された観測機器との同時観測データを用いて、大気重力波の上方伝搬、イオン温度と中性大気温度の比較、スプラディックナトリウム層の生成機構など数分から数時間の時間スケールの現象を取り扱ってきた。

近年、高エネルギー電子を振り込ませる脈動オーロラが注目されつつある。脈動オーロラを起こす高エネルギー電子は高度100 km以下まで侵入し、衝突を介して中性大気にエネルギーを受け渡す。このエネルギーの授受に伴う中性大気の応答過程は未だに明らかになっていない。これは、明滅周期が10秒前後である脈動オーロラに対して、十分な時間分解能で中性大気を計測するような観測が行われていないことに大きな問題があるためである。

本研究では、高出力・高感度のトロムソナトリウムライダーのデータ取得系及び、レーザー制御系に改良を加え、現システムの時間分解能を1桁以上向上させることに挑戦した。国内で開発した改良データ記録システムを2016年1月にトロムソのライダーシステムに導入し、試験的な観測をすでに開始、現在では秒/サブ秒オーダーでのナトリウム密度計測に成功している。本発表では、2015年1月～3月の試験的観測により得られたナトリウム密度の高時間分解能データの紹介を行う予定である。

キーワード：ナトリウムライダー、脈動オーロラ

Keywords: sodium lidar, pulsating aurora

南極昭和基地大型大気レーダーで得られた中間圏エコーと下部電離圏における電子密度モデルとの比較研究

Comparison study between coherent radar echo and empirically-modeled electron density in the mesosphere based on the PANSY radar

*西山 尚典^{1,2}、佐藤 薫³、中村 卓司^{1,2}、堤 雅基^{1,2}、佐藤 亨⁴、西村 耕司^{1,2}、田中 良昌^{1,2}、富川 喜弘^{1,2}、高麗 正史³

*Takanori Nishiyama^{1,2}, Kaoru Sato³, Takuji Nakamura^{1,2}, Masaki Tsutsumi^{1,2}, Toru Sato⁴, Koji Nishimura^{1,2}, Yoshimasa Tanaka^{1,2}, Yoshihiro Tomikawa^{1,2}, Masashi Kohma³

1.国立極地研究所、2.総合研究大学院大学、3.東京大学、4.京都大学

1.National Institute of Polar Research, 2.SOKENDAI (The Graduate University for Advanced Studies), 3.The University of Tokyo, 4.Kyoto University

Polar Mesosphere Winter Echo (PMWE) is known as back scatter echo from 55 to 85 km in the mesosphere, and it has been observed by MST and IS radar in polar region during non-summer period. Since density of free electrons as scatterer is low in the dark mesosphere during winter, it is suggested that PMWE requires strong ionization of neutral atmosphere associated with Energetic Particles Precipitations (EPPs) during Solar Proton Events [Kirkwood *et al.*, 2002] or during geomagnetically disturbed periods [Nishiyama *et al.*, 2015]. However, studies on relationship between occurrence of PMWE and background electron density has been limited yet [Lübken *et al.*, 2006], partly because the PMWE occurrence rate is known to be quite low (2.9%) [Zeller *et al.*, 2006].

The PANSY (Program of the Antarctic Syowa MST/IS) radar, which is the largest MST radar in Antarctica, observed many PMWE events since it has started mesosphere observations in June 2012. We established an application method of the PANSY radar as riometer, which makes it possible to estimate Cosmic Noise Absorptions (CNA) as proxy of relative variations on background electron density. In addition, electron density profiles from 60 to 150 km altitude are calculated by Ionospheric Model for the Auroral Zone (IMAZ) [McKinnell and Friedrich, 2007] and CNA estimated by the PANSY radar.

In this presentation, we would like to focus on strong PMWE during two big geomagnetic storm events, St. Patrick's Day and the Summer Solstice 2015 Event, in order to compare observed PMWE characteristics to model background electron density. On March 19 and 22, recovery phase of St. Patrick's Day Storm, sudden PMWE intensification was detected near 60 km by the PANSY radar. At the same time, strong Cosmic Noise Absorptions (CNA) of ~ 0.8 dB and 1.0 dB were measured, respectively. However, calculated electron density profiles did not necessarily demonstrate high electron density at the altitude where the PMWE intensifications were observed. On June 22, the Summer Solstice 2015 Event, strong nighttime PMWE near 80 km was detected around 16 UT, which is equal to 19 LT at Syowa station. Since PMWE observations are primarily confined to daytime because of relatively abundant free electrons in the illuminated mesosphere, this strong and long-lived nighttime PMWE implies that EPPs related to the storm caused the sporadic ionization sufficient for PMWE even in dark mesosphere. The modeled electron density profile agreed with the occurrence of the PMWE, showing density enhancement of $10^9 - 10^{10} \text{ cm}^{-3}$ from 60 to 80 km altitude. The difference between the above two events is thought to be caused by overestimations of CNA in St. Patrick's Day Storm, which leads to underestimation of modeled electron density. We are going to validate CNA estimated by the PANSY radar and discuss about discrepancy between modeled electron density and PMWE in detail.

キーワード：大型大気レーダー、中間圏、冬季極域中間圏エコー、電離圏D領域

Keywords: MST radar, Mesosphere, Polar Mesosphere Winter Echo, Ionospheric D-region

南極昭和基地SENSU SuperDARNレーダーの第IX期南極研究観測計画(2016-2023)

Antarctic Syowa station SENSU SuperDARN radars in the 9th phase of JARE Antarctic project (2016-2023)

*行松 彰^{1,2}

*Akira Sessai Yukimatu^{1,2}

1.国立極地研究所、2.総合研究大学院大学

1.National Institute of Polar Research, 2.Dept. Polar Sci., SOKENDAI (The Graduate University for Advanced Studies)

Syowa SENSU SuperDARN HF radars are important components of SuperDARN, the international HF radars network since 1995 and have significantly contributed to understanding not only magnetosphere-ionosphere coupling system but also MLT region dynamics. As SuperDARN radars were originally designed to reveal global polar ionospheric plasma convection patterns in both hemispheres in real time, its spatial resolution has been relatively low. As the number of new scientific targets like comparison with mid and small scale aurora phenomena, meso scale transient phenomena, elementary generation and decay process of field aligned irregularities, PMSEs and fine height profile of neutral wind has been increasing, higher spatial and temporal resolution observations have been essentially desired and of great importance. Imaging radar technique has been tried to be applied and developed to overcome these issues. We show the current status of the SENSU imaging radar system, and will discuss particularly on the scientific targets including coming JARE (Japanese Antarctic Research Expedition) project phase IX (2016-2023) and the future perspectives which can be revealed by this new technique using SuperDARN. In the phase IX JARE project. Our research group has proposed a research program to JARE project focused especially on possible grand minimum influences on geospace environment and inner magnetospheric dynamics and the detail plans including collaboration with other satellite and ground-based observation and its contribution to the project "Study of coupling processes in solar-terrestrial system" as well as the international VarSITI program will be discussed.

キーワード：SuperDARN、南極昭和基地、第IX期南極観測計画、グランドミニмум、内部磁気圏、VarSITI

Keywords: SuperDARN, Antarctic Syowa Station, Japanese Antarctic Research Expedition project phase 9, Grand minimum, inner magnetosphere, VarSITI