

## Orbital evolution of solid bodies in circumplanetary gas disks

\*Keiji Ohtsuki<sup>1</sup>, Shumpei Shimizu<sup>1,2</sup>, Hiroshi Kawamura<sup>1</sup>, Ryo Suetsugu<sup>1</sup>

1. Graduate School of Science, Kobe University, 2. NTT DATA Corporation

In the late stage of the formation of giant planets, sufficiently massive proto-giant planets capture gas and solids from the protoplanetary disk and form circumplanetary disks. Regular satellites of the giant planets such as the Galilean satellites of Jupiter are orbiting in the prograde direction in approximately circular and co-planer orbits, thus they are thought to be formed in the circumplanetary disks. Orbital decay of solid bodies is caused by different mechanisms depending on their sizes. When the solid bodies are small, aerodynamic gas drag is dominant (Adachi et al. 1976). Sufficiently small bodies are coupled to the gas and would be supplied to the circumplanetary disks with the inflowing gas (e.g., Canup & Ward 2002). Planetesimals that are large enough to become decoupled from the motion of the gas can be captured by gas drag from the circumplanetary gas disk (Fujita et al. 2013). It has been recently shown that the efficiency of capture of planetesimals from their heliocentric orbits by gas drag from the circumplanetary disk is the highest for planetesimals with radii of 10-100m (Tanigawa et al. 2014). While the so-called type-I migration is important in the late stage of satellite formation, orbital evolution by aerodynamic gas drag governs the orbital evolution of small solid bodies, and dynamical evolution of such small bodies in the circumplanetary gas disks is important for the growth of protosatellites. In the present work, we examine orbital evolution of planetesimals in circumplanetary gas disks, and the probability of capture of such small bodies by a growing protosatellite (Shimizu & Ohtsuki, in preparation).

We numerically evaluate the probability of collision of migrating planetesimals with the protosatellite, and its dependence on the size of planetesimals. We find that the collision probability has a peak at a certain size. This is because the time scale of the orbital decay varies depending on the size of planetesimals. We also examined cases of various masses and semi-major axes of the protosatellite, and obtained similar results. Finally, we will also discuss effects of gravitational interaction between planetesimals (Kawamura, Ohtsuki, Suetsugu, this meeting).

Keywords: Satellite formation

## Insulator to electronic conductor transition of synthetic planetary ices at interior conditions of icy giants compressed by laser-driven shock wave


\*Takuo Okuchi<sup>1</sup>, Norimasa Ozaki<sup>2</sup>, Mika Kita<sup>2</sup>, Tsuyoshi Ogawa<sup>2</sup>, Albertazzi Bruno<sup>2</sup>, Kohei Miyanishi<sup>2</sup>, Ryosuke Kodama<sup>2</sup>, Narangoo Purevjav<sup>1</sup>, Takayoshi Sano<sup>2</sup>, Youichi Sakawa<sup>2</sup>

1.Okayama Univ., 2.Osaka Univ.

Ices in the universe are consisting of hydrogen, oxygen, carbon, and nitrogen. These elements coalesced to develop into icy giant planets such as Uranus and Neptune in the solar system, as well as some water planets occurring in extrasolar planetary systems. Properties of such planetary ices at elevated pressures and temperatures are essential clues for understanding the layering structures and material circulations inside these giants. Here by using high-power nanosecond laser pulses from GEKKO-XII glass laser at Institute of Laser Engineering at Osaka University, such ices of several compositional types are dynamically compressed, where their equation-of-state (P-V-T) and optical reflectivity (R) are measured at in situ conditions using fast optical diagnostics (Figure). We have successfully observed the transition from insulator to electronic conductor state of the planetary ices which include carbon and/or nitrogen components, as well as that transition of pure H<sub>2</sub>O ice observed by our previous work [1]. We have also observed the transition at off-Hugoniot conditions using a sample-precompression technique with a diamond anvil cell, which was effectively coupled with laser-driven shock to increase the shock-pressure and to decrease the shock-temperature. These results provide us a new insight about the nature of conducting media inside the icy giants and about the origin of magnetic fields from deep inside of them.

[1] T. Kimura, N. Ozaki, T. Sano, T. Okuchi, et al., J. Chem. Phys., 142, 164504 (2015).

Keywords: icy giant planets, laser-driven shock compression, off-Hugoniot conditions, planetary magnetic field



## A massive primordial-atmosphere on proto-Titan formed in a gas-starved disk

\*Takashi Mikami<sup>1</sup>, Yasuto TAKAHASHI<sup>1</sup>, Kiyoshi Kuramoto<sup>1,2</sup>

1.Department of CosmoSciences, Graduate School of Science, Hokkaido University, 2.Center for Planetary Sciences, Kobe

Titan is known as a satellite with a thick atmosphere (1.5 bar at the surface) mainly composed of nitrogen. Although several hypotheses about the origin of atmosphere have been proposed, it remains an open question how and when such a thick atmosphere was generated. According to the recent satellite formation theory [e.g., Canup and Ward 2002], Titan formed within low temperature and pressure disk. We numerically investigate the property of the primordial atmosphere of Titan that grew in such a circum-planetary disk, especially in terms of the atmospheric mass and the blanketing effect. In spite of such a disk condition, Titan could capture a thick atmosphere strongly bounded by gravity, which is mainly composed of nebula gas components. This would cause a significant blanketing effect inducing differentiation of this satellite, and result in keeping the surface temperature high relatively (~200 K). This suggests that an ammonia-rich proto atmosphere could be kept on Titan even after the disk was dissipated. Titan's current nitrogen would be generated from ammonia in the proto atmosphere by photochemical reaction [Atreya et al., 1978]

Keywords: Titan, Atmosphere

## Predicting undiscovered species in Titan's stratosphere with chemical reaction network based on UMIST database

Chen Shen<sup>1</sup>, \*Hideko Nomura<sup>1</sup>, Jun Kimura<sup>2</sup>

1.Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2.Earth-Life Science Institute, Tokyo Institute of Technology

Titan, the largest moon of Saturn, is the only satellite that has a dense atmosphere in the solar system. It is known that Titan's atmosphere contains a wide variety of chemical species which mainly generated from the dissociation of two main components, molecular nitrogen and methane. Fractional abundances of these species have been studied well by the Voyager and Cassini-Huygens probes. Also, ALMA now starts to detect global distribution of some species in the Titan's atmosphere. In this study, we applied chemical reaction network based on UMIST database, which has been used in the studies of interstellar medium, to calculate the chemical evolution of Titan's upper stratosphere around 200 km from the satellite surface, where most of the observation data of molecular abundances heretofore located. In this chemical network calculation, 375 species are included, which are three times more than previous studies (Wilson et al. 2004 & Loison et al. 2015). We note that the effects of turbulent diffusion and three-body reactions are not included in the calculation. By comparing results of calculations with the observational data, a physical parameter set with moderate FUV flux, effect of cosmic ray and self-shielding of molecular nitrogen and methane is recommended. As a result, 17 nitrogen compounds (e.g. NH<sub>2</sub>CN CH<sub>3</sub>C<sub>5</sub>N HC<sub>7</sub>N) are abundant and could be detectable in the future observations of ALMA.

Keywords: Titan's atmosphere, chemical reaction network, ALMA observations

## Stability of subsurface ocean in Pluto

\*Jun Kimura<sup>1</sup>, Shunichi Kamata<sup>2</sup>

1.Earth-Life Science Institute, Tokyo Institute of Technology, 2.Hokkaido university

NASA's New Horizons spacecraft made its close flyby of the dwarf planet Pluto on July 14, 2015. The LORRI imaging system aboard spacecraft has acquired global images and unveiled a diverse range of landforms, from rugged mountainous region to extremely smooth plains, indicating geological processes that have substantially and recently modified the surface. Combining that accurate determinations of the mean radii of Pluto suggests that Pluto is a sphere, Pluto had or has a relatively warm interior (maybe an ocean) for the most part of its history.

Nitrogen (N<sub>2</sub>) ice, higher volatility (melting point of 63 K in vacuum) than water ice, has been known from ground-based spectroscopy and the New Horizons has confirmed that N<sub>2</sub> (and CH<sub>4</sub>) ice is enriched in the bright smooth plains, e.g., Sputnik Planum (SP). The ALICE, ultraviolet imaging spectrograph, has observed Pluto's nitrogen-rich atmosphere as far as 1,600 kilometers above the surface of the planet, demonstrating that nitrogen is volumetrically dominant on Pluto. In parallel, water ice is widely distributed on Pluto, in particular, on rugged mountainous region having relatively older age than SP. It implies that Pluto is covered by huge amount of water ice and few-km thickness nitrogen presents above water ice "bedrock" based on the molecular abundances in the Solar System. High volatility of nitrogen ice can lead to melting and rapid thinning of the ice shell (and forming an internal nitrogen ocean) and can drive tectonics on Pluto even if radiogenic heating is expected to be rather low at present and tidal heating is not efficient.

Here we are going to show the results of numerical simulation for the thermal history in Pluto, and this dwarf planet far away from the Sun has a potential to harbor an ocean and thin solid nitrogen crust might be able to support 2-3 km height ragged mountains even few-km thickness.

Keywords: Pluto, Subsurface ocean, Interior, Astrobiology

## The Effect of Surface Ice and Topography on the Atmospheric Circulation and Distribution of Nitrogen Ice on Pluto

\*Scot CR Rafkin<sup>1</sup>, Alejandro Soto<sup>1</sup>, Timothy Michaels<sup>2</sup>

1.Southwest Research Institute Boulder, 2.SETI Institute

A newly developed general circulation model (GCM) for Pluto is used to investigate the unexpected and highly heterogeneous distribution of nitrogen surface ice imaged by the New Horizons spacecraft on the surface of Pluto. The GCM is based on the GFDL Flexible Modeling System (FMS) dynamical core, solved on a discretized latitude/longitude horizontal grid and a pressure-based hybrid vertical coordinate. Model physics include a 3-band radiative scheme, molecular thermal conduction within the atmosphere, subsurface thermal conduction, and a nitrogen volatile cycle. The radiative-convective model takes into account the 2.3, 3.3 and 7.8 mm bands of CH<sub>4</sub>, including non-local thermodynamic equilibrium effects. The subsurface conduction model assumes a water ice regolith. In the case of nitrogen surface ice deposition, additional super-surface layers are added above the water ice regolith to properly account for conductive energy flow through the nitrogen ice. The nitrogen volatile cycle is based on a vapor pressure equilibrium assumption between the atmosphere and surface. Prior to the arrival of the New Horizons spacecraft, the expectation was that the volatile surface ice distribution on the surface of Pluto would be strongly controlled by the latitudinal temperature gradient resulting primarily from the slow seasonal variations of radiative-convective equilibrium. If this were the case, then Pluto would have broad latitudinal bands of both ice covered surface and ice free surface, as dictated by the season. Furthermore, the circulation, and thus the transport of volatiles, was thought to be driven almost exclusively by sublimation and deposition flows (so-called "condensation flows") associated with the volatile cycle. In contrast to expectations, images from New Horizon showed an extremely complex, heterogeneous distribution of surface ices draped over topography of substantial geologic diversity. To maintain such an ice distribution, the atmospheric circulation and volatile transport must be more complex than previously envisioned. Topography, the distribution of nitrogen ice itself, and an overall large-scale atmospheric circulation at least partially independent of the condensation flows must play a role. Simulations where topography, surface ice distributions, and volatile cycle physics are added individually and in various combinations are used to individually quantify the importance of the general circulation, topography, surface ice distributions and condensation flows. It is shown that even regional patches of ice or large craters, much like that of Tombaugh Regio, can have global impacts on the atmospheric circulation, the volatile cycle, and hence, the distribution of surface ices. This work demonstrates that explaining Pluto's volatile cycle and the expression of that cycle in the surface ice distribution requires consideration of atmospheric processes beyond simple vapor pressure equilibrium arguments.

Keywords: Pluto, Atmosphere, Dynamics

## Neutral pH of water on early Ceres

\*Yasuhito Sekine<sup>1</sup>, Takazo Shibuya<sup>2</sup>, Taro Funatsu<sup>1</sup>

1.Department of Earth and Planetary Science, University of Tokyo, 2.Laboratory of Ocean-Earth Life Evolution Research, JAMSTEC

Ceres, the ice-rich dwarf planet in the asteroid belt, would provide a clue to understand formation processes of the planets in the solar system, as it is considered as one of a few proto-planets remaining today (Castillo-Rogez and McCord, 2010). Ceres' surface reflectance spectra show a unique absorption at 3.06  $\mu\text{m}$ , which is recently found to be caused by the presence of  $\text{NH}_4$ -bearing hydrated silicates (e.g., mica) (De Sanctis et al., 2015). This in turn means that a large amount of  $\text{NH}_3$  should have been contained in Ceres' interior ocean formed in the early stage of its evolution, and that Ceres' building materials would have been originated from the outer solar system beyond the snowline of  $\text{NH}_3$  (De Sanctis et al., 2015). However, the formation of  $\text{NH}_4$ -bearing hydrated silicates would depend on not only the presence of  $\text{NH}_3$  in the ocean but also the chemical compositions and pH of the interior ocean where the hydrated silicates were formed. Here, we performed hydrothermal experiments to constrain pH of the water on early Ceres. Based on the chemical analysis and comparisons of infrared spectra of the produced hydrated silicates, together with the findings of carbonates on Ceres, we show that pH of water on early Ceres should be near neutral. This is because  $\text{NH}_4^+$  ions are incorporated into hydrated silicates under neutral pH conditions. To achieve neutral pH in the water, the rock compositions of Ceres would be different from that of carbonaceous chondrites. As sulfate salts were found on Ceres (Nathues et al., 2015), large amounts of sulfate ions may have worked as a major anion to keep the water pH as neutral. This further suggests that reducing sulfur in the core would have been oxidized by igneous activity on early Ceres sustained by short-lived radiogenic heating upon its early formation (within 3-5 Mys after CAIs).

Keywords: Ceres, hydrothermal reactions, water

## Exploration of Extraterrestrial Ocean Worlds –results from Cassini and perspectives

\*Christophe Sotin<sup>1</sup>

1. Jet Propulsion Laboratory - California Institute of Technology, Pasadena, CA (USA)

The Cassini mission, which has been in Saturn's system for more than a decade, has discovered two ocean worlds: Enceladus, a tiny moon 252 km in radius and Titan, a large moon 2575 km in radius which is the only moon in the solar system with a dense atmosphere. For each of these satellites of Saturn, the ocean is located beneath a thick icy crust. Enceladus is characterized by the presence of jets that eject samples of the deep ocean into space. The data gathered by the Cassini instruments suggest the presence of hydrothermal activity (Hsu et al., 2015). Recent measurements of Enceladus' librations are best explained by a decoupling between the icy crust and the inner core, suggesting the presence of a global ocean. Maintaining a global ocean during billions of years is a challenge for thermal evolution models of a tiny moon in a very cold environment. Titan is characterized by its dense atmosphere where large organic molecules are produced. These heavy molecules eventually form the organic haze that falls on Titan's surface and may constitute the sand of the equatorial dunes. Titan is also characterized by hydrocarbon seas at the North Pole making Titan, the only other object besides Earth with stable liquid bodies on its surface. Although the organic cycle has been studied by the Cassini missions, there are still major scientific questions such as the molecular and elementary composition of the heavy organics and the processes responsible for the replenishment in methane of Titan's atmosphere.

Enceladus can be compared with Europa in the sense that the global ocean is in contact with a rocky core where conditions are very similar to those existing on the terrestrial sea-floor. The fact that its ocean is being ejected into space makes measurements into the plume a science priority for future missions to Enceladus. Europa is the target of a planned NASA flagship mission for which the instruments to be mounted on the spacecraft have already been selected. This mission would characterize Europa's deep ocean. Although the presence of a plume (Roth et al., 2013) that would eject samples of Europa ocean into space is still debated, assets that would analyze this potential plume are being studied.

Titan can be compared to Callisto and Ganymede, Jupiter's largest moons, which have very similar mass and radius. Interpretation of the magnetic field measurements at Ganymede and Callisto by the Galileo mission suggests that they also have a deep ocean. Similarly, interpretation of (i) the electric signal recorded by the Huygens probe during its descent in Titan's atmosphere in January 2005 and (ii) the value of the tidal Love number  $k_2$ , suggest that an ocean is present beneath Titan's icy crust. On these three large moons, models based on the equation of state of H<sub>2</sub>O and values of the moment of inertia, a measure of mass repartition, suggest that this deep ocean would not be in contact with silicates because a high-pressure layer of ice would be present between the ocean and the rocky core. This paper will investigate the conditions under which the liquid water may have been in contact with the silicates in the past. This question is important to assess the habitability potential of these large icy moons. Ganymede and to a lesser degree Callisto will be the targets of the ESA JUICE mission, the first large mission of the ESA "Cosmic Vision" program. The primary goal of the JUICE mission is to characterize the oceans of Ganymede and Callisto. This work has been performed at the Jet Propulsion Laboratory, California Institute of Technology, under contract to NASA.

Keywords: Ocean Worlds, Enceladus, Titan, Europa





## The Science Case for the Ganymede Laser Altimeter (GALA) on ESA's Jupiter Icy Moons Explorer (JUICE)

\*Hussmann Hauke<sup>1</sup>, G. Steinbruegge<sup>1</sup>, A. Stark<sup>1</sup>, J. Oberst<sup>1</sup>, K. Lingenauber<sup>1</sup>, J. Kimura<sup>2</sup>, K. Enya<sup>3</sup>, N. Namiki<sup>4</sup>, M. Kobayashi<sup>5</sup>, K. Touhara<sup>3</sup>, N. Thomas<sup>6</sup>, K. Seiferlin<sup>6</sup>, L. Lara<sup>7</sup>

1.DLR Institute of Planetary Research, 2.ELSI, Tokyo, 3.ISAS/JAXA, 4.NAOJ, 5.Chiba Inst. Technology, 6.University of Bern, 7.IAA-CSIC

The Ganymede Laser Altimeter (GALA) is one of ten selected instruments for ESA's JUICE mission. Here we will give an overview on the scientific objectives for Europa, Callisto, and in particular, Ganymede to be accomplished by GALA. By obtaining range measurements during flybys at the moons and in orbit around Ganymede, geodetic and geophysical objectives will be addressed. Prime objectives at Ganymede are (1) determining the satellite's topography on global, regional, and local scales and (2) measuring tidal surface deformations. The topography measurements will provide information on the body's shape in relation to hydrostatic equilibrium. It will characterize topographic depressions and elevated regions for impact basins and other large scale features. Local topography will give insight into formation mechanisms for geologic features, e.g. the grooved terrain, cryo-volcanic sites, and impact structures. By obtaining range measurements at different tidal phases along Ganymede's orbit, the radial surface displacement can be measured. The tidal amplitudes are crucial to confirm a subsurface ocean on Ganymede. In addition the tidal phase-lag, if it can be detected, can provide key information on the deep interior of the satellite and the global dissipation inside Ganymede. Geodetic measurements will focus on the rotation state of Ganymede including the orientation of the pole and possible physical librations in longitude. By analyzing the spreading of the return pulse, the surface roughness on the scale of the laser footprint, i.e. approximately 50 m at the nominal 500-km orbit, can be determined. The slopes between individual laser spots along the track provide further information on surface roughness on larger scales, possibly correlated with geological features. The albedo at the laser wavelength of 1064 nm is obtained from each shot. Possible correlations with topographic heights and specific surface features can provide information on geological processes and on the interaction of the surface with the moon's radiation and particle environment. On Callisto, the measurements will be focused on a global shape determination to test whether this satellite which is only partially differentiated, is in a hydrostatic state. In addition impact structures of different sizes and types will be characterized by high-frequency along-track measurements. On Europa, the ground-tracks will focus on recently active sites and chaos regions. Combined with subsurface radar measurements and imaging data, this will provide insight in the formation mechanism of these unique features on Europa that are probably related to liquid water reservoirs near the subsurface. Constraints on the global shape of Europa will be provided by the flyby laser profiles with high spatial resolution. In addition surface roughness and reflectivity will be measured during the Callisto and Europa flybys.

Keywords: Jupiter, Ganymede, Laser Altimetry

## NASA's Flagship Mission to Jupiter's Moon Europa: Exploring a Potentially Habitable World

\*Louise M Prockter<sup>1</sup>, Robert T. Pappalardo<sup>2</sup>, David A. Senske<sup>2</sup>, Brian G. Paczkowski<sup>2</sup>, Rachel L. Klima<sup>1</sup>, Kathleen L. Craft<sup>1</sup>, Steven D. Vance<sup>2</sup>, Barry G. Goldstein<sup>2</sup>, Thomas J. Wagner<sup>1</sup>, Brian C. Cooke<sup>2</sup>

1.Johns Hopkins University Applied Physics Laboratory, 2.Jet Propulsion Laboratory

It is expected that Jupiter's moon Europa may have the three ingredients thought necessary for life to exist: liquid water, chemical elements from which to build organic molecules, and chemical energy. Europa is hypothesized to have the three ingredients in the form of: (1) an extensive saltwater ocean beneath an ice shell that is geodynamically active and relatively thin (several kilometers to several tens of kilometers thick); (2) key chemical elements derived from the primordial chondritic composition of the Jovian protoplanetary disk, plus delivery by asteroids and comets over time; and (3) a source of chemical energy for life, from the combination of irradiation of its surface to produce oxidants, plus hydrothermal activity and/or serpentinization at its ocean floor to produce reductants.

NASA recently approved the development of a flagship-level mission to explore Europa, with the specific goal of investigating its habitability. The spacecraft will launch some time in the next decade, and will arrive in the Jupiter system between 3 and 7 years later, depending on which launch vehicle and trajectory is selected. In order to survive the harsh Jovian radiation environment, the spacecraft will orbit Jupiter, dipping in and out of its radiation belts, and will encounter Europa at different positions in its orbit for a total of over 40 close flybys. This strategy allows data to be acquired from across most of the moon's surface and enables particles and fields measurements to be made in the local vicinity.

High-priority science will be accomplished through interrogations of the moon's ice shell, ocean, composition, geology, and current activity. The payload consists of five remote sensing instruments that cover the wavelength range from ultraviolet through radar and four *in-situ* instruments that measure fields and particles; moreover, gravity science can be achieved via the telecom system, and valuable scientific data could come from the spacecraft's planned engineering radiation monitoring system. The remote sensing instruments are: an ultraviolet spectrograph (Europa-UVS); a wide-angle and narrow-angle visible camera system (EIS); an infrared spectrometer (MISE); a thermal instrument (E-THEMIS); and an ice-penetrating radar (REASON). The *in-situ* instruments are: a magnetometer suite (ICEMAG); a plasma instrument (PIMS); a time-of-flight mass spectrometer (MASPEX); and a dust analyzer (SUDA). Taken together, the payload has the potential to test hypotheses and make discoveries relevant to the composition, interior, and geology of Europa, in order to address the potential habitability of this intriguing moon.

Keywords: Europa , Mission, Habitability, Spacecraft

## Deep Plume Interaction with Gas Giant Weather Layers: Preparing for Juno and Cassini Observations

\*Morgan E O'Neill<sup>1</sup>, Yohai Kaspi<sup>1</sup>, Eli Galanti<sup>1</sup>

1.Weizmann Institute of Science

The weather layers of Jupiter and Saturn receive both solar radiation and heat from the deep interior. Currently, most numerical models fall into two broad categories: Deep, dry and convecting interior models that lack a stably stratified troposphere above, or thin shells that represent only a troposphere, with parameterized heating from the lower boundary. Here we present the results from a new coupled model that allows resolved deep convective plumes to interact with a stable "weather layer". We demonstrate the relative importance of a stable tropospheric lapse rate and the magnitude of bottom heating on the strength and depth of the jets. Studies of this kind will benefit understanding of Jupiter's dynamics, in particular the depth of the cloud-level jets, in advance of Juno's 2016 arrival. Moreover, the difference between Saturn and Jupiter is explored using a parameter sweep of tropospheric stability, which acts as a proxy for tropospheric water abundance.

Keywords: Jupiter, Juno, Atmospheric Dynamics

Magnetic Field Observations; at Saturn with CASSINI and at Jupiter with JUICE.

\*Michele Karen Dougherty<sup>1</sup>

1.Imperial College London

Some of the highlights of discoveries made utilising magnetic field observations from the Cassini spacecraft at Saturn will be described. These include the planetary period oscillations that fill the Saturnian magnetosphere, that change over time, and that are different between the northern and southern hemisphere. The need to understand these observations is critical for Cassini end of mission science during which time the spacecraft will have 22 close flybys to Saturn, enabling resolution of the planetary dynamo field as well as how long a day is on Saturn. The discovery by the magnetometer team of a dynamic south polar plume on Enceladus, one of Saturn's icy moons, will also be described. We will also look to the future and the required measurements the magnetometer, onboard the ESA JUICE mission to Jupiter and its moons, will need to make. The most difficult measurement as well as the most interesting is that of the induced magnetic field signatures in a liquid subsurface ocean at Ganymede. Resolving these signatures at more than one inducing frequency will allow unambiguous determination of both the depth and conductivity of the ocean and potentially of its global extent.

Keywords: Jupiter, Saturn, Magnetic field, CASSINI, JUICE

Synergetic Multi-Wavelength Observation of Jupiter's Magnetosphere Driven by Hisaki:  
Recent Results and Plans for JUNO Mission

\*Tomoki Kimura<sup>1</sup>, Kazuo Yoshioka<sup>2</sup>, Go Murakami<sup>2</sup>, Atsushi Yamazaki<sup>2</sup>, Fuminori Tsuchiya<sup>3</sup>, Chihiro Tao<sup>4</sup>,  
Masaki Fujimoto<sup>2</sup>, Hisaki Science team

1.Nishina-Center for Accelerator Based Science, RIKEN, 2.JAXA/ISAS, 3.Tohoku University, 4.NICT

JAXA Hisaki satellite is an EUV space telescope dedicated for continuous monitoring of planetary atmospheric and plasma environments. Synergetic multi-wavelength observing campaigns for Jupiter's magnetosphere have been carried out by Hisaki with other ground-based and space telescopes from 2014 to the present. Here we report some highlights of the synergetic campaign and present plans for the coordinated observation with NASA JUNO mission in 2016-2017.

Keywords: Hisaki, Jupiter's magnetosphere, JUNO

## Jupiter's auroral observations by Hisaki/EXCEED and expectation toward collaborations with Juno

\*Chihiro Tao<sup>1</sup>, Tomoki Kimura<sup>2</sup>, Go Murakami<sup>3</sup>, Kazuo Yoshioka<sup>4</sup>, Fuminori Tsuchiya<sup>5</sup>, Hajime Kita<sup>5</sup>, Nicolas André<sup>6</sup>, Sarah Badman<sup>7</sup>, Michel Blanc<sup>6</sup>, Ichiro Yoshikawa<sup>4</sup>, Atsushi Yamazaki<sup>3</sup>, Daikou Shiota<sup>8</sup>, Hiroyasu Tadokoro<sup>9</sup>, Yasumasa Kasaba<sup>5</sup>, Masaki Fujimoto<sup>3</sup>

1.National Institute of Information and Communications Technology, 2.RIKEN, 3.ISAS/JAXA, 4.Tokyo University, 5.Tohoku University, 6.IRAP, 7.Lancaster University, 8.Nagoya University, 9.Musashino University

Ultraviolet (UV) emissions from atmospheric H<sub>2</sub> and H reflect powerful polar energetics at outer planets. UV spectra provides information related with the precipitating auroral electron energy. Auroral electron energy and flux relationship shows variety among Jupiter's auroral regions. The spectrometer EXCEED onboard JAXA's Earth-orbiting planetary telescope Hisaki monitors extreme UV emissions from Jovian aurora and Io plasma torus continuously. Hisaki succeeded to detect sporadic, large auroral power enhancements lasting both short- (<1 planetary rotation) and long-term (>a few rotations) variations and their modulations by an Io's volcanic activity over several weeks. The spectral information taken by Hisaki enables us to investigate (1) the time variation of the auroral electron during these emission enhancements, (2) statistical survey for occurrence of polar-dominant events, and (3) associated magnetospheric dynamics for these emission enhancement events using the Knight's aurora acceleration theory. Expecting collaborative observation with Juno will be discussed.

Keywords: Jupiter, Aurora, EXCEED/Hisaki

## Io's volcanic influence on the Jovian magnetosphere: HISAKI observation

\*Fuminori Tsuchiya<sup>1</sup>, Kazuo Yoshioka<sup>2</sup>, Tomoki Kimura<sup>3</sup>, Go Murakami<sup>4</sup>, Mizuki Yoneda<sup>5</sup>, Ryoichi Koga<sup>1</sup>, Masato Kagitani<sup>1</sup>, Hiromasa Nozawa<sup>6</sup>, Chihiro Tao<sup>7</sup>, Hiroaki Misawa<sup>1</sup>, Takeshi Sakanoi<sup>1</sup>, Yasumasa Kasaba<sup>8</sup>, Atsushi Yamazaki<sup>4</sup>, Ichiro Yoshikawa<sup>9</sup>

1.Planetary Plasma and Atmospheric Research Center, Tohoku University, 2.Graduate School of Science, The University of Tokyo, 3.RIKEN, 4.ISAS/JAXA, 5.Kiepenheuer Institute for Solar physics, 6.Kagoshima National College of Technology, 7.NICT, 8.Graduate School of Science, Tohoku University, 9.Graduate School of Frontier Sciences, The University of Tokyo

The satellite Io which has many active volcanos supplies volcanic gases to the Jovian magnetosphere with a typical rate of 1 ton/sec and has is a primary source of plasmas in the magnetosphere. Change in the volcanic activity would cause change in plasma supply rate to the magnetosphere and could affect structure of the magnetosphere and dynamics occurring in it. However, responses of the magnetosphere to the plasma supply rate change is still not fully understood. The extreme ultraviolet (EUV) spectroscopy, EXCEED, onboard the HISAKI satellite made observations of Io plasma torus and Jovian northern aurora from the end of Nov. 2014 to middle of May 2015 continuously. On middle of Jan. 2015, HISAKI detected gradual increase in intensity of  $S^+$  emission lines and decrease of  $S^{3+}$  ones in the plasma torus. The  $S^+$  intensity showed a maximum around the end of Feb. and  $S^{2+}$  and  $S^{3+}$  intensities also showed maxima subsequently. Simultaneous ground based observation of the sodium nebula showed increase of the emission intensity from the middle of Jan. to Feb. These observations suggest that the volcanic activity enhancement started at the middle of Jan. and increase neutral atom and ion densities in the Io torus. Change in radial structure of the plasma torus was also detected. The intensity of  $S^+$  ion began to increase around the orbit of Io (6 Jovian radii). The brightened region propagated outward and reached at 8.5 Jovian radii from Jupiter at the beginning of Feb. Further one month later, HISAKI found unusual activity of Jovian EUV aurora intensity. It began at the beginning of Mar. and continued for 1 month. The enhancement of the aurora activity may be caused by the enhanced loading of heavy ion plasma into the middle magnetosphere.



Radial distribution of sulfur and oxygen ions in the Io plasma torus observed by Hisaki spacecraft

\*Kazuo Yoshioka<sup>1</sup>, Fuminori Tsuchiya<sup>2</sup>, Tomoki Kimura<sup>3</sup>, Go Murakami<sup>4</sup>, Ichiro Yoshikawa<sup>1</sup>, Atsushi Yamazaki<sup>4</sup>, Yasumasa Kasaba<sup>2</sup>, Masaki Fujimoto<sup>4</sup>

1.The University of Tokyo, 2.Tohoku University, 3.RIKEN, 4.ISAS/JAXA

The imaging spectrum of Io plasma torus in extreme ultraviolet (50-150nm) has been observed by EXCEED on Hisaki. We analyzed them using the spectrum diagnosis method and deduced the column averaged plasma parameters (such as S<sup>+</sup>, S<sup>++</sup>, S<sup>+++</sup>, O<sup>+</sup>, etc.) from the radial distance of 6 to 8 Jovian radii. The local densities of those ions are deduced and the clear increase of S<sup>+</sup> (about 3 times) after the Io's outburst event in 2015 are seen.

Keywords: Io plasma torus, sulfur, oxygen

## Slope and roughness characteristics derived from high-resolution images of Galilean satellites

\*Shoko Oshigami<sup>1</sup>, Naoyuki Hirata<sup>2</sup>, Jun Kimura<sup>3</sup>, Noriyuki Namiki<sup>1</sup>

1.National Astronomical Observatory of Japan, 2.Kobe University, 3.Tokyo Institute of Technology

Topographic data is fundamental information to investigate geology at various scales. The Galileo spacecraft has obtained high-resolution images of the Galilean satellites at a scale up to approximately ten meters per pixel, which provide an insight into diverse geologies and surface materials associated with tectonics, cratering, and sublimation. Putative subsurface oceans on Europa, Ganymede, and Callisto indicated by inductive magnetic fields and aurorae shifting are top priority for the next planetary exploration. Measuring topography at a scale of ten-meter also could be essential for designing radar sounder, laser altimeter, and lander. These instruments are expected to detect putative subsurface oceans on Jovian icy satellites. Nevertheless, no quantitative topographic data at the scale has yet been obtained, except for the surface of Europa. This is mainly consequence of previous digital elevation models (DEMs) with spatial resolutions higher than 50 m being only available for the southeast region of the Tyre crater of Europa (33 m/pixel). Current knowledge of topographic features of the Galilean satellites is only derived from stereo image (SI) analysis or photoclinometry (PC) because of the absence of laser altimetry data. The slopes and roughness strongly depend on the spatial resolution of topographic data. In general, higher spatial resolution provides steeper slope histograms. We reexamined high-resolution images obtained by the SSI camera onboard the Galileo spacecraft using SI analysis and PC.

As for SI analysis, we used Integrated Software for Imagers and Spectrometers (ISIS3) produced by USGS to calibrate the SSI raw images radiometrically and perform bundle adjustment. Then we applied NASA's Ames Stereo Pipeline software (ASP) to compute DEMs. ASP is a suite of automated SI analysis tools developed by NASA and designed for processing planetary images. To compute the slope, we i) selected a pixel from a DEM, ii) computed a least squares plane in a seven-pixel square centered around the selected pixel, iii) obtained the gradient of the least squares plane (the slope is defined as the gradient), and iv) performed (i) to (iii) over the entire DEM. Then, the total surface area of each DEM was normalized to 1. To compute roughness, we defined the roughness as the Allan deviation of differences in height (i.e. RMS deviation). In detail, we i) measured the difference in height between two points separated by a given distance (i.e. window length) along lines of constant longitude, ii) performed (i) over the entire DEM and collected differences in height over the entire DEM, and (iii) computed the Allan deviation of all differences in height obtained in (ii).

PC uses apparent brightness to estimated slope at each pixel, assuming a photometric function. We applied a photometric function that fits lunar-like surface. In order to compute roughness, we converted the slopes into height differences for each pixel. Then we integrated the height differences along the line of sight to construct one-dimensional topographic cross-section from line to line. Here roughness is defined as the Allan deviation of differences in height between two points separated by a given distance along the line of sight. We showed an average of the Allan deviation values for each window length among the all one-dimensional topographic profiles.

As a result, we obtained the slope histograms and roughness from SI analysis and PC. We found that most of Ganymede and the region of Callisto showing abundant knobs appear to be very rough surfaces as steep as  $10^{\circ}$  to  $30^{\circ}$ , while Europa and the region of Callisto without knobs mostly appear to be smoother than  $10^{\circ}$ . These results are far from previous estimates based on topographic data with the lower resolution. Also, this implies that instrument performances are expected to be strongly

affected by the steep slopes in the former areas.

## Stability of subsurface ocean in Ganymede

\*Jun Kimura<sup>1</sup>, Shunichi Kamata<sup>2</sup>, Steven Vance<sup>3</sup>, Hauke Hussmann<sup>4</sup>

1.Earth-Life Science Institute, Tokyo Institute of Technology, 2.Hokkaido University,  
3.JPL/Caltech, 4.DLR

The outer solar system provides potential habitats for extra-terrestrial life. Past spacecraft's and telescopic observations support that the Jovian icy moons may harbor water oceans beneath the icy crusts. However evidence for oceans is not definitive and awaits confirmation measurements. Also their depth and composition remain unclear, as do their stability and variability with time. Here we focus on Ganymede, the largest moon in the Solar System and the primary target the Jupiter Icy Moons Explorer (JUICE). To investigate the stability of an ocean (structural, thermal and compositional change through time) assumed to be initially in an entirely liquid state, we performed numerical simulations for the internal thermal evolution using an one-dimensional spherically symmetric model for the convective and conductive heat transfer, with radial dependence of viscosity, heat source distribution, and other material properties. We take into account the energy due to decay of long-lived radioactive elements and also evaluate the effect of tidal heating. To see the temporal change of the boundary position between solid ice layers including ice shell and high-pressure ice mantle, we also evaluate the energy balance at the phase boundaries between the solid and liquid H<sub>2</sub>O layer, and the movements of the positions of these boundaries are calculated by evaluating the heat balance between incoming and outgoing flux at the boundaries considering with latent heat (classically known as a Stefan problem).

## Feasibility of the exploration of the subsurface structures of Jupiter's icy moons by Jovian hectometric radiation

\*Atsushi Kumamoto<sup>1</sup>, Yasumasa Kasaba<sup>1</sup>, Hiroaki Misawa<sup>2</sup>, Fuminori Tsuchiya<sup>2</sup>

1.Department of Geophysics, Graduate School of Science, Tohoku University, 2.Planetary Plasma and Atmospheric Research Center, Graduate School of Science, Tohoku University

A new method for detection of the subsurface structures in the ice crust of Jupiter's moons by using interference patterns found in the spectrogram of the Jovian hectometric radio emissions (HOM) have been proposed. In Jupiter icy moon explorer (JUICE) mission, plasma wave observation around icy moons are planned by using radio and plasma wave instrument (RPWI). In this observations, we will be able to obtain spectrograms of the HOM propagating from Jupiter. Because the emissions directly from Jupiter can be interfered with the emissions reflected at the icy moon's surface and subsurface boundaries, we will find interference patterns in the measured spectrograms. In case of the Earth's Moon, the lunar orbiter SELENE detected the interference patterns in the spectrograms of auroral kilometric radiation (AKR) [Ono et al., 2010; Goto et al., 2011]. Because the interference occurs between AKR directly from the earth and AKR reflected at the lunar surface, the amplitude of the interference patterns are almost constant. In case of Jupiter's icy moons, HOM directly from Jupiter, HOM reflected at the icy crust surface, and HOM reflected at the fully-frozen/partial-melted or high/low-porosity boundary in the ice crust. Due to slight phase difference between HOM emissions reflected at the surface and subsurface boundaries, the amplitude of the interference patterns will be modulated. The depth of the liquid ocean can be determined the frequency width of the modulation. Assuming that the frequency of HOM is ~10 MHz, the permittivity of the icy crust is 3, permittivity of the melted ice is 87, loss rate in the icy crust is 2-9 dB/km, and spacecraft height is 500 km, the maximum detection depth is estimated to be 6-23 km, which is less than the estimated ice thickness of the Ganymede, 150 km [Kivelson et al., 2002]. On the other hand, we can also expect lower attenuation rate than 2-9 dB/km in a depth range where the ice temperature is much lower than 240 K. The receiver's specifications needed for measurement of the interference patterns in the spectrogram are as follows: (1) Frequency resolution: 100 Hz, and (2) The interval of spectrum measurements: 30 sec. In addition, the following two issues have to be considered in actual application: (a) HOM itself has band structures in the spectrogram due to anisotropy of the emission at the source. (b) The roughness of the surface and subsurface boundaries has to be within the half wavelength (~15 m). (c) The delay by inhomogeneity of TEC of the moon's ionosphere has to be less than the half of the period of the HOM (~0.05msec), which corresponds to the dTEC ~  $9.3 \times 10^{12} \text{ m}^{-2}$ .

Keywords: Passive subsurface radar, Jupiter Icy Moon Explorer (JUICE), Radio and plasma wave instrument (RPWI)

## Software-type Wave-Particle Interaction Analyzer (SWPIA) by RPWI for JUICE: Science objectives and implementation

\*Yuto Katoh<sup>1</sup>, Hirotsugu Kojima<sup>2</sup>, Kazushi Asamura<sup>3</sup>, Yasumasa Kasaba<sup>1</sup>, Fuminori Tsuchiya<sup>1</sup>, Yoshiya Kasahara<sup>4</sup>, Tomohiko Imachi<sup>4</sup>, Hiroaki Misawa<sup>1</sup>, Atsushi Kumamoto<sup>1</sup>, Satoshi Yagitani<sup>4</sup>, Keigo Ishisaka<sup>5</sup>, Tomoki Kimura<sup>6</sup>, Yoshizumi Miyoshi<sup>7</sup>, Masafumi Shoji<sup>7</sup>, Masahiro Kitahara<sup>1</sup>, Ondrej Santolik<sup>8,9</sup>, Jan-Erik Wahlund<sup>10</sup>

1.Graduate School of Science, Tohoku University, 2.Research Institute for Sustainable Humanosphere, Kyoto University, 3.ISAS/JAXA, 4.Univ. Kanazawa, 5.Toyama Pref. Univ., 6.RIKEN, 7.ISEE, Nagoya Univ., 8.Institute of Atmospheric Physics, 9.Faculty of Mathematics and Physics, Charles University, 10.Swedish Institute of Space Physics, Uppsala

We present science objectives of Software-type Wave-Particle Interaction Analyzer (SWPIA), which will be realized as a software function of Low-Frequency receiver (LF) running on the DPU of RPWI (Radio and Plasma Waves Investigation; PI: J.-E. Wahlund, IRF-Uppsala, Sweden) for the ESA JUICE mission. SWPIA conducts onboard computations of physical quantities indicating the energy exchange between plasma waves and energetic ions. Onboard inter-instruments communications are necessary to realize SWPIA, which will be implemented by efforts of RPWI, PEP (Particle Environment Package; PI: Stas Barabash, IR-Kiruna, Sweden) and J-MAG (JUICE Magnetometer; PI: M. Dougherty, ICL, UK). By providing the direct evidence of ion energization processes by plasma waves around Jovian satellites, SWPIA contributes scientific output of JUICE as much as possible with keeping its impact on the telemetry data size to a minimum.

SWPIA measures the energy transfer process between energetic particles and electromagnetic plasma waves [Fukuhara et al., EPS 2009; Katoh et al., AnGeo 2013]. SWPIA will be firstly realized in the ERG satellite mission of JAXA to measure interactions between relativistic electrons and whistler-mode chorus in the Earth's inner magnetosphere. We will apply SWPIA to ion-scale wave-particle interactions occurring in the Jovian magnetosphere. SWPIA clarifies where/when/how heavy ions are energized by waves in the region close to Ganymede and other Jovian satellites. In SWPIA of RPWI for JUICE, we focus on the interactions between energetic ions (a few to tens of keV) and ion cyclotron waves (typically less than 1 Hz). SWPIA uses wave electromagnetic field and ion velocity vectors provided by RPWI sensors and PEP, respectively, with referring three-components of the background magnetic field detected by J-MAG. SWPIA measures a relative phase angle between the velocity vector  $v_i$  of  $i$ -th particle of charge  $q_i$  and the wave electric field vector at the timing of particle's detection ( $E(t_i)$ ) and computes an inner product of  $W(t_i) = q_i E(t_i) \cdot v_i$ , where  $W(t_i)$  corresponds to the variation of the kinetic energy of the  $i$ -th energetic particle. We accumulate  $W$  for detected particles to obtain  $W_{int} = \sum_i W(t_i)$ , and we expect statistically significant values of  $W_{int}$  for the case of the measurement at the site of efficient wave-particle interactions. In this presentation, we discuss details of the implementation of SWPIA of RPWI and inter-instruments communications among RPWI-PEP-J-MAG of JUICE.

Keywords: Jovian magnetosphere, Jovian satellite, wave-particle interactions

## Development of a low-energy energetic neutral atom analyzer (PEP/JNA) for JUICE

\*Kazushi Asamura<sup>1</sup>, Yoshifumi Saito<sup>1</sup>, Manabu Shimoyama<sup>2</sup>, Yoshifumi Futaana<sup>2</sup>, Yoshizumi Miyoshi<sup>3</sup>, Takeshi Sakanoi<sup>4</sup>

1.Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 2.Swedish Institute of Space Physics, 3.Nagoya U., 4.Tohoku U.

We are developing a low-energy (10eV-3keV) energetic neutral atom analyser (PEP/JNA) which is to be onboard European JUICE spacecraft. Ganymede has its own intrinsic magnetic moment. There is considered to be a mini-magnetosphere around Ganymede because of interactions between plasma in Jovian magnetosphere and Ganymede's magnetic field. However, its characteristics will be different from terrestrial one, since Alfvén Mach number of upstream plasma flow (corotational plasma flow around Jupiter) is small. JNA (Jovian Neutral Analyzer) will reveal characteristics of Ganymede's magnetosphere in terms of measurement of scattered/sputtered particles generated by precipitation of plasma particles onto Ganymede's surface. Measurement of these particles will provide spatial distribution of plasmas in remote sense, since electric/magnetic field do not affect trajectories of neutral particles. We will discuss current status of JNA.

Keywords: Energetic neutral atom, JUICE

## Responses of Io Plasma Torus to middle magnetosphere of Jupiter

\*Fumiharu Suzuki<sup>1</sup>, Kazuo Yoshioka<sup>1</sup>, Go Murakami<sup>2</sup>, Fuminori Tsuchiya<sup>3</sup>, Tomoki Kimura<sup>4</sup>, Ichiro Yoshikawa<sup>1</sup>

1.The University of Tokyo, 2.JAXA, 3.Tohoku University, 4.RIKEN

The satellite of Jupiter, Io, is volcanically active and ejects ionized gas such as sulfur and oxygen into space. This gas forms a torus encircling Jupiter along the orbit of Io, and called Io Plasma Torus (IPT). Jupiter has by far the most energetic and brightest aurorae in the solar system. During the Cassini spacecraft's flyby of Jupiter (October, 2000-March, 2001), the UV spectrometer witnessed a puzzling phenomenon. Both of IPT and Jupiter's aurorae were sometimes brightened almost simultaneously. The torus emissions reflect the state of the inner magnetosphere while the aurora emissions are an index of activity in the middle magnetosphere. This fact might suggest an energy transport process from the middle to inner magnetosphere, but it has not been understood yet. The dataset of Cassini was insufficient to reveal the process due to its low-temporal resolution. The HISAKI/EXCEED was launched in September 2013 by the Epsilon rocket. Now it is orbiting around the Earth. EXCEED is a space telescope dedicated for planets and has an advantage of long-term and continuous monitoring of the aurorae and IPT at the same time. We find that electron temperature in IPT increases 11 hours after the transient aurora brightening events. It suggests inward flow of hot electrons from the middle magnetosphere to IPT. On middle of January 2015, EXCEED observed gradual increase in EUV brightness of IPT. This observation suggests enhancement of Io's volcanic activity. Characteristics of the transient IPT brightening events also changed during the volcanic event. In this presentation, we will show the energy transport process in the Jovian magnetosphere by analyzing of the EXCEED observation.

Keywords: Jupiter, magnetosphere, aurora, Io plasma torus



## Horizontal and vertical structures of the Jovian IR aurora from plasma and neutral atmospheres: Observation by SUBARU/IRCS with Adaptive Optics

Shota Fujisawa<sup>1</sup>, Yasumasa Kasaba<sup>1</sup>, \*Hajime Kita<sup>1</sup>, Chihiro Tao<sup>2</sup>, Takeshi Sakanoi<sup>3</sup>, Masato Kagitani<sup>3</sup>

1.Dep. Geophysics, Tohoku Univ., 2.NiCT, 3.PPARC, Tohoku Univ.

In the rotational driven magnetosphere of Jupiter, the momentum and energy transfers between thermosphere and ionosphere is the key part of the magnetosphere, ionosphere and thermosphere (MIT) coupling. Jovian aurora, emitted from this region through this process, does not only shows the strength of magnetospheric activities but also affects to the ion-neutral interactions.

Previous observations have shown the morphological difference between plasma ( $H_3^+$ ) and neutral (IR  $H_2$ ) emissions. It suggests the difference between strong electron precipitating area exciting the plasma emissions and the heated area exciting the neutral emissions. In order to investigate their emission mechanisms and relationships to the energy injection processes, the comparison between  $H_3^+$  fundamental ( $v_2=1 \rightarrow 0$ ) and overtone ( $v_2=2 \rightarrow 0$ ) lines is also important. The different emission altitude of both emissions is caused by their different sensitivity to surrounding atmospheric temperature and density.

We first executed the quasi-simultaneous comparisons of the horizontal and vertical emission profiles in  $H_3^+$  fundamental,  $H_3^+$  overtone, and  $H_2$  polar emissions, by the near infrared spectroscopy of Jovian polar emissions using SUBARU/IRCS on 31 January 2015.  $H_2$  IR emission and  $H_3^+$  overtone emissions are seen simultaneously in K-band spectra (2.03-2.22  $\mu\text{m}$ ), and  $H_3^+$  fundamental emission in L-band (3.31-3.98  $\mu\text{m}$ ) is quasi-simultaneously taken by short interval,  $\sim 5$  min. We also simultaneously took the slit viewer image of the  $H_3^+$  fundamental line or K-band filter. During these observations, we used the adaptive optics system (AO188) when Galilean satellites could be used as a guide star, and achieved high spatial resolution,  $\sim 0.2$  arcsec ( $\sim 320$  km of Jupiter). The slit was set along the rotational axis when AO188 could be used.

First, we compared the horizontal flux profiles. The morphological difference between  $H_3^+$  fundamental and overtone emissions are small. Both have clear main oval emissions like the averaged UV aurora profile. On the other hand, IR  $H_2$  emission does not show clear enhancement at the main oval. We also derived the horizontal profiles of temperature and column density from those emissions.  $H_3^+$  fundamental lines have a better correlation with column density.  $H_3^+$  overtone lines are more related to temperature. On the other hand, IR  $H_2$  emission intensity does not show clear correlations.

Next, we derived the vertical structures of their volume emissivity profiles by "onion peeling" method. We confirm the result of Uno et al. (2014), the similar emission peak altitude between IR  $H_2$  and  $H_3^+$  overtone emissions in K-band. We also found that the peak altitude of  $H_3^+$  fundamental emission was lower than them. Although the derived  $H_3^+$  vertical emission profiles are not contradict to the theoretical models, and their derived temperatures represent those of emission peak altitudes, it is hard to explain the vertical profiles of IR  $H_2$  volume emissivity by a simple 'thermal excitation model'. It is also hard to explain the fact that the derived temperature from  $H_2$  emissions from higher altitude ( $\sim 700$  K) is lower than that from  $H_3^+$  fundamental emission in lower altitude ( $\sim 1,000$  K). We are now investigating possible scenarios for those points.

In May 2016, we will observe Jovian IR emission again, simultaneously with UV aurora by Hubble Space Telescope, EUV aurora and Io torus by Hisaki/EXCEED, and the upstream solar wind by NASA Juno spacecraft approaching to Jupiter. It will be the chance to solve the problems raised in this study.

Keywords: Jupiter, infrared aurora, horizontal profiles, vertical profiles

## Statistical study of the response of Jovian EUV aurora to the solar wind from Hisaki observations

Hajime Kita<sup>1</sup>, Tomoki Kimura<sup>2</sup>, Chihiro Tao<sup>3</sup>, \*Fuminori Tsuchiya<sup>1</sup>, Hiroaki Misawa<sup>1</sup>, Takeshi Sakanoi<sup>1</sup>, Yasumasa Kasaba<sup>1</sup>, Go Murakami<sup>4</sup>, Kazuo Yoshioka<sup>5</sup>, Atsushi Yamazaki<sup>4</sup>, Ichiro Yoshikawa<sup>6</sup>

1.Graduate School of Science, Tohoku University, 2.Nishina-Center for Accelerator Based Science, RIKEN, 3.National Institute of Information and Communications Technology, 4.Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 5.Department of Earth and Planetary Science, Graduate School of Science, University of Tokyo, 6.Department of Complexity Science and Engineering, University of Tokyo

In order to reveal the solar wind response of Jovian extreme ultraviolet (EUV) auroral activity, we made a statistical analysis of Jovian EUV aurora obtained from long term Hisaki observation. The EUV emission from hydrogen molecule is excited by collision with high energy electron. The main oval is one of the components of Jovian EUV aurora where the auroral particle precipitations are caused by the rotationally driven field-aligned current system. It is theoretically expected that angular velocity of magnetospheric plasma increases when the Jovian magnetosphere is compressed by enhanced solar wind pressure, which decreases the field-aligned current. Regarding this scenario, increase of the solar wind dynamic pressure is expected to be anti-correlated with the intensity of the EUV aurora. A previous observation such as that by International Ultraviolet Explorer (IUE) or Hubble Space Telescope (HST) showed the time variability of the EUV aurora, while their data still limited in continuity over solar wind variation with good time resolution. On the other hand, Hisaki satellite is an earth-orbiting EUV spectroscopy launched in 2013 which has been continuously monitoring Jovian EUV auroral activity. Therefore, the Hisaki data sets are effective for investigating the solar wind response of Jovian aurora.

The purpose of this study is to investigate the solar wind response of Jovian EUV aurora observed by Hisaki. We used the EUV data set obtained from Dec. 2013 to Feb. 2014 and from Dec. 2014 to Feb. 2015. We compare the total EUV power over 900-1480 Å and solar wind dynamic pressure which is extrapolated at Jupiter using a one-dimensional magnetohydrodynamic (MHD) model.

Superposed epoch analysis indicated that Jovian EUV aurora increases with the enhancement of the solar wind dynamic pressure. We also found a correlation between the total power of EUV aurora and the duration of the rarefaction region of the solar wind before the enhancement of the dynamic pressure. The similar trend could also be found in the thermal current, i.e., incoming electron flux increased with the duration of rarefaction region.

One possible scenario is that mass loading from Io increases the electron density in the Jovian middle magnetosphere and it also increases seed electron of the thermal current whose energy is several keV. The solar wind compression causes adiabatic acceleration of thermal current and then EUV aurora increases. However, it is still unclear how the angular velocity distribution and brightness distribution vary during the solar wind compression.

Keywords: Jupiter, Aurora, Solar wind response

## Variation in SII and SIII brightness distribution of Io plasma torus based on Hisaki/EXCEED and ground based observation data

\*Mika Shishido<sup>1</sup>, Takeshi Sakanoi<sup>1</sup>, Masato Kagitani<sup>1</sup>, Fuminori Tsuchiya<sup>1</sup>, Ichiro Yoshikawa<sup>5</sup>, Atsushi Yamazaki<sup>3</sup>, Kazuo Yoshioka<sup>2</sup>, Go Murakami<sup>3</sup>, Tomoki Kimura<sup>4</sup>

1.Graduate School of Science, Tohoku University, Department of Geophysics, 2.Graduate School of Science, The University of Tokyo, Department of Earth and Planetary Science, 3.ISAS/JAXA, 4.RIKEN, 5.Graduate School of Science, The University of Tokyo, Department of Complexity Science and Engineering

We report the time and spatial variation of sulfur ion emission line from the Io plasma torus ([SII]76.5nm and[SIII]68nm) to understand the dynamical process in the torus associated with Io's volcanic event during the period from December 2014 to March 2015, using the data obtained by Hisaki/EXCEED. The large quantities of gas were ejected by Io's volcanoes, principally oxygen and sulfur atoms and their compounds. Once they are ionized through electron impact and charge exchange, the ions are accelerated to the nearly corotational flow of the ambient plasma to form a torus of ions (the Io plasma torus, about  $6R_J$  from the center of Jupiter) surrounding Jupiter. The fresh ions lose their pickup energy to the ambient electrons through Coulomb collisions. Ultimately, the torus electrons lose energy by moving electrons bound to ions into excited states, leading to the prodigious extreme ultraviolet (EUV), ultraviolet, and visible emissions from the torus. Hisaki/EXCEED is an EUV spectrograph launched in September 2013. During the period from December 2014 to March 2015, Io's outburst was observed by EXCEED, and the increase in the pickup ions were anticipated along with the increase in the neutral gas. During this period, we identified the time variation of sulfur ion temperature associated with enhanced volcanic activities from images of sulfur ion emission at 76.5nm(SII) and 68nm(SIII) (an integration time for each image is about 50 min) to find increase in ion temperature through the ion-pickup process and cooling by transferring energy to electrons. In addition, we carried out the measurement of SII 673 nm emission with visible spectrograph on T60 telescope at Haleakala, Hawaii, which has high spatial resolution. In this talk we give the results on the scale height and ion temperature of sulfur torus derived from spatial distribution of Io torus with EXCEED associated with the Io's volcanic event, and the comparison with ground-based visible data obtained with T60.

time variation of 130.4nm atomic oxygen emission near Io observed by hisaki/EXCEED

\*Ryoichi Koga<sup>1</sup>, Fuminori Tsuchiya<sup>1</sup>, Masato Kagitani<sup>1</sup>, Takeshi Sakanoi<sup>1</sup>, Mizuki Yoneda<sup>1</sup>, Ichiro Yoshikawa<sup>2</sup>, Kazuo Yoshioka<sup>2</sup>, Go Murakami<sup>3</sup>, Atsushi Yamazaki<sup>3</sup>, Tomoki Kimura<sup>4</sup>

1.PPARC, Tohoku University, 2.University of Tokyo, 3.ISAS/JAXA, 4.RIKEN

Io, one of the Jupiter moon, is the most volcanically active body in the solar system because of its large tidal force from Jupiter. Main components of Io's atmosphere are SO<sub>2</sub>, and its dissociative sulfur and oxygen atoms. The rest components (a few percent) are neutral sodium and potassium. Origins of the atmosphere are thought to be gaseous plume driven by volcanic activity and sublimation of SO<sub>2</sub> frost, but it is not clear which is dominant. The object of this study is to understand the atmospheric generation process through the time variation of 130.4nm atomic oxygen emission near Io.

The brightening event of the extended sodium nebula was reported by the ground imaging observation from December 2014 to May 2015 (Yoneda et al., 2015). To show the behavior of atomic oxygen during the period, we analyzed the time variation of atomic oxygen, one of the main components of volcanic gas, -near Io (1 Jupiter radius) by using data observed by hisaki/EXCEED. We selected observed data when Io was in the dawn side (Io's phase angle of 45~135) and in the dusk side (225~315), and accumulated photons which came from around Io within a range of 60" on each day to acquire enough S/N. We carefully selected data when the local time of Hisaki was in the range of 20-4h to avoid the contamination from geocorona. As a result, it is found that the brightness of atomic oxygen emission was 11R at the beginning of January, but started to increase in the middle January and showed the maximum of ~32R in the middle of February. It decreased by the end of May and returned the normal brightness (10R). We also confirmed that the emission in the dusk side more intense than that in the dawn side by about 1.2 times throughout the period.

The result shows amount of atomic oxygen near Io increases when Io is volcanically active. To discuss the relation between the atmospheric generation process and volcanic activity, we need to derive atomic oxygen column density from emission brightness of atomic oxygen at 130.4nm. While both solar resonant scattering and electron impact excitation could contribute to 130.4nm atomic oxygen emission, we estimate the contribution of electron impact excitation is several hundred times as high as that of solar scattering. The estimation shows electron impact excitation is dominant and it is likely to explain dawn-dusk asymmetry of 130.4nm atomic oxygen emission because thermal electron temperature in the dusk side is higher than that in the dawn side. In this talk, we derive the atomic oxygen column density and discuss how volcanic activities influence characteristic of Io's atmosphere according to the estimation.

Keywords: Jupiter, Io, hisaki/EXCEED, volcanic activity

## Science Experiments with the Trojan Asteroid Lander in the Solar Powered Sail Mission

\*Tatsuaki Okada<sup>1</sup>, Jean-Pierre Bibring<sup>2</sup>, Stephahn Ulamec<sup>3</sup>, Jens Biele<sup>3</sup>, Ryosuke Nakamura<sup>4</sup>, Hajime Yano<sup>1</sup>, Yoko Kebukawa<sup>5</sup>, Jun Aoki<sup>6</sup>, Yosuke Kawai<sup>6</sup>, Hikaru Yabuta<sup>6</sup>, Motoo Ito<sup>7</sup>, Yoshifumi Saito<sup>1</sup>, Shoichiro Yokota<sup>1</sup>, Noel Grand<sup>8</sup>, Herve Cottin<sup>8</sup>, Laurent Thirkell<sup>9</sup>, Christelle Briois<sup>9</sup>, Takahiro Iwata<sup>1</sup>, Ayako Matsuoka<sup>1</sup>, Jun Matsumoto<sup>10</sup>, Hideki Kato<sup>1</sup>, Osamu Mori<sup>1</sup>, Junichiro Kawaguchi<sup>1</sup>

1.Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 2.Institut d'Astropysique Spatiale, Orsay, 3.German Aerospace Center, 4.National Institute of Advanced Industrial Science and Technology, 5.Yokohama National University, 6.Osaka University, 7.Kochi Institute for Core Sample Research, Japan Agency for Marine-Earth Science and Technology, 8.LISA, Univ-Paris-XII, 9.Centre National de la Recherche Scientifique, Orleans, 10.University of Tokyo

Scientific exploration on the Jupiter Trojan asteroid is under study for the solar-powered sail (SPS) mission. This mission includes a scientific lander jointly studied by Japanese and European engineers and scientists [1]. We present the objectives and the strawman payloads for this mission. The SPS is a candidate as the next medium class space science mission in Japan. This engineering mission is based on the technologies such as the solar sail and the ion engine system inherited from Ikaros and Hayabusa missions, respectively. With this hybrid propulsion system, the spacecraft will cruise to the Jupiter and beyond, even if a radioisotope thermoelectric generator (RTG) is not used. A Trojan asteroid will be investigated by remote sensing after rendezvous, and then a small lander will be deployed from the mothership to conduct *in situ* experiments on the asteroid. As an option, sample will be returned to the Earth. Mission duration is typically 15 years to arrive at the Trojan asteroid, and 30 years in total for Earth return. The shortest one way trip to the asteroid is less than 12 years. The lander should be designed within 100 kg wet mass. Total mission payloads should be within 20 kg, including all the science payloads, sampling and sample return systems [2].

Jupiter Trojan asteroids are located around the Sun-Jupiter Lagrange points. Most of them are volatile-rich D- or P-type asteroids, and their origin and evolution, composition and physical conditions still remain unknown. In a classical model of solar system evolution, they formed around the Jupiter orbit and survive until now. But in a recent model such as Nice model [3], they formed at the far end of the solar system and transferred inward due to dynamical migration of giant planets. Physical, mineralogical, and isotopic studies of surface materials could solve their origin and evolution processes, as well as the solar system formation [4]. To achieve these goals, *in situ* observations using the lander is planned, as well as the asteroid global characterization with a near-infrared hyperspectral imager.

Geological, mineralogical, and geophysical observations will be conducted to characterize the landing site, by using a panoramic camera, an infrared hyperspectral imager, a magnetometer, and a thermal radiometer. The surface conditions and composition will be investigated with a close-up imager and a Raman spectrometry. The imager is also used to check the conditions whether the sampling could be done or not. If the configuration is unsuitable for sampling, the lander must relocate and change the configuration. The surface and subsurface materials will be collected into a carousel by bullet-type and pneumatic drill type samplers, respectively. Samples in the each case of carousel will be viewed by infrared microscope to identify them. Those samples will be transferred for evaporation of volatiles for high resolution mass spectrometry (HRMS). Some samples will be heated for pyrolysis for isotopic analysis. Mass resolution  $m/\Delta m > 30,000$  is required to investigate isotopic ratios of D/H,  $^{15}\text{N}/^{14}\text{N}$ , and  $^{18}\text{O}/^{16}\text{O}$ , as well as molecules from organic matters ( $M = 30$  to  $1000$ ). The MULTUM type in Japan and the Cosmorbitrap type in France are being

investigated for the HRMS. A set of strawman payloads are now considered to meet the science, mission, and system requirements and constraints (total mass < 20kg, and total energy consumption < 600 WHr). They will be finally determined by the international announce of opportunity.

References: [1] Mori O. et al. (2015) *11<sup>th</sup> Low-Cost Planetary Missions Conf.*, S3-10. [2] Saiki T. et al. (2015) *ISSFD2015*, S19-3, #84. [3] Morbidelli A. et al. (2005) *Nature* 435, 462-466. [4] Yano H. et al., (2014) *COSPAR 2014*, B0.4-2-14. [5] Mori O. et al. (2016) *Lunar Planet. Sci. Conf.* , submitted.

Keywords: Trojan asteroid, Solar Sail, Lander, High Resolution Mass Spectrometry, Solar-Powered Sail, Subsurface sample