

火星探査衛星MAVEN搭載IUVSと大気大循環モデルの比較で得られた熱圏密度温度構造

Comparison of the Martian thermospheric density and temperature from IUVS/MAVEN data and general circulation modeling

*中川 広務¹、Medvedev Alexander³、Mockel Chris³、Yigit Erdal³、黒田 剛史¹、Hartogh Paul³、寺田 香織¹、寺田 直樹¹、関 華奈子²、Schneider Nicholas⁴、Jain Sonal⁴、Evans Scott⁵、Deighan Justin⁴、McClintock William⁴、Lo Daniel⁶

*Hiromu Nakagawa¹, Alexander S Medvedev³, Chris Mockel³, Erdal Yigit³, Takeshi Kuroda¹, Paul Hartogh³, Kaori Terada¹, Naoki Terada¹, Kanako Seki², Nichlas M Schneider⁴, Sonal Jain⁴, Scott Evans⁵, Justin I Deighan⁴, William McClintock⁴, Daniel Lo⁶

1.東北大学 大学院理学研究科 地球物理学専攻太陽惑星空間物理学講座 惑星大気物理学分野、2.東京大学 大学院理学系研究科、3.Max Plank Institute for Solar System Reserach, Gottingen, Germany、4.Laboratory for Atmospheric and Space Physics, University of Colorado Boulder、5.Computational Physics, Inc.、6.Lunar and Planetary Laboratory, University of Arizona
1.Planetary Atmosphere Physics Laboratory, Department of Geophysics, Graduate School of Science, Tohoku University, 2.Graduate School of Science, University of Tokyo, 3.Max Plank Institute for Solar System Reserach, Gottingen, Germany, 4.Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, 5.Computational Physics, Inc., 6.Lunar and Planetary Laboratory, University of Arizona

The Mars Atmosphere and Volatile Evolution (MAVEN) Mission, which operates for slightly over one year to date, has been specifically designed for investigating the upper atmosphere. The Imaging Ultraviolet Spectrograph (IUVS) onboard MAVEN measures spectra of mid- and far UV atmospheric emission, which are used for retrieving vertical density profiles of CO₂ and other species. Newly released IUVS/MAVEN measurements of CO₂ density in the Martian thermosphere have been used for comparison with the predictions of the Max Planck Institute Martian General Circulation Model (MPI-MGCM). In this study, we focus on the October 2014 campaign in which a total of 122 density profiles were obtained for the period between 18 and 22 October (Ls=216.68-218.94). IUVS nicely covers the thermosphere in the altitude range of 130-220 km. The MGCM demonstrated the sensitivity of simulated density and temperature profiles on (i) solar flux, (ii) atomic oxygen, and (iii) small-scale gravity waves (GWs). It is the only MGCM to date that includes a parameterization of effects of subgrid-scale GWs with broad spectra.

The simulations reproduced (within one standard deviation) the available zonal mean density and derived temperature above 130 km. The comparison shows a great role of gravity waves in the thermosphere, and in bringing the simulated density and temperature closer to observations. The MGCM replicated the observed dominant zonal wavenumber-3 non-migrating tide, which was already reported by MAVEN measurements (Lo et al., 2015). The simulations also demonstrated that it represents a non-moving imprint of the topography in the thermosphere.

This comparison confirms that, with the current state of knowledge of the Martian thermospheric physics, MGCM can reproduce its state and variability. Further observations will help to constrain physical parameterizations and improve modeling capabilities.

キーワード：重力波、熱潮汐、MAVEN

Keywords: Gravity waves, Tides, MAVEN

Runaway acidification on early Mars triggered by atmospheric evolution



Runaway acidification on early Mars triggered by atmospheric evolution

*田畑 陽久¹、関根 康人¹*Haruhisa Tabata¹, Yasuhito Sekine¹

1. 東京大学

1. University of Tokyo

Mineralogical and chemical observational data of sedimentary rocks deposited on early Mars suggest that chemistry of Mars' surface water had shifted from neutral pH (pH~7-9) to highly acidic (pH~2) at around 3.5 billion years ago (Ga) (Bibring et al., 2006; Ehlmann et al., 2011). One proposed mechanism for this acidification is photo-oxidation of ferrous iron dissolved in surface water (Hurowitz et al., 2010). When UV light is irradiated to surface water, ferrous iron are oxidized to ferric iron by producing H⁺ with precipitation of Fe³⁺ minerals (see Equations 1 and 2 below). Hurowitz et al. (2010) show that H⁺ produced via this mechanism are quantitatively sufficient to explain the mineralogical and chemical compositions of sedimentary rocks in Meridiani Planum observed by Mars Exploration Rover, Opportunity (e.g., Tosca et al. 2005).



Although this mechanism would have promoted acidification at 3.5 Ga on Mars, it remains unsolved why surface water with neutral pH had been maintained before 3.5 Ga, and what the trigger for the acidification was. In this study, we propose a new hypothetical scenario to explain the chemical transition of surface water from neutral to acidic pH by considering positive and negative feedbacks in association with photo-oxidation of ferrous iron.

Ferrous iron has two stable dissolved species; Fe²⁺ and Fe(OH)⁺, depending on pH of water. It has been reported that both species are oxidized when irradiated with UV (< 200 nm). But, Fe(OH)⁺, which becomes the dominant species at pH > 9, is also oxidized with irradiation of visible light (300-400 nm) (Braterman et al., 1983). In a thick CO₂ atmosphere (~1 bar) with several ppms of SO₂, UV light with short wavelength (< 300 nm) is shielded by the atmospheric gas species. In this case, photo-oxidation of Fe(OH)⁺ proceeds in surface water, whereas that of Fe²⁺ is limited. In surface water with neutral to alkaline pH, photo-oxidation of Fe(OH)⁺ acidifies the water forming H⁺. However, the concentration of Fe(OH)⁺ drastically decreases around pH~6.5 via the conversion into Fe²⁺. This decrease in Fe(OH)⁺ concentration, in turn, results in dampening Fe(OH)⁺ photo-oxidation. Consequently, in a thick atmosphere, pH dependence of Fe(OH)⁺ photo-oxidation works as a negative feedback to maintain surface water pH around neutral.

On the other hand, in a thin CO₂ atmosphere (0.1 bar or less) with < 1 ppm of SO₂, UV light reaches to the surface water, leading to photo-oxidation of both Fe²⁺ and Fe(OH)⁺ in surface water. In this case, even if the concentration of Fe(OH)⁺ in surface water decrease at pH~6.5, acidification proceeds via photo-oxidation of Fe²⁺. As a result, a runaway acidification to highly acidic surface water occurs via this positive feedback.

Here, we discuss the above possibility more quantitatively using the previous laboratory data on photo-oxidation rates of Fe²⁺ under acidic conditions by Jortner et al. (1962) and those by Braterman et al. (1983) under neutral pH conditions. We calculate the total production rate of H⁺ both via the photo-oxidation of Fe²⁺ and Fe(OH)⁺ for various atmospheric compositions and pressures. Based on our sensitivity study of the abundances of CO₂ and SO₂ to acidification of surface water, the critical atmospheric compositions to drive the runaway acidification are discussed.

キーワード：火星、酸性化、大気進化、光酸化、地球化学

Keywords: Mars, acidification, atmospheric evolution, photooxidation, geochemistry

Warmer Wetter Mars in the Past?

*Sushil K Atreya¹, Paul Mahaffy², Christopher Webster³, Pamela Conrad⁴, Michael Wong⁵

1.University of Michigan Ann Arbor, 2.NASA/GSFC, 3.NASA/JPL, 4.GSFC, 5.U-Michigan

The isotopes of CO₂ (¹³C/¹²C, ¹⁸O/¹⁶O), H₂O (D/H), N₂ (¹⁵N/¹⁴N), Ar (³⁸,⁴⁰Ar/³⁶Ar), Kr and Xe are excellent indicators of climate change in the atmosphere of Mars. Recent high precision measurements of those isotopes with the quadrupole mass spectrometer and the tunable laser spectrometer of the Sample Analysis (SAM) suite of instruments on the Curiosity Rover clearly show that the atmosphere of Mars has been substantially depleted over the past four billion years. At the same time, both geological evidence and a comparison of the D/H isotope ratios in water vapor in the atmosphere with the Hesperian-era Gale Crater smectite rock fines, and even older Mars meteorites suggest a relatively large abundance of (liquid) water on the surface of Mars in the past. With the exception of xenon, the above isotopes inform about the change since roughly 4 Ga. The isotopic fractionations in xenon suggest, in addition, a massive H₂-driven hydrodynamic escape very early in the geologic history of Mars. Employing the isotopic record in the atmosphere and rock fines, we investigate a scenario of Mars where atmospheric composition and relatively high atmospheric pressure resulted in warmer conditions necessary for maintaining surface liquid water at least intermittently through late Noachian/early Hesperian, followed by a gradual loss of the atmosphere by escape since then, hence warmer and wetter conditions in the past compared to Mars' present cold and arid state.

Keywords: Mars, Climate Evolution, Isotopes

Viking Lander 2搭載地震計データ復元の現状

Current status of data restoration of the seismometer onboard Viking Lander 2

*山本 幸生¹、山田 竜平²*Yukio Yamamoto¹, Ryuhei Yamada²


1.国立研究開発法人宇宙航空研究開発機構宇宙科学研究所、2.自然科学研究機構国立天文台

1.The Institute of Space and Astronautical Science, 2.National Astronomical Observatory of Japan

1975年に打ち上げられたViking Lander 2には地震計が搭載された。40年が経過した現在も、人類が保有する唯一の火星における地震計データである。このデータは火星での地震活動を発見するに至らなかったが、データそのものが整備されておらず、検証を行える状況になかった。我々はこの地震計データの復元を試み、過去の論文で図示されたデータと一致する結果を得ることができた。これを元に全データを処理したところ、源泉データに幾つかの問題点が存在し、そのままでは科学解析を行う上で前処理として多大な労力が必要となることが判明した。本研究では、これらの課題を克服し、新しいデータセットを提供することを目的とし、データ処理を進めている。

キーワード：ヴァイキング、地震計、データ

Keywords: Viking, Seismometer, data



火星大気中音波検出器の開発のための過酷環境試験下マイク動作特性の解析
Analysis of microphones operating characteristics under harsh environment
in order to develop a sound detector in Martian atmosphere

*藤津 裕亮¹、山本 真行¹

*Hiroaki Fujitsu¹, Masa-yuki Yamamoto¹

1.高知工科大学

1.Kochi University of Technology

1. 背景

2020年代、宇宙航空研究開発機構(JAXA)により火星探査機の打ち上げが計画されており、シリーズ的な火星探査の実現が期待されている。2016年現在、火星大気中における音波観測は未だ行われていない。そこで探査用ローバーに本研究を反映して設計・開発するマイクを搭載すれば、ダスト現象に伴う火星大気中の音の計測だけでなく、大気中の物理量の間接計測も可能になるとともに、春先の季節に発生する可能性のあるガス放出現象など最近注目され始めた火星地表面活動にもフォーカスしたりリモートセンシングに応用できると期待される。

2. 目的

火星探査機に搭載するマイクの過酷環境下での試験を行い、コンデンサマイクおよびMEMS (Micro Electro Mechanical Systems)マイクの二種類のマイク評価モデル(BBM)を試作し、どちらが火星大気での動作に適しているか評価・選定を行うこと、およびマイクアレイによる検知性能向上について調整することを目的とする。

3. 実験概要

火星地表大気条件として、CO₂成分が95%を占め、地表面気圧7 hPa、夜間気温-120 °Cという過酷環境を模した試験を2015年1月に千葉工業大学にて行った。さらに高知工科大学にてマイクBBMの較正実験を行った。可聴音の下限である20 Hz以下の音をインフラサウンドと呼び、将来の火星大気中でのインフラサウンド信号の検知も目標としているため、可聴音の他にインフラサウンドを入力し、較正データとして用いた。真空チャンバを密閉容器として使用し、これに接続した微小容積のシリンジを使って周期的に圧縮・吸引を行い、大気圧に対する微気圧変動を精密波形として作り出し、この波形をインフラサウンド入力とし較正実験を行った。実験系に周波数0.1 Hz、振幅1 Paの波形を入力した時の結果から既存のインフラサウンドセンサ(Chaparral Physics製 Model25)とマイクBBMの性能評価を行った。

4. 実験結果

高知工科大学で行った実験より、インフラサウンドを用いた測定結果では共に0.1 Hzに一致するスペクトルピークが得られ、波形の最大振幅はChaparral Physics製Model25が0.05989 Pa、コンデンサマイクが0.07657 Paとなった。このことより、使用したコンデンサマイクは0.1 Hz帯では既存のインフラサウンドセンサより高い検出精度をもつことが分かった。その後、同じ0.1 Hz、1 Paの波形入力にてコンデンサマイクとMEMSマイクの比較を行った。コンデンサマイクは0.1 Hzのインフラサウンドを検出できているがMEMSマイクはほとんど検出できていないことが分かった。

5. 考察

千葉工業大学の実験では模擬火星大気中での動作が確認されたが、さらに低圧の1 hPaではMEMSマイクのみ出力信号が乱れ動作が確認できなかった。さらに高知工科大学の実験より、0.1 Hzの波形に対してコンデンサマイクBBMはModel25インフラサウンドセンサに対し約1.28倍の受信性能を持っていることが分かる。これよりコンデンサマイクには0.1 Hz帯での優れたインフラサウンド受信性能があることが分かった。2種類のマイクBBMを比較するとコンデンサマイクは0.1 Hzを入力値の約1/10の圧力で検出することが可能であるがMEMSマイクは同帯域を検出できていないことが分かる。千葉工業大学、高知工科大学にて実施した両実験より、耐苛酷、環境面、検出性能面においてMEMSマイクに比べ、コンデンサマイクが火星大気中の音波検出器として適していると言える。

6. 結論

火星大気模擬環境下において、コンデンサマイク、MEMSマイクの2種類のBBMを試作して動作比較を行い、大気

を模擬した環境下での動作に適したマイクの選定を行った。火星探査における最終的な観測方法は長距離伝搬可能なインフラサウンド帯域での多地点観測に重点を置いている。そのため高知工科大学での実験結果の通り、0.1 Hz程度までのインフラサウンドを観測可能なコンデンサマイクを火星探査用マイクに使用するべきであると結論付ける。

参考文献:

Jeffrey B. Johnson et al., Volcanic eruptions observed with infrasound, GEOPHYSICAL RESEARCH LETTERS, VOL. 31, L14604, doi:10.1029/2004GL020020, 2004.

平塚市博物館火星の大気と気候

http://www.hirahaku.jp/hakubutsukan_archive/tenmon/00000025/72.html, 2015年12月参照。

キーワード：火星、インフラサウンド

Keywords: Mars, Infrasound

火星表面での細胞のその場観察：生命探査顕微鏡

LDM (Life Detection Microscope): In situ imaging of living cells on surface of Mars

*山岸 明彦¹、佐藤 毅彦²、宮川 厚夫¹、佐々木 聡³、吉村 義隆⁴、今井 栄一⁵、長沼 毅⁶、小林 憲正⁷、癸生川 陽子⁷、藪田 ひかる⁸、三田 肇⁹、出村 裕英¹⁰、大野 宗祐¹¹、小林 正規¹¹、波多 英寛¹²

*Akihiko Yamagishi¹, Takehiko Satoh², Atsuo Miyakawa¹, Sasaki Satoshi³, Yoshitaka Yoshimura⁴, Eiichi Imai⁵, Takeshi Naganuma⁶, Kensei Kobayashi⁷, Yoko Kebukawa⁷, Hikaru Yabuta⁸, Hajime Mita⁹, Hirohide Demura¹⁰, Sohsuke Ohno¹¹, Masanori Kobayashi¹¹, Hidehiro Hata¹²

1.東京薬科大学、2.宇宙航空研究開発機構宇宙科学研究所、3.東京工科大学、4.玉川大学、5.長岡技術科学大学、6.広島大学、7.横浜国立大学、8.大阪大学、9.福岡工業大学、10.会津大学、11.千葉工業大学、12.熊本大学

1.Tokyo University of Pharmacy and Life Science, Department of Molecular Biology, 2.Japan Aerospace Exploration Agency, 3.Tokyo University of Technology, 4.Tamagawa University, 5.Nagaoka University of Technology, 6.Hiroshima University, 7.Yokohama National University, 8.Osaka University, 9.Fukuoka Institute of Technology, 10.Aizu Univerisy, 11.Chiba Institute of Technology, 12.Kumamoto University

Past trial of detection of life on Mars by 1970's Viking mission ended up with a negative conclusion [1]. Whereas, numbers of new finding provided by Mars exploration missions in the last decade indicate that there are good reasons to perform another life detection program. The sensitivity of GC-MS onboard the Viking mission was not very high, and was not able to detect the microbes 10^{*6} cells in 1 gram clay [2,3]. Here we propose Life Detection Microscope (LDM) that has much higher sensitivity than the instrument onboard Viking.

Recent observations on Mars have found the evidences of past water activities. MSL Curiosity has reported the temporal increase of methane concentration in Martian atmosphere [4]. The presence of reduced sulfur compound such as pyrite in Martian soil was also detected by MSL [5]. Methane and reduced sulfur compound can be the energy source to support the growth of chemoautotrophic microbes [6]. Possible presence of liquid water at Recurring Slope Lineae has been supported by the detection of hydrated salts [7]. The presence of organic compounds of Martian origin has been reported [8]. These evidences tend to support the possible presence of living microbes near the surface of Mars.

Physical and chemical limits for terrestrial life have been major foci in astrobiology [9], and are summarized in ref. [6]. Combining the environmental factors, anywhere in the Martian environment where we can find the three components, water molecules, reducing compounds and oxidative compounds could be an environment where life can be sustained for long periods of time, if other factors such as temperature, pressure, UV and other radiations permit [6]. Among these factors, most of the factors including ionic radiation, can be endured by terrestrial extremophiles. Only UV can kill the most UV-resistant microbes within minutes. However, UV can be shielded by a-few-centimeter sail layer. These evaluation lead to the conclusion that the Martian soil under a few cm can be the place to support the growth of microbes, if the water activity is higher than 0.6.

We will report the current status of the development of the LDM. We propose to search for cells from a depth of about 5 - 10 cm below the surface, which is feasible with current technology.

Microscopic observation has the potential to detect single cells. We have developed the solution and combination of fluorescence pigments to detect organic compounds, and to differentiate organic compounds surrounded by membrane. The subsequent analysis of amino acids, in the following mission, will provide the information needed to elucidate the origin of the cell.

LDM that we propose here could detect less than 10^{*4} cells in 1 gram clay [6]. LDM is capable of

identifying what we think to be the most fundamental features that a cell should possess to constitute life. Our Investigation Goals are the followings. 1) Identify cell-like structure in which organic compounds are enveloped by membrane, which may represent Martian life. 2) Search for any type of organic compounds in Mars surface samples. The compounds include cells, other biological materials, and abiotic polycyclic aromatic hydrocarbon (PAH). 3) High-resolution characterization of regolith and dust particles.

References: [1] Margulis, L. et al. *J. Mol. Evol.* 14, 223-232 (1979). [2] Glavin, et al, *Earth Planet. Scie. Lett.*, 185, 1-5 (2001). [3] Navarro-González, et al., *Proc. Natl. Acad. Sci. USA.* 103, 16089-16094 (2006). [4] Webster, C.R. et al. *Science Express* Dec. 16 (2014). [5] Ming, D.W. et al. *Science Express* Dec.19 (2013). [6] Yamagishi, A. et al. *Biol. Scie. Space*, 24, 67-82 (2010). [7] Ojha, L. et al. *Nature Geoscie.* 8, 829-832 (2015). [8] Freissient, C. et al. *J. Geophys. Res. Planets* (2015). [9] Marion, G.M. et al. *Astrobiol.* 3, 785-811 (2003).

キーワード：火星、蛍光顕微鏡、生命探査

Keywords: Mars, Fluorescence microscope, Life search

Tracking the MSL-SAM methane detection source location Through Mars Regional Atmospheric Modeling System (MRAMS)

*Jorge Pla-García¹, Scot Rafkin

1. Centro de Astrobiología (INTA-CSIC)

The putative in situ detection of methane by SAM instrument has garnered significant attention. There are many major unresolved questions regarding this detection: 1) Where is the release location? 2) How spatially extensive is the release? 3) For how long is CH₄ released? In an effort to better address the potential mixing and remaining questions, atmospheric circulation studies of Gale Crater were performed with the Mars Regional Atmospheric Modeling System (MRAMS). The model was focused on rover locations using nested grids with a spacing of 330 meters on the innermost grid that is centered over the landing. MRAMS is ideally suited for this investigation. In order to characterize seasonal mixing changes throughout the Martian year, simulations were conducted at Ls 0, 90, 180 and 270. Two additional simulations at Ls 225 and 315 were explored to better understand the unique meteorological setting centered around Ls 270. Ls 270 was shown to be an anomalous season when air within and outside the crater was well mixed by strong, flushing, northerly flow and large amplitude breaking mountain waves: air flowing downslope at night is cold enough to penetrate all the way to the surface. At other seasons, the air in the crater is more isolated -but not completely- from the surrounding environment: mesoscale simulations indicate that the air flowing down the crater rims does not easily make it to the crater floor. Instead, the air encounters very cold and stable air pooled in the bottom of the crater, which forces the air to glide right over the colder, more dense air below. Thus, the mixing of near surface crater air with the external environment is potentially more limited at seasons other than around Ls 270. The rise in CH₄ concentration was reported to start around sol 300 (~Ls 336), peaked shortly after sol 520 (~Ls 82), and then dropped to background values prior to sol 575 (~Ls 103). Two scenarios are considered in the context of the circulations predicted by MRAMS. The first scenario is the release of methane from somewhere outside the crater. The second is a release of methane within the crater. In both cases, the release is assumed to take place near the season when the rise of concentration was first noted (~Ls 336). This is a transitional time at Gale Crater, when the flushing winds are giving way to the more isolated crater scenario. Some preliminary work, including tracer gases into the model, is being performed to establish the amount of mixing during the limited mixing epochs. Preliminary results may support the idea that during periods of limited mixing, there could be enough time for methane to bind to activated mineral surfaces through wind erosion.

Keywords: Mars meteorology, Mars atmosphere, Mars Science Laboratory, Gale crater, methane

火星ユートピア平原におけるフィロケイ酸塩分布

Distribution of phyllosilicates on Utopia Planitia, Mars

菅原 咲季¹、*小川 佳子¹、出村 裕英¹Saki Sugawara¹, *Yoshiko Ogawa¹, Hirohide Demura¹

1.会津大学

1.The University of Aizu

Many studies suggest that abundant liquid water existed in the past on the Martian surface, although there is currently no liquid water on the surface of Mars. The hydrated minerals are generally formed through long-term contact with water or alternation by water. We can get a key to understand the water environment of Mars by studying the observed hydrated minerals.

The wider distribution of hydrated minerals on the southern hemisphere on Mars has been reported in previous studies [e.g. Carter et al., 2013], however, only a limited number of outcrops of hydrated minerals are detected so far in the northern lowlands. The surface of northern lowlands is basically young being covered with lava materials in Hesperian through Amazonian ages. We expect that the deposition below the younger crust of the northern lowlands should host hydrated minerals commonly. The purpose of this study is to examine the distribution of hydrated minerals in the northern lowlands of Mars.

We focus on the comparatively larger craters which should expose the subsurface minerals by impact gardening in the northern lowlands of Mars. Our analysis targets are the craters with diameters ≥ 10 km in Utopia Planitia ranging 25-50N, 90-140E. The number of the craters amounts to 14 in total. We used the data from CRISM (Compact Reconnaissance Imaging Spectrometer for Mars) onboard Mars Reconnaissance Orbiter. The spectral data cover near-infrared wavelengths. We tried to detect the absorption features characteristic to the phyllosilicates. We used CAT (CRISM Analysis Toolkit) combined with ENVI spectral analyst tool to analyze the latest CRISM data observed in the study area in detail.

As a result, phyllosilicates-bearing minerals were detected at 5 impact craters among 14 impact craters examined in this study. The 5 craters are: 2 craters where phyllosilicate hadn't been detected by Fairén et al. [2012], 2 craters not examined so far and 1 crater where phyllosilicates are already detected by Fairén et al. [2012]. The four kinds of phyllosilicates were detected: illite, smectite, vermiculate and a small number of saponite. We saw no discrete distribution according to the kind of minerals but we found that the distribution of phyllosilicates-bearing minerals strongly connects with erosional areas. We observed that phyllosilicates distribute at the rim, wall, floor, ejecta and around the central peak of the craters. These detected phyllosilicates are interpreted that deposited under the younger crust of the northern lowlands are emitted on the surface by impact gardening by Carter et al. [2010].

The number of samples in this study is still small, but the detection ratio of phyllosilicates seems significantly larger than Carter et al. [2013]. The results imply that hydrated minerals are possibly more widespread in the northern lowlands of Mars. Such detailed examination that uses the latest observation data and smart tools should result in increasing ratio of detecting phyllosilicates and contribute to clarification of water environment in the northern lowlands of Mars.

キーワード：火星、含水鉱物、フィロケイ酸塩、赤外分光、CRISM/MRO

Keywords: Mars, hydrated minerals, phyllosilicate, infrared spectra, CRISM/MRO

火星境界層の高解像度LES実験

High resolution Large eddy simulation on Martian planetary boundary layer

*西澤 誠也¹、小高 正嗣²、高橋 芳幸³、杉山 耕一朗⁴、中島 健介⁵、石渡 正樹²、竹広 真一⁶、八代 尚¹、富田 浩文¹、林 祥介³

*Seiya Nishizawa¹, Masatsugu Odaka², Yoshiyuki O. Takahashi³, Ko-ichiro SUGIYAMA⁴, Kensuke Nakajima⁵, Masaki Ishiwatari², Shin-ichi Takehiro⁶, Hisashi Yashiro¹, Hirofumi Tomita¹, Yoshi-Yuki Hayashi³

1.理化学研究所計算科学研究機構、2.北海道大学、3.神戸大学、4.宇宙科学研究所、5.九州大学、6.京都大学
1.RIKEN Advanced Institute for Computational Science, 2.Hokkaido Univ., 3.Kobe Univ., 4.ISAS, 5.Kyushu Univ., 6.Kyoto Univ.

A large-eddy simulation (LES) on Martian planetary boundary layer has been performed to examine structure and characteristics of turbulence in the layer and their dependency on experimental resolution. SCALE-LES, which is an LES model for large domain and high resolution experiments developed in RIKEN AICS (<http://scale.aics.riken.jp/>), is used in this study. Spatial resolution is swept from 100m to 5m. Domain size is about 20km in horizontal and vertical. Horizontal boundary condition is double periodic condition. Heating rate calculated by an offline one-dimensional experiment (Odaka et al. 2001) is used instead of explicit calculation of the radiative transform process.

Well-known features of the boundary layer, such as hexagonal structure of convective cells and the -5/3 energy spectrum, are reasonably simulated. Dependency of several physical quantities at 14:00-15:00 local time, when the boundary layer is almost mature, on the resolution is analyzed. We found that vertical heat flux and variance of vertical velocity of resolved component show convergence or systematic tendency with the resolution. Convective vortices are developed and most of them are located near the upward reasion.

キーワード：火星境界層

Keywords: Martian Planetary Boundary Layer

火星大気北半球冬季におけるCO₂過飽和の経度依存性

Longitudinal dependence of CO₂ supersaturation during northern winter in the Martian atmosphere

*野口 克行¹、森井 靖子¹、小田 尚香¹、黒田 剛史²

*Katsuyuki Noguchi¹, Yasuko Morii¹, Naoka Oda¹, Takeshi Kuroda²

1.奈良女子大学、2.東北大学

1.Nara Women's University, 2.Tohoku University

米国の火星探査機であるマーズ・グローバル・サーベイヤー（MGS）で行われた電波掩蔽観測では、長期間（1998-2007）にわたる気温の高度分布データが得られている。このデータを解析した結果、北半球冬季の高緯度（北緯60-80度）において、大気主成分であるCO₂の過飽和の発生に強い経度依存性が見られることがわかった。CO₂過飽和が頻発するのは、約400-30 Paの高度領域においては東経90-180度と270-360度の地域であった。気温の東西偏差と比較すると、およそ低温部に一致することがわかった。

キーワード：火星、過飽和、二酸化炭素

Keywords: Mars, supersaturation, CO₂

惑星大気大循環モデル DCPAM を用いた火星大気中の水循環の数値シミュレーション

Numerical simulation of water cycle in a Martian atmosphere by the use of a planetary atmosphere general circulation model, DCPAM

*高橋 芳幸¹、林 祥介¹

*Yoshiyuki O. Takahashi¹, Yoshi-Yuki Hayashi¹

1.神戸大学大学院理学研究科

1.Graduate School of Science, Kobe University

Spacecraft observations of Mars revealed distributions of water vapor and water ice cloud in the Martian atmosphere. Those observations covered an almost global region and a long period. For example, the Mars Global Surveyor (MGS) observed column densities of water vapor and water ice cloud for about 9 Martian years. The MGS observation characterizes the seasonal variation of the water distribution in the Martian atmosphere. On the other hand, general circulation models (GCMs) have been used to investigate water cycle in the Martian atmosphere. Those models represented distributions of water vapor and water ice cloud consistent with observed ones, successfully. In our group, we have been developing an atmosphere GCM, DCPAM, which is applicable to planetary atmospheres. By the use of the DCPAM, we have been performed simulations of Martian atmosphere without water cycle. In this study, the simulations of water cycle in the Martian atmosphere are performed by implementing relevant processes in the DCPAM. By performing the simulations, features of water cycle in the Martian atmosphere will be investigated and the model will be validated under a condition of Mars.

The DCPAM used in this study consists of a dynamical core based on the primitive equation system and physical processes relevant to Martian atmosphere. The dynamical core solves the primitive equation system by the use of spectral transform method with the finite difference method in vertical direction. The included physical processes are the radiation, the turbulent mixing, the surface processes, the CO₂ and H₂O condensation, and gravitational sedimentation. In the model, the radius of cloud particles is assumed to be a constant. By the use of a "Mars mode" of this model, several experiments have been performed. In the experiments, the dust distribution in the atmosphere is prescribed. In the vertical direction, the Conrath-type distribution is assumed. In the horizontal direction, the optical depth is prescribed following observations. In order to simulate water cycle, large amount of water ice is placed north of 80N. Further, the surface temperature south of 85S is fixed to 145 K to represent a permanent CO₂ ice cap. Those H₂O and CO₂ ices at southern and northern high latitude regions act as source and sink of the water. The resolutions used for this study is T21L36, which is equivalent to about 5.6 degrees longitude-latitude grid and has 36 vertical levels. Under these conditions, the model is integrated for 10 Mars years from an initial condition of isothermal atmosphere at rest. The result during the last Martian year is analyzed.

The model is evaluated by comparing the column densities of water vapor and water ice cloud simulated by the model with those observed by the MGS. The simulations with ice cloud radius of 7 micron meter show following features of seasonal variation of those values which are roughly consistent with observations. From northern summer to northern fall, the water vapor is transported from northern cap to low latitude. At the same time, water ice cloud forms in northern low latitude where the ascending motion of Hadley circulation occurs. However, the low latitude water ice cloud from northern winter to northern spring is slightly thicker in the model than observed one. This may be caused by crude treatment of ice cloud in the model. In the presentation, features of water cycle represented in the model will be presented in more details.

キーワード：惑星大気、大気大循環モデル、火星、水循環

Keywords: planetary atmosphere, general circulation model, Mars, water cycle

火星大気大循環モデルを用いた表面ダスト巻き上げフラックス診断実験：地形の効果によって受ける影響についての考察

Diagnostic experiments of lifted dust flux at the surface with Mars GCM: Consideration of the effects of topography

*荻原 弘堯¹、高橋 芳幸²、石渡 正樹¹

*Hiroataka Ogihara¹, Yoshiyuki O. Takahashi², Masaki Ishiwatari¹

1.北海道大学大学院理学院宇宙理学専攻、2.神戸大学大学院理学研究科

1.Department of CosmoSciences, Graduate school of Science, Hokkaido University, 2.Graduate School of Science, Kobe University

火星ではダストが大気熱構造に影響を与えている（例えば、Liu et al., 2003）。各国の研究グループで火星のダスト循環を大気大循環モデルで再現しようという試みが行われている（例えば、Newman et al., 2002, Basu et al., 2004, Kahre et al., 2006）。我々もこれまで開発を進めてきた大気大循環モデル DCPAM（高橋他, 2014）へモデルで直接表現できる風応力によるダスト巻き上げと直接表現できない対流渦（ダストデビル）によるダスト巻き上げを実装しダスト巻き上げ実験をおこなった（荻原他, 2014）。そこで得られたダスト巻き上げフラックスの特徴は先行研究と大まかに一致した。しかし、ダスト巻き上げパラメタリゼーション、特にダストデビルによるダスト巻き上げパラメタリゼーションの振る舞いは地形によって複雑な挙動を示し、分布がどのように決まるかまでは理解することはできなかった。そのために本研究では、地形がどのようにダストデビルによるダスト巻き上げパラメタリゼーションの振る舞いへ影響を及ぼすか理解することを目指す。それゆえに単純化した地形で考える。ここでは、水平一様な地形と東西平均された地形という二つの単純な地形を用いてダスト巻き上げ実験を行い、その結果を比較する。

ここで用いるモデルは DCPAM である。DCPAM では 3 次元球面プリミティブ方程式を用いている。放射過程では CO₂ と大気ダストを考慮し、Takahashi et al.(2003, 2006) のスキームを用いる。放射スキームに与えるダスト分布は固定したものをを用いる。乱流過程は Mellor and Yamada (1982) の方法に従って決めた鉛直拡散係数を用いて評価する。地表面過程は Beljaars and Holtslag (1991), Beljarrs (1994) の方法に従って評価する。ダストデビルによるダスト巻き上げスキームとして DDA スキーム (Newman et al., 2002) を用いた。このスキームはダストデビルを熱エンジンと見なし、顕熱フラックスと対流層の厚さに依存する熱力学効率を用いてダスト巻き上げ量を決める。水平離散化にはスペクトル法を用い全波数 21 で打ち切りを行う、鉛直方向には差分法を用い、鉛直層数は 36 とする。積分時間は 4 火星年であり、最後の 1 火星年分の解析を行った。北緯 25 度と南緯 25 度の二つの緯度領域について調べる。共に、ダスト巻き上げフラックスが増加するそれぞれの半球における春から夏の季節に着目する。地表面熱慣性、地表面アルベドは固定した。地形高度として水平一様な地形を用いた場合（実験 F）とグローバルサーベイヤーの観測結果を東西平均した地形を用いた場合（実験 Z）で、ダストデビルによるダスト巻き上げパラメタリゼーションのダスト巻き上げ実験を行い、それぞれの結果を比べてみた。

まず、北緯 25 度付近の場合の結果を述べる。東西平均したダスト巻き上げフラックスを比べてみると、実験 Z の場合は 実験 F の場合のよりも小さくなった。熱収支の解析から、実験 Z の場合は 実験 F の場合に比べて下層で対流調節が余り働かず上層を加熱できなく、子午面循環が弱められる。それにより、大気が安定状態になり顕熱フラックスが小さくなったことによってダスト巻き上げフラックスが小さくなった。

次に南緯 25 度付近の場合の結果を述べる。同様に、東西平均したダスト巻き上げフラックスを比べてみると、実験 Z の場合は 実験 F の場合に比べて大きくなった。熱収支の解析から、実験 Z の場合は 実験 F の場合に比べて下層の対流調節が良く働き上層を加熱し、子午面循環が強くなり、対流層が厚くなった。よって、ダスト巻き上げフラックスが大きくなった。

これらの結果から、どの季節でも対流調節が上層を加熱し、子午面循環が強められるとダストデビルによるダストの巻き上げが大きくなると思われる。

今後は観測された地形における東西方向の起伏の違いによるダストデビルのダスト巻き上げフラックスへの影響を考察する予定である。

キーワード：火星、ダスト、大気大循環モデル、ダストデビル、ダスト巻き上げ

Keywords: Mars, Dust, General Circulation Model, Dust Devil, Dust lifting

高分解能大気大循環モデルを用いた火星大気重力波の励起と伝播の評価

Assessment of the generation and propagation of the gravity waves in the Martian atmosphere using a high-resolution general circulation model

*黒田 剛史¹、Alexander Medvedev²、Erdal Yiğit³、Paul Hartogh²

*Takeshi Kuroda¹、Alexander S. Medvedev²、Erdal Yigit³、Paul Hartogh²

1.東北大学大学院理学研究科 地球物理学専攻、2.マックスプランク太陽系研究所、3.ジョージメイソン大学
1.Department of Geophysics, Tohoku University, 2.Max Planck Institute for Solar System Research,
3.George Mason University

Gravity waves (GWs) are small-scale atmospheric waves generated by various geophysical processes, such as topography, convection, and dynamical instability. On Mars, several observations and simulations have revealed that GWs strongly affect temperature and wind fields in the middle and upper atmosphere. Our previous study using the Max Planck Institute Mars General Circulation Model (MGCM) and the nonlinear spectral whole atmosphere parameterization of small-scale GWs by Yigit et al. [2008] have shown that the dynamical forcing of GWs significantly change the winds, reversing its direction above ~100 km [Medvedev et al., 2011]. We also have shown that the thermal effects induced by GWs can be the main source of cooling above ~120 km, reproducing the observed temperature structure on Mars [Medvedev and Yigit, 2012]. Similar physical importance of GWs has previously been demonstrated for the general circulation of Earth's upper atmosphere using the whole atmosphere parameterization [Yigit et al., 2009; Yigit and Medvedev, 2009]. Despite numerous observations however, the global picture of GW activity is yet to be revealed both on Earth and Mars.

In order to investigate the global distribution of small-scale GWs in the Martian atmosphere, we have conducted the first simulations with a high-resolution MGCM, using the DRAMATIC (Dynamics, RAdiation, MAterial Transport and their mutual InteraCtions) MGCM [e.g., Kuroda et al., 2005, 2013]. The MGCM was run at the T106 spectral truncation, which corresponds approximately to a $1.1^\circ \times 1.1^\circ$ (or ~60 km) horizontal resolution. In the vertical direction, the model domain extends from the surface to ~80-100 km and is represented by 49 sigma-levels. Such setup allows for realistically capturing generation and propagation of GWs with horizontal wavelengths of ~180 km and longer and, to some extent, their vertical attenuation due to nonlinear processes. We considered horizontal-scale fluctuations with a total wave number of larger than 60 (horizontal wavelengths of less than ~350 km) as GW-induced disturbances.

We investigated the spatial distributions of potential and kinetic energies associated with GW activity in the northern winter solstice. The simulated GW potential energy distribution is in a good agreement with available radio occultation data [Creasey et al., 2006] in the lower atmosphere between 10 and 30 km. The model reveals a latitudinal asymmetry with stronger wave generation in the winter hemisphere, and investigations from the ratio of potential and kinetic energies show that there are two distinctive sources of GWs: mountainous regions and the meandering winter polar jet. Orographic GWs are filtered upon propagating upward, and the mesosphere is primarily dominated by waves with faster horizontal phase velocities. Wave fluxes are directed mainly against the local wind, with a clear relation between wave dissipation and wind acceleration. GW dissipation in the upper mesosphere generates a body force per unit mass of tens of m s^{-1} per Martian solar day (sol^{-1}), which tends to close the simulated jets. Effects of horizontal propagation of GWs on the acceleration are much smaller than those of vertical propagation, and the results of acceleration rates are comparable to those obtained from the application of the GW parameterization by Yigit et al. [2008], which considers only the vertical propagations of a broad spectrum of GWs.

The results represent a realistic surrogate for missing observations, which can be used to further constrain existing GW parameterizations and validate GCMs. Also the observational investigations of GW signatures in the thermosphere by the MAVEN mission would help better understand propagation and dissipation mechanisms of GWs.

キーワード：火星、大気力学、重力波、大気大循環モデル、MAVEN

Keywords: Mars, Atmospheric dynamics, Gravity waves, General circulation model, MAVEN

Fluorescence Life-Time (FLiT) instrument for space missions

Fluorescence Life-Time (FLiT) instrument for space missions

*佐藤 毅彦¹、山岸 明彦²、宮川 厚夫²、佐々木 聡³、藪田 ひかる⁴、癸生川 陽子⁵、小林 憲正⁵

*Takehiko Satoh¹, Akihiko Yamagishi², Atsuo Miyakawa², Sasaki Satoshi³, Hikaru Yabuta⁴, Yoko Kebukawa⁵, Kensei Kobayashi⁵

1.宇宙航空研究開発機構・宇宙科学研究所、2.東京薬科大学、3.東京工科大学、4.大阪大学、5.横浜国立大学
1.Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 2.Tokyo University of Pharmacy and Life Science, 3.Tokyo University of Technology, 4.Osaka University, 5.Yokohama National University

While we detail the design of the Life-Detection Microscope (LDM), a high spatial resolution fluorescence microscope, alternative ways of detecting extraterrestrial life have been under consideration. One of such is the "fluorescence life-time (FLT)" measurement. FLT has been widely used in biophysical studies of proteins as FLT by its nature is one of the most robust fluorescence parameters. To examine the possibility of applying the FLT measurement to the space missions, we have developed a Bread Board Model (BBM), called FLiT. The method we use is the "time-domain" method in which the sample is illuminated with a short pulse laser and the decay time of the fluorescence is measured.

The pulse laser of FLiT is a 488-nm laser diode (Nichia NDS4116) to which pulses of 1-ns full width at half maximum is fed by the driver electronics. One pulse may excite just one fluorophore and the photon from it will be detected by an avalanche photo diode (APD) in the photon-counting mode (MPPC C13001-01 from Hamamatsu Photonics). The time delay from the start trigger (1-ns pulse) to the stop trigger (photon detection in MPPC) is measured by the time-to-digital converter (TDC7200 from TI). Such measurement will be repeated and a histogram of delays is obtained from which the fluorescence life-time of the sample material is inferred. This potentially allows us to distinguish organisms from minerals in the Martian soil. The FLiT BBM is now under the characterization phase, to evaluate possible delays in the electronics (including cables) and their stability. Details of the FLiT BBM as well as results of initial tests will be presented.

キーワード：蛍光寿命、火星生命、生命探査顕微鏡

Keywords: Fluorescence life-time, Martian life, Life-detection microscope