

## 東京湾不整合と万田野寒冷期

## The Tokyo Bay Unconformity and the Mandano Ice- Age

\*楡井 久<sup>1</sup>、風岡 修<sup>2</sup>、木村 英人<sup>3</sup>、吉田 剛<sup>2</sup>、藤田 寛、楡井 亘<sup>4</sup>

\*hisashi nirei<sup>1</sup>, Osamu Kazaoka<sup>2</sup>, Hideto Kimura<sup>3</sup>, Takeshi Yoshida<sup>2</sup>, Hiroshi Fujita, Wataru Nirei<sup>4</sup>

1.国際地質科学連合環境管理研究委員会、2.千葉県地質環境研究室、3.東邦地水株式会社、4.東京情報大学

1.International Union of Geological Science for Environmental Management, 2.Research institute of Environmental Geology, Chiba, 3.Tohochisui Co., 4.Tokyo university of information science

The Kanto fore-arc basin in the Kato plain is an extremely deep submarine basin that was formed during the early Pleistocene. The sediments in the basin change from deep-sea sediments to lacustrine-alluvial sediments, the Mandano formation which is up to 95 m in thickness and overlays the Tokyo Bay unconformity that is widely distributed under Tokyo bay area. The formation consists of three parts. The lithofacies in the lower part on the unconformity gradually change, with decreasing depth, from sand to gravel. The lithofacies of the middle part are muddy. The upper part changes with increasing depth from gravel to sandy silt by transgression. The lower part and lower half of the middle part comprise sediments characteristic of a topset fan delta (Nirei H., 1997) in the regression stage. The upper half of the middle part and upper parts are composed of transgression sediments. The lower half of the middle part contains sediments from the ice age regression stage, evidence of which is provided by the cold-index plant remains, *Picea maximowiczii*, *Tsuga diversifolia*, *Fagus crenata*, *Cryptomeria japonica*, etc., present in the uppermost part of the lower half. Geological analysis reveals the Kanto continental shelf to extend widely under the Kanto Plain; it is also underlaid by a range of bottom set sediments in distinct formations and forest sediments, also in distinct formations, in ascending order under the conformity.

キーワード：東京湾不整合、万田野寒冷期

Keywords: Tokyo bay unconformity, Mandano ice-age

## 千葉セクションにおける松山-ブルン極性反転境界記録

Revised Matuyama-Brunhes polarity transition record from a marine succession at the Chiba section, a Lower-Middle Pleistocene GSSP candidate

\*岡田 誠<sup>1</sup>、菅沼 悠介<sup>2,3</sup>、羽田 裕貴<sup>1</sup>、風岡 修<sup>4</sup>

\*Makoto Okada<sup>1</sup>, Yusuke Suganuma<sup>2,3</sup>, Yuki Haneda<sup>1</sup>, Osamu Kazaoka<sup>4</sup>

1.茨城大学理学部理学科、2.国立極地研究所、3.総合研究大学院大学極域科学専攻、4.千葉県環境研究センター

1.Department of Earth Sciences, College of Science, Ibaraki University, 2.National Institute of Polar Research, 3.Department of Polar Science, The Graduate University for Advanced Studies, 4.Research Institute of Environmental Geology

We report revised paleomagnetic records of the Matuyama-Brunhes boundary (MBB) from a continuous marine succession at the Chiba section of the Kokumoto Formation, Kazusa Group. The Chiba section is the one of the candidate sites for the Lower-Middle Pleistocene Boundary GSSP. In the section, a wide spread tephra bed named as Byk-E is intercalated just 80 cm below the MBB. In order to provide globally comparable VGP (virtual geomagnetic pole) and paleointensity (past geomagnetic field intensities) records from the Chiba section, we have taken oriented mini-cores from a 13 meters succession with 10-cm intervals across the Byk-E tephra bed. Thermal magnetic experiments suggest that the samples include iron sulfides, magnetites but no hematite. Measurements of magnetic hysteresis indicate that the magnetic domain state is PSD. Progressive alternating field demagnetization (AFD) indicate a reversed to normal polarity transition boundary is at around 1.5 meter below the Byk-E bed as well as previous studies, however the transition boundary is observed at around 0.8 meter above the Byk-E bed in thermal demagnetization (ThD) results. Therefore, the reversed to normal polarity transition boundary seen below the Byk-E bed is thought to be overprint. This overprint, which might be carried by iron sulfide, is particularly observed in a transitional interval. Since iron sulfides generally decompose and oxidized into magnetites due to heating during ThD, the yielded magnetites have no magnetic signal but provide an over estimate of magnetic grain amount which prevents to estimate paleointensities. To provide a reliable paleointensity record, we applied to use a composite demagnetization technique consisting of a 300°C ThD and a regular progressive AFD sequence. After the 300°C ThD, most of the overprint has been removed but the magnetic susceptibility has not changed even in the air condition, indicating that iron sulfides just lose magnetic signals due to the ThD but not to change the amount of magnetic grains. The VGP latitudes and preliminary derived paleointensities using the composite demagnetization technique from the Chiba section quite match well with the U1308 records. To use the both independent techniques of oxygen isotope and paleointensity will provide a further reliable stratigraphic correlation across the Lower-Middle Pleistocene Boundary.

キーワード：古地磁気学、古地磁気強度、国際標準模式層断面および地点

Keywords: Paleomagnetism, Paleointensity, GSSP

千葉県銚子より掘削された陸上ボーリングコアを用いたMIS 19における高解像度酸素・炭素同位体記録（予察）

A preliminary report for high-resolution foraminiferal oxygen and carbon stable isotope records in MIS 19 from an on land core drilled at the Choshi city, central Japan.

早田 達哉<sup>1</sup>、\*羽田 裕貴<sup>2</sup>、岡田 誠<sup>1</sup>、久保田 好美<sup>3</sup>

Tatsuya Hayata<sup>1</sup>, \*Yuki Haneda<sup>2</sup>, Makoto Okada<sup>1</sup>, Yoshimi Kubota<sup>3</sup>

1.茨城大学理学部理学科、2.茨城大学大学院理工学研究科、3.国立科学博物館

1.Department of Earth Sciences, Faculty of Science, Ibaraki University, 2.Graduate School of Science and Engineering, Ibaraki University, 3.National Museum of Nature and Science

The Plio-Pleistocene Inubo Group, distributed in the Choshi city, Chiba prefecture, central Japan, is thought to be a suitable marine succession to investigate paleoceanographic and paleoclimatic changes around the northwestern Pacific Ocean, because a lot of wide spread key tephra beds are intercalated, and microfossils and pollens are abundant. In 1998, a continuous, well recovered on land core drilled through the Obama, Yokone, Kurahashi and Toyosato Formations in the Inubo Group was obtained (after Choshi core). Kameo et al. (2006) studied calcareous nannofossil, paleomagnetic and planktonic foraminiferal oxygen isotope stratigraphies of the Choshi core, and reported that the core corresponded to a period between MIS 11 and 24 based on a correlation with the LR04 stack curve (Lisiecki and Raymo, 2005). In this study, we show a new high-resolution stable isotope record using benthic foraminifers from a section across the Lower-Middle Pleistocene boundary of the Choshi core. This record corresponds to MIS 18-20 with a time resolution of ca. 500 years. The average oxygen isotopic value of the Choshi core is about 0.5 ‰ lighter than that of LR04 during the period of MIS 19, and the difference becomes larger as the age becomes younger, indicating that the accumulation depth of the Choshi core was getting shallower due to uplifting and/or burring up the basin. Further analysis on the core will show some paleoceanographic findings at around the north western Pacific margin during the MIS 19 period.

キーワード：MIS 19、有孔虫安定同位記録、銚子コア

Keywords: MIS 19, Foraminiferal stable isotope record, Choshi core

房総半島中部に分布する上総層群下部-中部更新統の石灰質ナノ化石層序と古海洋環境  
Calcareous nannofossil biostratigraphy of the Lower-Middle Pleistocene in the Kazusa  
Group, central part of the Boso Peninsula, and estimated sea surface environments

\*渡辺 賢人<sup>1</sup>、亀尾 浩司<sup>1</sup>

\*Kento Watanabe<sup>1</sup>, Koji Kameo<sup>1</sup>

1. 千葉大学理学研究科地球生命圏科学専攻地球科学コース

1. Department of Earth Sciences, Division of Geosystem and Biosystem Sciences, Graduate School of  
Science, Chiba University

本邦中部太平洋側に位置する房総半島には第四系の海成堆積物から構成される上総層群が分布する。露頭条件が良いことや化石を多く含むことから、岩相層序学的研究や古地磁気層序学、テフラ層序学、微化石層序学のような年代層序学的な研究が行われてきた（新妻, 1976; 佐藤ほか, 1988; 五十嵐, 1994など）。このうち下部-中部更新統に相当する国本層には、ブリュンヌ/松山境界（Matuyama/Brunhes Boundary: MBB）が記録されており、GSSP（Global Boundary Stratotype Section and Point）模式地の有力な候補のうちの一つとなっている（Kazaoka et al., 2015）。そのため、前期-中期更新世の正確な年代を決定するために、より詳細な年代層序の検討が必要であると言える。ところが、国本層に産出する石灰質ナノ化石は、必ずしも高い時間解像度で解析されているわけではない（佐藤ほか, 1988）。そのため、当層準における石灰質ナノ化石層序は十分に検討されておらず、化石基準面も明らかにされていない。よって本研究ではMBB相当層準の石灰質ナノ化石基準面を明らかにするとともに、群集変化に基づいて同層準堆積当時の北西太平洋の表層海洋環境を推定するために、国本層の石灰質ナノ化石の検討を行った。検討した66試料からは、10属19種の石灰質ナノ化石が産出した。検討層準を通じて種組成に変化はないが、そのうち、小型の*Gephyrocapsa*属が多産する層準がMBB直上に位置し、この多産イベントは顕著である。この層準のイベントはイタリア南部のMontalbano JonicoやValle di Mancheでも指摘されていることから（Girone et al., 2013）、グローバルに追跡可能なイベントである可能性がある。さらに、寒冷な環境指標種である*Coccolithus pelagicus*や温暖な環境指標種である*Umbilicosphaera sibogae*の産出には逆相関する関係が見られた。層位分布をみると*Umbilicosphaera sibogae*が全層準を通して増減を繰り返す傾向が見られることから、黒潮水域と混合水域の北上、南下が交互に起きていたと考えられる。

#### 引用文献

Girone et al., 2013, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 371, 62-79.

<http://www.elsevier.com>

五十嵐, 1994, *地質学雑誌*, 100, 348-359

Kazaoka et al., 2015, *Quaternary International*, 383, 116-135.

<http://dx.doi.org/10.1016/j.quaint.2015.02.065>

新妻, 1976, *地質学雑誌*, 82, 163-181

佐藤ほか, 1988, *石油技術協会誌*, 53, 475-491

## 更新統下部—中部境界を含む国本層上部の詳細層序と堆積環境：千葉セクション

Detailed litho-stratigraphy and sedimentary environment of upper part of Kokumoto Formation with the L-M Pleistocene boundary: the Chiba section, Central Japan

\*風岡 修<sup>1</sup>、荻津 達<sup>1</sup>、八武崎 寿史<sup>1</sup>、本田 恵理<sup>1</sup>、吉田 剛<sup>1</sup>、亀山 瞬<sup>2</sup>、香川 淳<sup>1</sup>、森崎 正昭<sup>1</sup>、竹下 欣宏<sup>3</sup>、里口 保文<sup>4</sup>、中里 裕臣<sup>5</sup>、西田 尚久<sup>6</sup>、岡田 誠<sup>7</sup>、菅沼 悠介<sup>8</sup>、泉 賢太郎<sup>9</sup>、熊井 久雄<sup>10</sup>、楡井 久<sup>11</sup>

\*Osamu Kazaoka<sup>1</sup>, Itaru Ogitsu<sup>1</sup>, Hisashi Yabusaki<sup>1</sup>, Eri Honda<sup>1</sup>, Takeshi Yoshida<sup>1</sup>, Syun Kameyama<sup>2</sup>, Atsushi Kagawa<sup>1</sup>, Masaaki Morisaki<sup>1</sup>, Yoshihiro Takeshita<sup>3</sup>, Yasuhumi Satoguchi<sup>4</sup>, Yasuomi Nakazato<sup>5</sup>, Naohisa Nishida<sup>6</sup>, Makoto Okada<sup>7</sup>, Yusuke Suganuma<sup>8</sup>, Kentaro Izumi<sup>9</sup>, Hisao Kumai<sup>10</sup>, Hisashi Nirei<sup>11</sup>

1.千葉県環境研究センター地質環境研究室、2.千葉県環境生活部、3.信州大学教育学部、4.琵琶湖博物館、5.農業工学研究所、6.産業技術総合研究所地質調査総合センター、7.茨城大学、8.国立極地研究所、9.国立環境研究所、10.大阪市立大学、11.GEM-IUGS日本支部

1.Research Institute of Environmental Geology, Chiba, 2.Environmental Protection division of Chiba Prefectural Government, 3.Institute of Education, Shinshu University, 4.Lake Biwa Museum, 5.National Institute for Rural Engineering, 6.Geological Survey of Japan, AIST, 7. Ibaraki University, 8. National Institute of Polar Research, 9.National Institute for Environmental Studies, 10.Osaka City University, 11.Japan Branch of GEM-IUGS

The Lower -Middle Pleistocene Kazusa Group, deposited on mainly bathyal -shelf in the Pacific Ocean with many kind of fossils, distributes widely in Boso peninsula. The Kazusa Group exposes continuously along Yoro river, Chiba section, type section of the group. The Kazusa Group consists of Kurotaki Formation (mainly tuffaceous gravelly sandstone), Katsuura F. (mainly alternation of sandstone with slump bed), Namihana F. (mainly siltstone with slump bed), Ohara F. (muddy alternation of sandstone and siltstone), Kiwada F. (muddy alternation of sandstone and siltstone with slump bed), Otadai F. (alternation of sandy alternation and muddy alternation of sandstone and siltstone), Umegase F. (mainly sandy alternation of sandstone and siltstone), Kokumoto F. (alternation of thick siltstone and sandy alternation of sandstone and siltstone), Kakinokidai F. (sandysiltstone with sandstone) and Chonan F. (alternation of thin sandstone and thin siltstone) in ascending order (Mitsunashi et al., 1959). Total thickness of the Kazusa Group is over 2,000 meters with over 50 marker tephra beds. Trepid depositional rates of 2.0-2.5 m/ky are obtained for the Kazusa Group. Therefore the Chiba section have high potential for international stratotype section (Kazaoka et al., 2015).

Kokumoto Formation, about 350 meter thick, is composed of the lowermost part, the lower part, the upper part and the uppermost part in ascending order. The lowermost part, about 60 meter thick, consists of thick siltstone with thin sandstone bed and marker tephra, Ku6 and ku5. The lower part, about 120 meter thick, consists of sandy alternation of sandstone and siltstone with Ku3 tephra. The upper part, about 80 meter thick, consists of thick siltstone with thin sandstone and marker tephra (Byakubi zone (Byk-G, Byk-F, Byk-E, Byk-D, Byk-C, Byk-B and Byk-A), Koss2, Koss1-B, Koss1-A, Kosp-C, Kosp-B, Kosp-A, Tap-B, Tap-A, Tas-C, Tas-B, Tas-A, Ku2 and Ku1). Especially maker tephra are interbedded every 0.1-7.0 ky in the thick siltstone from Byk-E to Ku2 horizon. The Matuyama-Brunhes boundary is between Byk-C and Byk-B. Uppermost part, about 90 meter thick, consists of sandy alternation of sandstone and siltstone with Ku0.1 tephra.

The upper part, thick siltstone, is interbedded with thin, 1-3cm thick, sandstone every 0.3-3 m thick and thin, 1-5 cm thick, sandysiltstone every 0.1-0.25 m thick without slump bed and thick mudflow bed. The siltstone have bathyal and sublittoral benthic foraminifera and many trace fossils. Grain size distribution in the siltstone have bimodal grain group (Nishida et al., 2015). Main grain group is composed of fine silt and sub group consists of very fine sand. These

characteristics show hemipelagic sedimentary environment in deep sea and very fine sand flow often into, namely deep sea slope. The thickness from Byk-G to Byk-A change little laterally in the central part of Boso Peninsula. This show that the fracks deposited approximately uniformly. It is presumed that clastic sediments were supplied little around here from source mountain area, because this horizon is warm stage, MIS 19.

キーワード：更新統下部－上部境界、国本層、上総層群、火山灰鍵層、Byk-E、MIS 19

Keywords: L-M Pleistocene boundary, Kokumoto Formation, Kazusa Group, Maker Tephra, Byk-E, MIS 19