

B2型MgOの格子拡散挙動

Lattice diffusion in B2-type MgO

*原田 隆史¹、土屋 卓久¹*Takafumi Harada¹, Taku Tsuchiya¹

1.愛媛大学地球深部ダイナミクス研究センター

1.Geodynamics Research Center, Ehime University

High-pressure and high-temperature rheology is essential for understanding the dynamics in planets. Diffusion creep might be one of the dominant viscoplastic mechanisms and lattice diffusion coefficient D is a key property in this deformation process (e.g., Karato, 2011). Experimental measurements of D however still remain technically difficult under deep planetary conditions. Theoretical approaches therefore play a substantial role.

B2 (CsCl)-type MgO is a high-pressure phase of B1 (NaCl)-type MgO and expected to be one of the major constituents in super-Earths' mantle and giant planetary core (Guillot, 1999; Tsuchiya and Tsuchiya, 2011). Although diffusion creep viscosity of super Earths' mantle is usually assumed to increase with depth monotonically, a previous study (Karato, 2011) suggested that it could decrease associated with the B1-B2 transition of MgO. However, this idea is obtained based on measured plasticity of analog materials and thus D of actual B2-type MgO is still underdetermined.

In this study, we calculate D of B1- and B2-type MgO based on first principles constant-temperature molecular dynamics method combined with static lattice energy calculations. We identify distinctly larger D of both Mg and O in B2-type than in B1-type at the same pressure, suggesting that B2-type would be less viscous as expected. The mechanisms of increase in D will be presented.

キーワード：格子拡散、第一原理計算、スーパーアース

Keywords: Lattice diffusion, First principles calculation, Super Earth

転位クリープ変形下における含水かんらん岩の組織発達と変形特性

Microstructural development in olivine aggregates during dislocation creep under hydrous conditions

*田阪 美樹^{1,2}、Mark Zimmerman²、Kohlstedt David²

*Miki Tasaka^{1,2}, Mark Zimmerman², David Kohlstedt²

1.新潟大学、2.ミネソタ大学

1.Niigata University, 2.University of Minnesota

Since hydrogen plays an important role in dynamic processes in the mantle, we conducted high-strain torsional shear experiments on aggregates of Fe-bearing olivine [(Mg,Fe)₂SiO₄; Fo50] under hydrous condition. Olivine with a composition of Fo50 was used because of its enhanced grain growth kinetics and low strength relative to Fo90. Two pieces of an oriented San Carlos olivine crystal were embedded in each aggregate to monitor water fugacity both before and after deformation. We deformed samples to high enough shear strain ≈ 5 , to achieve a steady-state microstructure.

A non-linear, least-squares fit to the stress versus strain rate data yielded a stress exponent of $n \approx 3.5$, indicative of deformation involving dislocations. The water content determined from Fourier transform infrared (FTIR) spectroscopy analyses of the single crystals demonstrated that the samples were water saturated after deformation. Fabric analyses of the polycrystalline olivine samples, determined using electron backscatter diffraction (EBSD), indicate that the strength of the lattice preferred orientation (LPO) increases with increasing strain. Further, the LPO of olivine changes as a function of strain due to competition among three slip systems: (010)[100], (100)[001], and (001)[100]. Observed strain weakening can be attributed to geometrical softening due to LPO development, which reduces the stress by $\sim 1/3$ from its peak value in constant strain rate experiments. The evolution of fabric can be applied to investigations of upper mantle seismic anisotropy especially in a mantle wedge or in a shear zone, locations in which hydrous conditions prevail.

キーワード：かんらん石、高歪み変形実験、転位クリープ、結晶配向形成による歪み弱化

Keywords: olivine, high strain deformation, dislocation creep, geometrical softening due to LPO development

下部マントル条件下におけるMORBとパイロライト物質の多相粒成長実験

Differences in grain growth kinetics between MORB and pyrolitic materials under lower mantle conditions: preliminary results

*久保 友明¹、今村 公裕¹、加藤 工¹

*Tomoaki Kubo¹, Masahiro Imamura¹, Takumi Kato¹

1.九州大理

1.Kyushu University

Convective mixing and persistence of chemical heterogeneities such as subducting slabs in the lower mantle largely depends on their density and viscosity contrasts. In contrast to detailed studies on the density, those on the viscosity of deep slab materials have been limited so far due to difficulties of direct quantitative deformation experiments. Previous studies suggest that both MORB and peridotite regions of deep slabs across the upper and lower mantle boundary cause significant grain-size reduction through the post-spinel and post-garnet transformations, respectively, resulting that the grain-size sensitive creep becomes dominant as the deformation mechanism (e.g., Kubo et al., PEPI2008; EPSL2009; AGU2011). Therefore, the grain growth is an important process controlling the viscosity of slab materials in the lower mantle. Here we report preliminary results on grain growth experiments of MORB and pyrolitic materials under lower mantle conditions.

Each material consists of 3-4 phases after the transformations; those are Mg-perovskite, Ca-perovskite, stishovite, and aluminous phase in MORB, and Mg-perovskite, Ca-perovskite, and ferropericlasite (+majoritic garnet at the top of the lower mantle) in pyrolite. We conducted grain growth experiments in these assemblages using a Kawai-type multi-anvil apparatus at ~25-28 GPa, 1873-2373K, and for 1-600 min. SEM observations of recovered samples revealed that these assemblages exhibit relatively homogeneous equi-granular texture except for the short-duration annealing in the pyrolitic material. At the present stage, the average grain size was measured without distinction among phases in the case of MORB material, whereas the grain size in each phase was measured for the pyrolitic material. Preliminary analysis on the grain growth data indicates that the grain growth exponent is about 3.5 for both the MORB material and the major phase of Mg-perovskite in the pyrolitic material, suggesting that the grain growth kinetics in these multi-phase assemblages are controlled by an Ostwald ripening process. The grain size in the MORB material is smaller than that in the pyrolitic material, which corresponds to the difference of 100-150K in temperature. The difference in grain size evolution may lead to the viscosity contrast between MORB and pyrolitic materials in the lower mantle, however further detailed studies are needed to assess this issue.

In-situ X-ray diffraction measurement of growth kinetics of reaction rim at high pressures
In-situ X-ray diffraction measurement of growth kinetics of reaction rim at high pressures

岩澤 圭希¹、*西 真之¹、入船 徹男¹、肥後 祐司²

Yoshiki Iwazawa¹, *Masayuki Nishi¹, Tetsuo Irifune¹, Yuji Higo²

1.愛媛大学地球深部ダイナミクス研究センター、2.高輝度光科学研究センター

1.Geodynamics Research Center, Ehime University, 2.Japan Synchrotron Radiation Research Institute

Diffusion rates of elements in minerals provide important constraints for understanding many physical and chemical processes in the Earth's interior, including mantle rheology and chemical transportation. Therefore, many researchers showed the experimental studies on the growth kinetics of reaction rims between minerals, which are often controlled by diffusion of elements. Most of the previous studies succeeded to determine the rim growth kinetics based on the direct measurement of the thickness of reaction layers of recovered samples. However, it is difficult to obtain the precise growth kinetics in this method due to the small uncertainty of temperature and water contents during each experiment.

Here we use in situ X-ray measurements in conjunction with a multi-anvil apparatus to obtain the precise kinetic data of the rim growth of MgAl_2O_4 spinel between MgO periclase and Al_2O_3 corundum. Time resolved X-ray diffraction patterns enable us to obtain the information with the constant temperature, pressure, and water contents.

We succeeded to obtain the time-resolved X-ray diffraction data during the rim growth of MgAl_2O_4 spinel. However, the obtained growth kinetics contains large uncertainty because of the grain growth of the sample minerals due to high water contents. Further experimental improvement would be required to obtain the precise growth rate of the reaction rims.

キーワード：反応帯成長速度、マントル、マルチアンビル型装置

Keywords: rim growth kinetics, mantle, multi-anvil apparatus

金属メルトのマントルレオロジーに対する効果

Effect of the metallic melt on the mantle rheology

*岡本 篤郎¹、平賀 岳彦¹*Atsuro Okamoto¹, Takehiko Hiraga¹

1. 東京大学地震研究所

1. Earthquake Research Institute, The University of Tokyo

コア - マントル境界域では、主にポストペロブスカイト相からなるマントル物質に外核物質である鉄合金メルトが混ざった状態であると推定されている。しかしながら、マントル物質とコア物質の化学的・力学的相互作用についてはよくわかっていないことが多い。本研究では、こうした問題を考える上で重要な、金属メルトがマントル物質の流動に与える影響を実験により検証した。

Hirth & Kohlstedt (2003) によれば、一定応力下でのメルトを含む多結晶体のクリープ速度はメルトの体積分率の指数関数で表わされ ($d\epsilon/dt \propto \exp(\alpha\phi)$)、メルト分率が多いほど柔らかい。濡れ性の良い玄武岩質メルトでは、高い α 値を取るのに対して (Scott & Kohlstedt, 2006)、Hustoft et al. (2007) は濡れ性の悪い金や硫化鉄の入ったオリビン多結晶体の変形実験を行い、低い α 値を得た。メルトと固相の濡れ性が α 値を決めていることが分かっているものの、結晶に対するメルトの相対的な大きさや歪の効果は未だ不明である。本研究では、Hustoft et al. (2007) と同様に、オリビン多結晶体に金を混ぜた物質の変形実験を行い、その粘性率と先行研究のそれとを比較することで、これらの因子がマントル物質のレオロジーにどのような影響を与えるのかを調べた。

水酸化マグネシウム、二酸化ケイ素及び金の微粒粉を混合し、固相反応からフォルステライト+エンスタタイトを合成し、得られた鉱物粉を真空中で焼結することで金入りオリビン多結晶体を作製した。フォルステライト、エンスタタイト、金の体積分率はそれぞれ81 %, 9 %, 10 %であった。また、オリビンの平均粒径は0.7 μ m、金の平均粒径は0.8 μ mであった。この試料を用いて、大気圧下で一軸圧縮の高温クリープ実験を行った。一定温度下(1200°C・1300°C)で 10 MPa, 20 MPa, 40 MPa, 80 MPaと階段状に応力を変化させ、その時の歪速度を測定した。時間と歪の関係が線形になったところを試料の定常クリープとみなし、その際の歪速度を得た。実験後、回収試料の微細構造を走査型電子顕微鏡によって観察した。

得られた応力-歪み速度の関係から、 $d\epsilon/dt \propto \sigma^{1.7}$ を得た。また、等方形状をした金粒子が、実験後に圧縮方向に対し垂直につぶれていることを観察した。また、金を含まないフォルステライト+エンスタタイト試料のクリープ強度と比較すると、本研究の試料は4 ~ 8倍柔らかかった。さらに、Hustoftの結果と比較しても、同じメルト分率で本研究の試料が 3 ~ 4倍柔らかかった。

Hustoftの実験では変形による金粒子の変形が見られないのに対し、本研究では、金の著しい変形が観察された。本研究の試料が柔らかかったことは、オリビンに対する金メルトの粒径が大きいために金メルトが大きく変形し、それに伴ってオリビンにかかる応力が増大したことで説明できるかもしれない。

キーワード：粘性率、金属メルト、オリビン

Keywords: viscosity, metallic melt, olivine

超音波パルスエコー法を用いた15GPaまでのFe-S融体の音速測定手法の開発

Technical development and improvement for sound velocity measurements of liquid Fe-S up to 15 GPa using ultrasonic pulse-echo method

*西田 圭佑¹、鈴木 昭夫²、柴崎 裕樹³、若林 大佑⁴、船守 展正⁴

*Keisuke Nishida¹, Akio Suzuki², Yuki Shibasaki³, Daisuke Wakabayashi⁴, Nobumasa Funamori⁴

1.東京大学大学院理学系研究科地球惑星科学専攻、2.東北大学大学院理学研究科地学専攻、3.東北大学 学際科学フロンティア研究所、4.高エネルギー加速器研究機構 物質構造科学研究所

1.Department of Earth and Planetary Science, The University of Tokyo, 2.Department of Earth and Planetary Materials Science, Faculty of Science, Tohoku University, 3.Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, 4.Institute of Materials Structure Science, High Energy Accelerator Research Organization

Knowledge of the physical properties of liquid iron alloys is important for understanding the liquid core of the Earth and other terrestrial planets and satellites. Sound velocity is a key physical property to know the structure and composition of these cores because it can be directly compared with seismic observations. However, sound velocity measurements of liquid iron alloy by ultrasonic methods combined with multi anvil apparatus have been limited to below 8 GPa (Nishida et al. 2013; Jing et al. 2014; Kuwabara et al. 2016). Therefore, we have been developing and improving techniques that enable us to measure sound velocities of liquid iron alloys up to 20 GPa. Here we report newly established techniques for sound velocity measurements of liquid Fe-S up to 15 GPa. High-pressure and high-temperature experiments were conducted at the AR-NE7A beamline at the KEK PF-AR synchrotron facility. High pressure was generated by Kawai-type multi anvil apparatus (MAX-III). High temperature was generated using cylindrical resistive heater made of $\text{Al}_2\text{O}_3 + \text{TiC}$ composite. The sample was enclosed in a flat-bottomed cylinder container made of BN with a buffer-rod and a backing plate made of sapphire single crystal. We determined the pressure and temperature simultaneously without a thermocouple from the unit-cell volumes of NaCl and MgO by employing their equations of state. Sound velocity was measured by ultrasonic pulse-echo overlap method. The sample melting was identified during the experiments using X-ray diffraction, and was confirmed afterwards from textural observations of the run products.

Preliminary results show the error in sound velocity of liquid Fe-S at 15 GPa is approximately 2.5% under good conditions. In our previous experiments, the error in sound velocity below 7 GPa was approximately 1% under the best conditions. Taking it into account, our newly techniques can provide satisfactory accuracy. Details and latest experimental results will be presented.

キーワード：核、音速、Fe-S融体、高圧

Keywords: core, sound velocity, liquid Fe-S, high pressure

到来方向検出器開発の現状と地球科学への応用

Development for Neutrino Directional Measurement in Liquid Scintillator

*白旗 豊¹*YUTAKA SHIRAHATA¹

1.東北大学, ニュートリノ科学研究センター

1.RCNS, Tohoku University

Liquid scintillator detector has a sensitivity to geo-neutrino. However, this detector can't observe direction of anti-neutrino. So we develop new-type detector that can know direction of anti-neutrino. In my poster, I talk about development of ^6Li loaded liquid scintillator and high vertex resolution imaging detector.

キーワード：ニュートリノ

Keywords: neutrino

ダイヤモンドアンビルセルを用いた高圧その場X線ラミノグラフィー撮像法の開発

High-pressure *in situ* X-ray laminography using diamond anvil cell*野村 龍一¹、上杉 健太郎²*Ryuichi Nomura¹, Kentaro Uesugi²

1.東京工業大学、2.高輝度光科学研究センター

1.Tokyo Institute of Technology, 2.JASRI/SPring-8

The diamond anvil cell (DAC) is a powerful tool to reproduce high-pressure (P) and high-temperature (T) conditions, corresponding to those of the deep Earth interior, in a laboratory. Various types of measurements such as *in situ* high-P-T spectroscopic measurements and *ex situ* chemical analysis have been conducted using DACs to understand the structure and evolution of the Earth's interior. Among these techniques, 3D visualization and textural/chemical characterization of the internal structure of samples at high-P-T is of great importance. Recently, the dihedral angle of molten iron between bridgmanite was investigated by imaging recovered DAC samples using X-ray computed tomography (CT), and the results provide important insights about the physical process of the Earth's core-mantle separation (Shi et al., 2013). *In situ* high-P X-ray CT has been developed by transmitting X-rays through a light metal gasket, such as Be, between diamond anvils. To date, the applications have been limited to physical purposes, such as the changes of the volume or shape of the sample with pressure (Liu et al., 2008; Wang et al., 2012).

On the other hand, Tsuchiyama et al. (2013) developed a 3D chemical imaging technique, known as analytical dual energy microtomography, in which two X-ray energies below and above the absorption energy of a key element, such as Fe, are used for CT. We applied this technique to recovered DAC samples to determine the solidus temperature of pyrolitic mantle, using incompatible Fe enrichment as a signature of melting (Nomura et al., 2014). This dual energy technique has an additional advantage that artifacts in the reconstructed images, which are a typical problem in CT, can be avoided. The next step should be to carry out *in situ* high-P-T dual energy X-ray imaging, which remains challenging because deep Earth is composed of light elements (e.g. Fe, O, Si, Mg) with an X-ray absorption edge far below the hard X-ray energy range, in which the light metal gasket absorbs the incident X-rays crucially.

A high-pressure *in situ* X-ray laminography technique was developed using a newly designed, laterally open diamond anvil cell. A low X-ray beam of 8 keV energy was used, aiming at future application to dual energy X-ray chemical imaging techniques. The effects of the inclination angle and the imaging angle range were evaluated at ambient pressure using the apparatus. Sectional images of ruby ball samples were successfully reconstructed at high pressures, up to approximately 50 GPa. The high-pressure *in situ* X-ray laminography technique is expected to provide new insights into the deep Earth sciences.

キーワード：高圧、X線イメージング、ダイヤモンドアンビルセル

Keywords: High pressure, X-ray imaging, diamond anvil cell

ボロンドープダイヤモンドヒーターを用いた抵抗加熱式ダイヤモンドアンビルセルの開発 Development of resistance-heated diamond anvil cell using boron-doped diamond heater

*中島 悠貴¹、野村 龍一¹、Xie Longjian²、廣瀬 敬³、米田 明²

*Yuki Nakashima¹, Ryuichi Nomura¹, Longjian Xie², Kei Hirose³, Akira Yoneda²

1.東京工業大学、2.岡山大学地球物質科学研究センター、3.東京工業大学大学院理工学研究科地球惑星科学専攻

1.Tokyo Institute of Technology, 2.Institute for Study of the Earth's Interior, Okayama University, 3.Department of Earth and Planetary Sciences, Tokyo Institute of Technology

Laser-heated diamond anvil cell is the most successful method for reproducing the pressures and temperatures of the Earth's deep interior entirely in a laboratory (e.g. Tateno et al., 2010 Science). However, it is well known that laser-heating results in steep temperature gradients within a sample (e.g. Rainey et al., 2013). Such steep temperature gradients cause the solid-state chemical segregation called as 'Soret diffusion'. This effect is well known in gas chemistry such that heavier elements or elements with larger ionic radii migrate from hot to cold regions, while the lighter elements move in the opposite direction (Grew and Ibbs, 1952). The Soret diffusion takes place not only in gas but also in liquid and solid. Therefore, Soret diffusion makes it difficult to perform chemical equilibrium experiments using laser-heated diamond anvil cell (Sinmyo et al., 2008 JGR).

The technique for homogeneous heating has been developed, that is internally-heated diamond anvil cell. In this method, the heater is put into a sample chamber and compressed together with the sample. Because the diamond has a high thermal conductivity, diamond anvils do not become a high temperature and the heated zone is limited to near the sample chamber. Therefore, internally-heated diamond anvil cell has a potential to generate temperature more than 2000 K because the diamond anvils do not transform to the graphite. A sample was used also as a heater in previous studies (e.g. Liu and Bassett, 1975; Boehler et al., 1986). Therefore, only electric conductor can be used as a sample for the experiment. Zha and Bassett (2003) overcame this situation by making a small hole in a Re heater and filled the hole with sample. Both metal and nonmetal can be used as the sample in this method. The study showed that temperature gradients of Re heater near sample room were gentler than that of laser-heating studies. However, its temperature gradients were still steeper than multi-anvil's temperature gradients (Canil, 1994).

On the other hand, the boron-doped semiconductor diamond heater is known to be able to have a much smaller temperature gradient than that of metallic heater (Yoneda et al., 2014). In multi-anvil experiments, Yoneda et al. (2014) successfully generate high temperature more than 3000 K, which Re and LaCrO₃ heaters cannot generate.

In this study, I developed the internally-heated diamond anvil cell technique using boron-doped diamond heater. The results showed that temperature gradients of boron-doped diamond heater in diamond anvil cell were smaller than that of metallic heater used in previous study (Zha and Bassett, 2003). In addition, its temperature gradients are as steep as multi-anvil's temperature gradients. In this presentation, we will show our recent progress on this study.

キーワード：ダイヤモンドアンビルセル、ボロンドープダイヤモンド、内部抵抗加熱

Keywords: Diamond anvil cell, Boron-doped diamond, Internal resistive heating

酸化水酸化鉄による地球内部への水輸送

Transportation of hydrogen by iron oxide-hydroxide in the Earth's interior

*鈴木 昭夫¹*Akio Suzuki¹

1.東北大学大学院理学研究科地学専攻

1.Department of Earth and Planetary Materials Science, Faculty of Science, Tohoku University

Water (hydrogen) plays important roles in dynamics in the Earth's interior. It is expected that hydrogen is transported from the surface to the interior by the subducting slab. Hydrogen is also a candidate of the light element of outer core. Therefore, the study of the hydrogen in the Earth's interior is quite important. Terasaki et al. (2012) investigated the reaction between Fe-Ni alloy and delta-AlOOH up to the pressure of core-mantle boundary. They showed that hydrogen is partitioned into Fe-Ni alloy and suggested that the core was hydrogenated by the subducting delta-AlOOH. (Dobson and Brodholt (2005) proposed that the banded iron formation subducted to the core-mantle boundary and stagnated there. The banded iron formation contains iron oxide-hydroxide. Therefore, it is important to study the stability and properties of iron oxide-hydroxide under high pressures and high temperatures to discuss the transportation of hydrogen in the Earth's interior. We carried out X-ray diffraction study and X-ray absorption measurement in the Photon Factory, Tsukuba, Japan. Goethite (alpha-FeOOH) is stable at ambient condition. However, it transformed to epsilon-FeOOH at 7.8+-0.5 GPa and 873K. Our previous study showed that epsilon-FeOOH was stable under the condition of the lower mantle. We suggest that epsilon-FeOOH can transport hydrogen to the core-mantle boundary and supply hydrogen to the outer core.

キーワード：含水相、スラブ、マントル、縞状鉄鉱層、外核

Keywords: hydrous phase, slab, mantle, banded iron formation, outer core

マントル組成の東西半球構造と地球ニュートリノの関係性

Possible link between East-west mantle geochemical hemispheres and Geoneutrino

*上木 賢太¹、中村 仁美¹、岩森 光¹*Kenta Ueki¹, Hitomi Nakamura¹, Hikaru Iwamori¹

1.独立行政法人海洋研究開発機構地球内部物質循環研究分野

1.Japan Agency for Marine-Earth Science and Technology, Department of Solid Earth Geochemistry

There has been an extensive debate concerning the compositional structure of the mantle: e.g., two-layered mantle model, stratified/zoned mantle model, plum-pudding mantle model, and marble-cake mantle model (e.g., Tackley, 2008; Iwamori, 2016, for review). The compositional structure is important as it reflects the flow pattern of mantle convection and differentiation processes during material cycling, yet poorly constrained at present. An geochemical end-member model argues that the mantle convection chaotically stirs geochemical heterogeneity brought by plate subduction, resulting in ubiquitously heterogeneous mantle (e.g., plum-pudding or marble-cake mantle, Zindler et al., 1984; Allègre and Turcotte, 1986). At the same time, a large-scale heterogeneity has been argued to exist, e.g., Dupal-anomaly in the southern hemisphere (Dupré and Allègre, 1983; Hart, 1984), or East-west geochemical hemispheres in terms of hydrophilic components (Iwamori and Nakamura, 2012; 2015).

Distributions and abundances of the radiogenic isotopes such as uranium (U) -238, -235 and thorium (Th) -232 in the mantle are key to constrain cooling history and mode of mantle convection of the Earth (e.g., McKenzie et al., 1974; Davies, 1999; Korenaga, 2013), because of the radiogenic heating. However, detailed information of distributions and abundances of such radiogenic elements in the Earth's interior are not well constrained. Recently, Kamioka Liquid-Scintillator Antineutrino Detector measured the geoneutrino flux from decay of U-238 and Th-232. The observations indicate that the heat from radioactive isotopes might account for about a half of Earth's total heat flux (The KamLAND Collaboration, 2011). In addition, Tanaka and Watanabe (2014) proposed Li-loaded directionally sensitive detector for possible geo-neutrino-graphic imaging. Based on these on-going measurements and new methods, the fundamental questions concerning the mantle compositional structures described above could be addressed.

East-west geochemical hemispheres of the mantle have been proposed based on statistical analysis (Independent Component Analysis) of the global isotopic data set of young basalts (Iwamori and Nakamura, 2012; 2015), suggesting that the eastern hemisphere is enriched in "anciently subducted fluid component". If this is the case, we expect a systematic hemispherical difference in elemental abundances and ratios, including U and Th. Although the absolute abundances of these elements in the mantle are not readily constrained by the basalt data (unless the degrees of partial melting of basalts are tightly determined), by combining the geochemical analyses and geoneutrino measurements, we might be able to constrain the distribution and abundances of crucial elements, which will lead us to evaluate Geoneutrino.

キーワード：マントル組成、地球ニュートリノ

Keywords: Mantle composition, Geoneutrino

高温高圧下におけるphase Eggの安定領域

～Phase Eggはマントル遷移層における水の貯蔵庫になり得るか～

Stability field of phase Egg under high temperature and high pressure:

Possibility of phase Egg as a water reservoir in mantle transition zone

*福山 鴻¹、大谷 栄治²、柴崎 裕樹³、小澤 信²、鈴木 昭夫²*Ko Fukuyama¹, Eiji Ohtani², Yuki Shibazaki³, Shin Ozawa², Akio Suzuki²

1.東京大学大学院理学系研究科附属地殻化学実験施設、2.東北大学大学院理学研究科地学専攻、3.東北大学学際科学フロンティア研究所

1.Geochemical Research Center Graduate School of Science, The University of Tokyo, 2.Department of Earth Science, Graduate School of Science, Tohoku University, 3.Frontier Research Institute for Interdisciplinary Sciences, Tohoku University

Water in the earth's interior is one of the main research topics because the water is known to affect physical properties of materials in the earth's interior such as rheology, electric conductivity, seismic velocity, density, and melting point. The sedimentary layer of the oceanic crust transfers water into the deep earth mantle via subducting slabs (Peacock, 1990). Phase Egg, $\text{AlSiO}_3(\text{OH})$, is one of the important hydrous phases in the mantle originating from the sedimentary layer and can contain H_2O of 7.5 wt% as hydroxyl. However, two previous studies (Sano et al., 2004; Pamato et al., 2014) have reported different stability fields about phase Egg. This inconsistency leads to different earth-scientific outlooks on water cycling system via subducting slabs: whether the phase Egg can reserve water in the top of the lower mantle or not and where the superdeep diamond containing phase Egg originates (Wirth et al., 2007). Phase Egg as an inclusion in diamond might indicate a possibility that a top of the lower mantle might be wet.

Here, we conducted high-pressure experiments using Kawai-type 3000 ton multi-anvil apparatus and 1000 ton multi-anvil apparatus at Tohoku University in order to determine the stability field of phase Egg. Experiments were performed in the pressure range of approximately 17-21GPa and in the temperature range of 1000-1200°C. Starting material was a mixture of Al_2O_3 , $\text{Al}(\text{OH})_3$, and SiO_2 compounded similarly to ideal phase Egg composition, which was different from those of two previous studies (Sano et al., 2004; Pamato et al., 2014).

In this study, we found that phase Egg decomposed under the pressure corresponding to the mantle transition zone at 1000°C. This indicates that phase Egg is unstable in the top of lower mantle and can be a water reservoir in the mantle transition zone. In addition, this implies that the superdeep diamond, which Wirth et al. (2007) reported, does not originate from the lower mantle but from the wet mantle transition zone.

キーワード：地球内部の水循環、phase Egg、マルチアンビル型高圧装置、高温高圧、マントル遷移層、superdeep diamond

Keywords: water cycle in the earth's interior, phase Egg, multi-anvil apparatus, high pressure and high temperature, mantle transition zone, superdeep diamond

地球深部における玄武岩マグマの粘度

Viscosity of basaltic magma at high pressure

*鈴木 昭夫¹*Akio Suzuki¹

1. 東北大学大学院理学研究科地学専攻

1. Department of Earth and Planetary Materials Science, Faculty of Science, Tohoku University

Viscosity and density control the mobility of magma. Sakamaki et al. (2013) measured the viscosity of basaltic magma at high pressure and reported the viscosity minimum around 4 GPa. They also measured the density of magma and found a quick elevation of the density around the pressure. On the basis of the results they proposed a model of stagnation of magma around the lithosphere-asthenosphere boundary (melt pond model). However, the pressure range of their measurement was limited to 7 GPa. Reid et al. (2003) carried out viscosity measurement of diopside melt to 13 GPa and reported the viscosity maximum around 10 GPa. They suggested that the decrease in viscosity above 10 GPa was caused by the structural change of melt. Therefore, the change in viscosity is also expected in basaltic magma. Here, we report the result of viscosity measurement of basaltic magma above 10 GPa. We adopted the falling sphere method using the X-ray radiography. Experiments were carried out at the beamline BL-04B1 in SPring-8. We found the decrease in viscosity between 7 and 10 GPa. Above 10 GPa the viscosity increased to 13 GPa. On the basis of the observation of seismic wave, the existence of melt around the base of the upper mantle (e.g., Revenaugh and Sipkin, 1994). We suggest the stagnation of melt by the change in viscosity.

キーワード：マグマ、粘度、マントル

Keywords: magma, viscosity, mantle

Thermal conductivity of lower mantle minerals from *ab initio* anharmonic lattice dynamics
Thermal conductivity of lower mantle minerals from *ab initio* anharmonic lattice dynamics

*出倉 春彦¹、土屋 卓久¹

*Haruhiko DEKURA¹, Taku Tsuchiya¹

1.愛媛大学地球深部ダイナミクス研究センター

1.Geodynamics Research Center, Ehime University

Determination of lattice thermal conductivity (κ) of lower mantle minerals is a key to understanding the dynamics of the Earth's interior. Although determination of κ was impractical in the deep Earth P , T condition for a long time, recent experimental and computational developments have been extending the accessible P and T ranges (e.g. H. Dekura, T. Tsuchiya and J. Tsuchiya, Phys. Rev. Lett. 110, 025904, 2013). *Ab initio* prediction of κ requires understanding of the phonon-phonon interaction associated with the lattice anharmonicity. We recently succeeded in developing an efficient method to calculate it based on the density-functional perturbation theory combined with anharmonic lattice dynamics theory, and applying to MgSiO_3 perovskite in the whole lower mantle P , T range for the first time. Next we extend our techniques to other lower mantle minerals such as MgSiO_3 post-perovskite, and now calculations of more realistic Fe-bearing systems are also started. In this presentation, we introduce the current situation of our research on κ .

キーワード：地球下部マントル、フォノン輸送特性、第一原理計算

Keywords: Earth's lower mantle, Phonon transport property, First-principles calculation

Ab initio prediction of the incongruent melting relation in the MgO-SiO₂ system at multi-megabar

Ab initio prediction of the incongruent melting relation in the MgO-SiO₂ system at multi-megabar

*谷内 聖¹、土屋 卓久¹

*Takashi Taniuchi¹, Taku Tsuchiya¹

1.愛媛大学地球深部ダイナミクス研究センター

1.Geodynamics Research Center, Ehime University

Magnesium silicates are thought to be the major components of the mantle of terrestrial planets and the core of giant planets (Guillot, 1999; Seager *et al.*, 2007). However, the thermodynamic phase equilibrium in the MgO-SiO₂ system is still not well studied at multi-megabar, including melting relations. A recent laser shock experience reported two discontinuous phase changes in MgSiO₃ at 300-400 GPa (Spaulding *et al.*, 2012), but an *ab initio* molecular dynamics study identified no clear transition in MgSiO₃ liquid (Militzer, 2013). Boates and Bonev (2013), on the other hand, examined a decomposition reaction of liquid MgSiO₃ into solid MgO and liquid SiO₂ and reported that liquid MgSiO₃ is dissociated at ~300 GPa. This result implies a possible incongruent melting. However, the reaction they considered is too simple and unrealistic. The detailed phase diagram in the MgO-SiO₂ system is therefore required to be clarified at multi-megabar. In this study, we perform *ab initio* free energy calculations based on the thermodynamic integration method (Kirkwood, 1935) and determine the melting phase relation in this binary system.

キーワード：第一原理計算、MgO-SiO₂ 系、非調和融解、マルチメガバール

Keywords: *ab initio* calculation, MgO-SiO₂ system, incongruent melting, multi-megabar

地球電磁場加速度の時空間分布とマントル最下部電気伝導度

Geoelectromagnetic jerks produced by heterogeneous electrical conductivity in the D" layer

*清水 久芳¹、歌田 久司¹*Hisayoshi Shimizu¹, Hisashi Utada¹

1. 東京大学地震研究所

1. Earthquake Research Institute, University of Tokyo

Geomagnetic jerks are identified at 2003, 2007 and 2011 in the Atlantic and Indian Ocean. It has been suggested by Chulliat et al. (2015) that the regional nature of the geomagnetic jerks and spatio-temporal variations of geomagnetic secular acceleration can be explained by fast equatorial magnetohydrodynamic waves propagating near the surface of the Earth's core. On the other hand, the geoelectric field observed using long baseline submarine cables in the northwestern Pacific also showed sudden change of its secular variation at 2006. In a previous study, we showed using a simplified mantle conductivity models that the geoelectric field variation and the geomagnetic jerk in 2007 can have the same origin; the variations can be generated by a sudden change of the toroidal magnetic field secular variation in the core and influence of high electrical conductivity region in the D" layer beneath the area where the geomagnetic field variation was evident. Since the geoelectric and geomagnetic field variations can have the same origin, we call the variations a geoelectromagnetic jerk. In this presentation, we show results of numerical experiment on the electromagnetic field in the mantle due to the toroidal magnetic field variation in the core employing more realistic distribution of the electrical conductivity in the D" layer based on the SH-wave velocity model obtained by Takeuchi (2012). Conductivity heterogeneity at the Earth's surface due to the conductivity contrast between the land and seawater on the electric field is also included in the numerical experiment to examine its effect on the voltages observed by submarine cables. Typical spatio-temporal variations of the observed and simulated geoelectromagnetic field are examined to consider plausibility of geoelectromagnetic jerk hypothesis.

Chulliat, A., Alken, P. and Maus, S. (2015) Fast equatorial waves propagating at the top of the Earth's core, *Geophys. Res. Lett.*, 42, 3321-3329.

Takeuchi, N. (2012) Detection of ridge-like structures in the Pacific Large Low-Shear-Velocity Province, *Earth and Planet. Sci. Lett.*, 319-320, 55-64.

Experimental and theoretical thermal equations of state of MgSiO_3 post-perovskite at multi-megabar pressures

Experimental and theoretical thermal equations of state of MgSiO_3 post-perovskite at multi-megabar pressures

*境 毅¹、出倉 春彦¹

*Takeshi Sakai¹, Haruhiko Dekura¹

1.愛媛大学地球深部ダイナミクス研究センター

1.Geodynamics Research Center, Ehime University

The MgSiO_3 post-perovskite phase is the most abundant silicate phase in a super-Earth's mantle, although it only exists within the Earth's lowermost mantle. We established the thermal equations of state (EoS) of the MgSiO_3 post-perovskite phase, which were determined by using both laser-heated diamond anvil cell (LHDAC) and density-functional theoretical techniques, within a multi-megabar pressure range, corresponding to the conditions of a super-Earth's mantle. The LHDAC experiments were performed at up to a pressure of 265 GPa at a temperature of 300 K, and 170 GPa at 2560 K. The *ab initio* calculations were performed at up to 1.2 TPa and 5000 K. The Keane and AP2 EoS models, which include parameters that limit to infinity at high pressure, were adopted for the first time to extract meaningful physical properties. The experimental volume data in a wide pressure-temperature range enabled us to determine the fully experimentally based parameters for the Mie-Grüneisen-Debye model. The Grüneisen parameter and its volume dependency were found to be consistent with their theoretically obtained values. Both the experimental and theoretical EoS are also found to be in very good agreement with one another, within 0.1% in volume at the earth's core-mantle boundary condition, and the relation is maintained within 0.8% even up to a pressure and temperature of 300 GPa and 5000 K, respectively. Our newly developed EoS should be applicable to a super-Earth's mantle, as well as the Earth's core-mantle boundary region.

キーワード：ポストペロブスカイト、状態方程式、スーパーアース

Keywords: post-perovskite, equation of state, super-Earth

Sensitivity of Core Phases on F-layer

Sensitivity of Core Phases on F-layer

*大滝 壽樹¹、金嶋 聡²*Toshiki Ohtaki¹, Satoshi Kaneshima²

1.産業技術総合研究所地質情報研究部門、2.九州大学大学院理学研究院地球惑星科学部門

1.Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology (AIST), 2.Department of Earth and Planetary Sciences, Kyusyu University

Last year we showed that the dispersion in PKPbc and differential traveltimes between PKiKP and PKPbc are particularly sensitive to the F-layer structure (lowermost outer core) and are insensitive to the structure of the other parts of the Earth (Ohtaki and Kaneshima, 2015). In previous studies, the Vp structure of the F-layer have been investigated using absolute traveltimes of PKPbc/c-diff, differential traveltimes between PKPbc/c-diff and PKiKP, amplitude ratios between PKPbc/c-diff and PKiKP, and the position of the C-cusp. PKiKP pierces the inner-core boundary and turns in the inner core; PKPbc turns in the lower part of the outer core; PKPc-diff diffracts on the inner-core boundary beyond the C-cusp; PKiKP reflects on the inner core boundary. In this study we discuss the sensitivity of various core phases (PKiKP, PKPbc, PKPc-diff, and PKiKP) to the F-layer structure in detail. Among these observations, absolute traveltimes of PKPbc are affected by crustal and mantle structures that are strongly heterogeneous and are not precisely known, which indicates benefits of analyzing differential travels times. Differential traveltimes suppress the effects of heterogeneous structures as well as the discrepancy between a reference seismic model and the real Earth above the turning depths of rays. However, it is difficult to discriminate the P-wave velocity of the F-layer from that of the inner core using the differential traveltimes between PKPbc/c-diff and PKiKP, because the inner core is more heterogeneous than the F-layer. Fine structure of the F-layer is also poorly constrained by the amplitude ratios because of the low sensitivity of the ratios to the Vp gradient and of a trade-off between the Vp profile of the F-layer and the attenuation values in the inner core. The C-cusp position can be constrained only poorly by the amplitude observations, and there exist many velocity profiles that yield the same C-cusp position. In summary, conventional observations are obviously insufficient to resolve detailed F-layer structure.

In our previous study (Ohtaki et al., 2012), we examined the seismic structure near the inner core boundary beneath the South Pole. In that study, we investigated the velocity above the inner core boundary using the amplitude ratios between PKiKP and PKPbc/c-diff, assuming a constant velocity in the F-layer, because the ratio is not so sensitive to the velocity gradient, as mentioned above. In this study, we also examine a tolerance level of a velocity gradient there.

キーワード : Seismic velocity, Outer Core

Keywords: Seismic velocity, Outer Core

Anomalously large PKiKP/PcP amplitude ratios at frequency of around 1 Hz observed by USArray

Anomalously large PKiKP/PcP amplitude ratios at frequency of around 1 Hz observed by USArray

*田中 聡¹

*Satoru Tanaka¹

1. 海洋研究開発機構 地球深部ダイナミクス研究分野

1. Department of Deep Earth Structure and Dynamics Research Japan Agency for Marine-Earth Science and Technology

Tanaka and Tkalcic (2015) observed the frequency dependent PKiKP/PcP amplitude ratios by Hi-net, of which ray paths are passing below the Western Pacific. They showed the existence of a spectral peak at approximately 2 to 3 Hz, a spectral hole at 1 and 3 Hz, and no peak and hole, suggesting lateral variations at the inner core surface. Here I report frequency dependent PKiKP/PcP amplitude ratios observed by USArray, of which reflection points are located below the Central America. I find also a large scatter of the frequency characteristics. Interestingly, the spectral peaks in the PKiKP/PcP spectral ratios around 1-1.5 Hz are detected when the reflection points are located below the east of Mexico that is not observed by Hi-net. The peak amplitude is about 2 times greater than that around 2 Hz. Based on the finite difference simulations by Tanaka and Tkalcic (2015), this observed peak can be explained by ICB sinusoidal topography with wavelength and height of 0.5 km, or ICB spiky topography with wavelength and height of 1.5 km. This observation suggests that the crystallization at the inner core surface or inner core growth system below the east of Mexico is different from that below the Western Pacific.

キーワード：内核、USArray、PKiKP/PcP

Keywords: Inner core, USArray, PKiKP/PcP

On the possible scenario of thermal evolution of Earth's core with high thermal conductivity in a coupled core-mantle evolution model

On the possible scenario of thermal evolution of Earth's core with high thermal conductivity in a coupled core-mantle evolution model

中川 貴司¹、*松井 宏晃²

Takashi Nakagawa¹, *Hiroaki Matsui²

1.海洋研究開発機構数理科学・先端技術研究分野、2.カリフォルニア大学デービス校

1.MAT, JAMSTEC, 2.UC Davis

The thermal conductivity measurement of iron alloy from high P-T physics suggested to range from 60 to 150 W/m/K under temperature and pressure condition in Earth's core [e.g. Gomi et al., 2013]. The previous study by Nakagawa and Tackley [2015] indicated that the CMB heat flow was just only 6 TW and, as a result, the magnetic evolution would be failed with high thermal conductivity and colder CMB temperature (~3500 K) caused by large adiabatic temperature gradient across the Earth's core (~1 K/km) [Labrosse, 2015]. Here we assume smaller adiabatic temperature gradient across the Earth's core (0.5 to 0.7 K/km taken from lower-bound value in Ichikawa et al. [2014]) as well as high thermal conductivity of Earth's core set as 120 W/m/K. For the successful scenario of a coupled core-mantle thermal evolution matching the current size of the inner core and continuous magnetic field generation, the CMB heat flow at the present time-scale would be around 12 TW because the CMB temperature is still high (~4000 K) for finding the current size of the inner core and the age of the inner core would be 1.2 billion year, which seems to be a bit older age of inner core compared to other studies [Labrosse, 2015; Davis, 2015]. To find the successful scenario of thermal evolution of Earth's core, the adiabatic temperature gradient across the Earth's core prescribed by Grueneisen parameter and bulk modulus would be quite important in terms of high CMB heat flow than could find the continuous magnetic evolution under the high thermal conductivity. With the latest update of core-mantle evolution, the adiabatic temperature across Earth's core would be consistent with the range found from first principle calculation rather than that used in the core evolution models approximated heat transfer across the CMB.

キーワード：地球コア、コアーマントル熱流量、断熱温度勾配、熱伝導率

Keywords: Earth's core, Heat flow across the core-mantle boundary, Adiabatic temperature gradient, Thermal conductivity