

## 物性物理学の観点から説明される地球の内部構造

## The Earth's internal structure explained from a viewpoint of the condensed matter physics

\*唐澤 信司<sup>1</sup>\*Shinji Karasawa<sup>1</sup>

1.宮城工業高等専門学校 名誉教授

1.Miyagi National College of Technology Professor emeritus

地球の深さの増加に応じて物質に及ぼす圧力が増加します。外部から圧力を受けると物質のサイズは縮小し、凝集力のポテンシャルエネルギーは大きくなります。物質は低いエネルギーの状態になる時に低下する分のエネルギーを放出します。ところで、隕石が地表に衝突する際には衝突により圧縮された物質は収縮して発熱しますが、その熱は直後の膨張で吸収されてしまいます。隕石が衝突するとクレーターを作りますが、衝突の周辺の地域の地面が溶融していた形跡はありません。ちなみに、地球の中心部の定常温度は最高で7000℃であり、0.63 eV の電子状態に相当します。地球の中心部の温度をそのような状態にするのは衝突のエネルギーではなくて、集積した物質の性質によるとして説明できます。

地球の地下深くなるほど電子軌道間の相互作用が大きくなり、より低いエネルギーの状態が可能になります。熱運動により隣接原子を交換して置かれた状況に適した物質になることが可能であれば物質はその変化をします。物質はエネルギーが低い状態になる時に、前後の差のエネルギーによって発熱します。その熱はマントルを經由して地殻に放出されます。熱の移動に伴って物質も移動するのでマントル対流が起こります。その対流は海のプレートの下のマントルの方が大陸の地殻の下より放熱が盛んであるので、海底に湧き出します。地球の深い内部の物質の性質は次のようになります。地球内部の高い圧力で物質の電子軌道がオーバーラップすると、同じエネルギーレベルに複数の電子軌道が存在する縮重状態になります。縮重した電子状態は電気伝導性と流動性を持ち、その物質は液体金属のような性質を示します。それが地球内部の外核の状態です。ところが物質の縮重状態の電子のエネルギーレベルは原子の配置の対称性を低くすると分裂して、さらに低いエネルギーの電子状態が得られるというヤーン・テラー効果があります。そのヤーン・テラー効果によって横波を伝える固体の状態になったのが地球の中心部の内核の状態です。

このように集まった物質の電子状態を考慮して次の事柄を説明しました。(1)マントル対流の熱源、(2)海のプレートが沈み込む付近の火山の熱源、(3)地球の外殻の液体の状態、(4)地球の内核の固体の状態、(5)太陽系の形成、及び(6)地球型惑星の形成。詳しくは以下のWebsite をご覧ください。

([https://www.youtube.com/watch?v=Wi5G2F\\_pDXM&feature=youtu.be](https://www.youtube.com/watch?v=Wi5G2F_pDXM&feature=youtu.be) )

キーワード：マントル対流、外核、内核、断熱圧縮、縮重電子状態、ヤーン・テラー効果

Keywords: Mantle convection, Outer core, Inner core, Adiabatic compression, Degenerated state of electron, Jahn-Teller-Effect

低粘性層を有する月の深部構造：月の潮汐応答に関する最新の測地学的係数に基づいて修正された制約条件

The deep lunar interior with a low-viscosity zone: Revised constraints from recent geodetic parameters on the tidal response of the Moon

\*原田 雄司<sup>1,2</sup>、ホーセンス サンダー<sup>3</sup>、松本 晃治<sup>4</sup>、エン 建国<sup>5</sup>、平 勁松<sup>6</sup>、野田 寛大<sup>4</sup>、春山 純一<sup>7</sup>

\*Yuji Harada<sup>1,2</sup>, Sander Goossens<sup>3</sup>, Koji Matsumoto<sup>4</sup>, Jianguo Yan<sup>5</sup>, Jinsong Ping<sup>6</sup>, Hiroto Noda<sup>4</sup>, Junichi Haruyama<sup>7</sup>

1. 澳門科技大学太空科学研究所、2. 東京大学地震研究所、3. メリーランド大学ボルチモア郡校宇宙科学技術研究探査センター、4. 自然科学研究機構国立天文台、5. 武漢大学測絵遥感信息工程国家重点実験室、6. 中国科学院国家天文台、7. 宇宙航空研究開発機構宇宙科学研究所

1.SSI, MUST, 2.ERI, UT, 3.CRESST, UMBC, 4.NAOJ, NINS, 5.LIESMARS, WHU, 6.NAOC, CAS, 7.ISAS, JAXA

We revisit the constraints on the deep lunar interior with a possible low-viscosity zone at the core-mantle boundary obtained from our previous forward modeling of the tidal response of the Moon, by comparing a numerical model with several tidal parameters that have been improved or are newly determined by recent geodetic observations and analyses from GRAIL (gravity), LRO (shape), and LLR (rotation). Our results are in principle consistent with the latest data and lead to a thicker low-viscosity layer (with an outer radius of about 540 to 560 km, which is much larger than that of about 500 km in our earlier investigation) which reaches just below or inside the place where many seismic nests of deep moonquakes are located.

キーワード：粘性、月、内部、潮汐、ラブ数、クオリティ係数

Keywords: viscosity, Moon, interior, tide, Love number, quality factor

## Seafloor subsidence and mantle dynamics

## Seafloor subsidence and mantle dynamics

\*Adam Claudia<sup>1</sup>、King S<sup>1</sup>、Rabinowicz M<sup>2</sup>、Vidal Valerie<sup>3</sup>、Jalobeanu A<sup>4</sup>、吉田 晶樹<sup>5</sup>

\*Claudia Adam<sup>1</sup>, S King<sup>1</sup>, M Rabinowicz<sup>2</sup>, Valerie Vidal<sup>3</sup>, A Jalobeanu<sup>4</sup>, Masaki Yoshida<sup>5</sup>

1.Virginia Polytechnic Institute and State University、2.GET, Observatoire Midi-Pyrénées, Toulouse、3.Laboratoire de Physique, ENS Lyon、4.Bayes Map Solutions, LLC, Castro Valley, CA、5.独立行政法人海洋研究開発機構 地球深部ダイナミクス研究分野

1.Virginia Polytechnic Institute and State University, 2.GET, Observatoire Midi-Pyrénées, Toulouse, 3.Laboratoire de Physique, ENS Lyon, 4.Bayes Map Solutions, LLC, Castro Valley, CA, 5.Department of Deep Earth Structure and Dynamics Research, Japan Agency for Marine-Earth Science and Technology (JAMSTEC)

The subsidence of seafloor is generally considered as a passive phenomenon related to the conductive cooling of the lithosphere since its creation at mid-oceanic ridges. Recent alternative theories suggest that the mantle dynamics plays an important role in the structure and depth of the oceanic lithosphere. However, the link between mantle dynamics and seafloor subsidence has still to be quantitatively assessed. Here we provide a statistic study of the subsidence parameters (subsidence rate and ridge depth) for all the oceans. These parameters are retrieved through the positive outliers method, a classical method used in signal processing. We also model the mantle convection pattern from the S40RTS tomography model. The density anomalies derived from this model are used to compute the instantaneous flow in a global 3D spherical geometry, and the induced dynamic topography.

The variations of the mid-oceanic ridge depths are well recovered by the modeled dynamic topography. The systematic fit of the bathymetry allows the recovery of the subsidence rate, from which we derive the effective thermal conductivity,  $k_{\text{eff}}$ . This parameter ranges between 1 and 7  $\text{Wm}^{-1}\text{K}^{-1}$ . We show that departures from the  $k_{\text{eff}}=3 \text{ Wm}^{-1}\text{K}^{-1}$  standard value are systematically related to mantle convection and not to the lithospheric structure. Regions characterized by  $k_{\text{eff}}>3 \text{ Wm}^{-1}\text{K}^{-1}$  are associated with the uplift of mantle plumes. Regions characterized by  $k_{\text{eff}}<3 \text{ Wm}^{-1}\text{K}^{-1}$  are related to large scale mantle downwellings such as the Australia-Antarctic Discordance (ADD) or the return flow from the South Pacific Superswell to the East Pacific rise. This demonstrates that the mantle dynamics plays a major role in the shaping of the oceanic seafloor. In particular, the parameters generally considered to quantify the lithosphere structure, such as the thermal conductivity, are not only representative of this structure but also incorporate signals from the mantle convection occurring beneath the lithosphere.

## Effect of along-trench slab dimension on subduction-induced upper mantle flow: Insights from 3-D laboratory models

\*Vincent Strak<sup>1,2</sup>, Wouter Pieter Schellart<sup>1</sup>

1.Monash University, 2.Earthquake Research Institute

Three-dimensional self-evolving subduction models with various along-trench dimensions have been quantitatively analysed in the laboratory by means of a stereoscopic Particle Image Velocimetry (sPIV) technique. The purpose was (1) to provide information on the pattern of the upper mantle flow induced by subduction, particularly focusing on the location and magnitude of upwellings around the lateral slab edges and in the mantle wedge, (2) to study the evolution of mantle upwellings in terms of location and magnitude, and (3) to study the effect of along-trench slab dimension on upwelling location and magnitude. The model results show that 4 types of upwelling are generated by subduction in a Newtonian upper mantle. One of these upwellings occurs laterally away from the sub-slab domain and could potentially trigger decompression melting, thereby explaining the occurrence of a certain type of intraplate volcanism. Two other significant upwellings are observed in the mantle wedge. The tested along-trench slab dimensions were comparable to slabs of narrow (e.g., Calabria) to wider (e.g., Tonga-Kermadec-Hikurangi) subduction zones. The results indicate that both the location and magnitude of the upwelling occurring laterally away from the sub-slab domain are affected by the along-trench slab dimension, with wider slabs producing faster and more focused upwellings that are located closer to the lateral slab edges. The models also seem to show that the along-trench slab dimension controls the intensity of upwellings in the mantle wedge.

Keywords: Subduction, Mantle flow, Upwellings

## Seismic structure and dynamics beneath eastern Tibet: Insight into large earthquakes and Tengchong volcano

\*Lei Jianshe<sup>1</sup>, Dapeng Zhao<sup>2</sup>

1.Key Laboratory of Crustal Dynamics, Institute of Crustal Dynamics, China Earthquake Administration, 2.Department of Geophysics, Tohoku University

We review recent studies of seismic tomography and earthquake sequences, and discuss their implications for seismotectonics and mantle dynamics beneath eastern Tibet. The crustal structures in the source areas of the 2008 *M*<sub>s</sub> 8.0 Wenchuan and the 2013 *M*<sub>s</sub> 7.0 Lushan earthquakes are similar and exhibit prominent low-velocity (low-*V*) and high-Poisson's ratio anomalies in the source areas, indicating that in addition to compositional variations, fluid-filled rock matrices exist in the Longmenshan fault zone, which may have influenced the nucleation of the two earthquakes. Significant low-*V* anomalies are revealed between the Wenchuan and Lushan mainshocks, which may explain why their aftershock zones extend northward and southward, respectively. The relocated aftershocks of the 2011 *M*<sub>s</sub> 5.8 Yingjiang and the 2014 *M*<sub>s</sub> 6.5 Ludian earthquakes show a conjugate-shaped distribution, which may explain why the two moderate-sized earthquakes caused heavy damage. The large earthquakes in eastern Tibet are located at boundaries of low- or high-velocity anomalies in the upper mantle. The structural heterogeneities in the crust and upper mantle are associated with hot and wet upwelling and corner flows in the big mantle wedge above the subducting Indian slab beneath eastern Tibet, as well as slab dehydration, which affect the seismogenesis in the region.

Keywords: Seismic structure, Mantle dynamics, Eastern Tibet, Large earthquake, Tengchong volcano

## Transformational Faulting as a Deep-Focus Earthquake Mechanism: Correlating In-Situ Acoustic Emission Locations at High P-T with Post-Mortem Fault Imaging Using Synchrotron X-Ray Microtomography in Controlled Deformation Experiments

\*Yanbin Wang<sup>1</sup>, Lupei Zhu<sup>2</sup>, Feng Shi<sup>1</sup>, Tony Yu<sup>1</sup>, Mark L Rivers<sup>1</sup>, Alexandre Schubnel<sup>3</sup>, Nadege Hilairet<sup>4</sup>, Fabrice Brunet<sup>5</sup>

1.Center for Advanced Radiation Sources, University of Chicago, Chicago, IL, USA, 2.Department of Earth Sciences, St. Louis University, St. Louis, MO, USA, 3.Laboratoire de Géologie, CNRS UMR 8538, Ecole Normale Supérieure, Paris, France, 4.Université des Sciences et Technologies de Lille, Lille, France, 5.CNRS-ISTerre, Université de Grenoble, Grenoble, France

One of the possible mechanisms responsible for deep-focus earthquakes occurring at depths below 350 km is faulting induced or triggered by phase transformation from metastable olivine to its high-pressure phases (wadsleyite and ringwoodite, both are related to the general spinel structure) in the mantle transition zone. We have studied transformational faulting from olivine to spinel in  $Mg_2GeO_4$ , a close analog of the silicate olivine  $(Mg,Fe)_2SiO_4$ , using a high-pressure deformation apparatus in conjunction with in-situ acoustic emission (AE) monitoring [1]. Synchrotron X-ray microtomography (XMT) [2] was used to image the samples recovered from high-pressure deformation experiments, with spatial resolution of  $\sim 0.005$  mm. In this study, we establish spatial correlations between AE events observed during in-situ deformation experiments and faults imaged by XMT post-mortem. The nature of high-pressure experiments limits the sample size to the order of 2 mm in linear dimensions, and the acoustic system we used has a maximum sampling rate of 50 MHz (i.e., 20 ns between adjacent sampling points), which limits spatial resolution of AE locations to about 0.3 mm. This makes it difficult to locate AE events accurately. A well-developed double-difference cross-correlation (CC) algorithm (hypoDD [3]) developed for seismological studies has been successfully adapted in the analysis of AE locations, improving AE allocation spatial resolution by a factor of  $\sim 10$ . This algorithm also helps separate events with various waveforms, which are related to different fault planes and faulting directions. The CC algorithm classifies AE events into various groups. Events within each group share common characteristics and may be considered occurring within the same faulting mechanism. These groups of events display excellent correlations with faults imaged by XMT, with two major groups of AE events correlating well with two conjugated faults in XMT images. Furthermore, time sequence of AE events can be examined to investigate details of formation of macroscopic faults from the micro AE events. These results help understand the dynamic process of transformational faulting.

### References:

- [1] A. Schubnel *et al.*, Deep-Focus Earthquake Analogs Recorded at High Pressure and Temperature in the Laboratory. *Science* 341, 1377 (2013).
- [2] M. L. Rivers, Y. Wang. (2006), Recent developments in microtomography at GeoSoilEnviroCARS, in: *Developments in X-Ray Tomography V*, U. Bonse (ed.), *Proc. SPIE* vol. 6318, pp. 63180J-63180J-15
- [3] F. Waldhauser, W. L. Ellsworth, A Double-Difference Earthquake Location Algorithm: Method and Application to the Northern Hayward Fault, California. *Bulletin of the Seismological Society of America* 90, 1353 (2000).

Keywords: Deep-focus earthquakes, High P-T deformation, Acoustic emission, Synchrotron x-ray diffraction and imaging, Earthquake mechanisms

## Behavior of basalt-carbonate melts at high pressures

## Behavior of basalt-carbonate melts at high pressures

\*Kono Yoshio<sup>1</sup>, Ohfuji Hiroaki<sup>2</sup>, Stagno Vincenzo<sup>2</sup>, Kenney-Benson Curtis<sup>1</sup>, Wang Yanbin<sup>3</sup>, Shen Guoyin<sup>1</sup>

\*Yoshio Kono<sup>1</sup>, Hiroaki Ohfuji<sup>2</sup>, Vincenzo Stagno<sup>2</sup>, Curtis Kenney-Benson<sup>1</sup>, Yanbin Wang<sup>3</sup>, Guoyin Shen<sup>1</sup>

1.Carnegie Institution of Washington, 2.Ehime University, 3.The University of Chicago

1.Carnegie Institution of Washington, 2.Ehime University, 3.The University of Chicago

Knowledge of the occurrence and mobility of carbonate-rich melts in the Earth's mantle is important for understanding the deep carbon cycle and related geochemical and geophysical processes.

Recently, Kono et al. (2014) find that viscosities of carbonate melts are surprisingly low, in the range of 0.006-0.010 Pa s, which are ~2 to 3 orders of magnitude lower than those of basaltic melts in the upper mantle. As a result, the mobility of carbonate melts (defined as the ratio of melt-solid density contrast to melt viscosity) is ~2 to 3 orders of magnitude higher than that of basaltic melts. Such high mobility of carbonate melts may have significant influence on several magmatic processes, such as fast melt migration and effective melt extraction beneath mid-ocean ridges. However, the behaviour of carbonate-rich silicate melt may be complex, as demonstrated in the reported immiscible behaviour of silicate-carbonate melts. To predict potential implications of carbonate-rich silicate melt as they migrate up in the upper mantle, the stability should be investigated at those pressures. Here we investigate behavior of basalt-dolomite melts with 20-61 vol.% dolomite compositions at high pressures to 7.3GPa by using newly developed X-ray phase contrast imaging technique in Paris-Edinburgh cell at beamline 16BMB, HPCAT, in the Advanced Photon Source. We find that basalt plus 32-61 vol.% dolomite becomes single melt at pressure >~5.5 GPa, while it separates into silicate-dominant melt plus CO<sub>2</sub> at lower pressures. The abrupt change of stability of carbonate-rich silicate melt may play a role of paramount importance in migration velocity of upwelling carbonate-silicate magmas due to separation of carbonate component from silicate melt, with 2-3 orders of magnitude difference of viscosity between basalt and carbonate melts.

## Reference

Kono, Y., C. Kenney-Benson, D. Hummer, H. Ohfuji, C. Park, G. Shen, Y. Wang, A. Kavner, and C. E. Manning (2014), Ultralow viscosity of carbonate melts at high pressures, *Nature Communications*, 5, 5091.

キーワード : carbonate、melt、high pressure

Keywords: carbonate, melt, high pressure

## 沈み込むスラブの構造と運動学的観測から見た滞留スラブの形成と進化

Stagnant slab formation and evolution inferred from kinematic and geometrical observations of the subduction zone

\*中久喜 伴益<sup>1</sup>

\*Tomoeeki Nakakuki<sup>1</sup>

1. 広島大学大学院理学研究科地球惑星システム学専攻

1. Department of Earth and Planetary Systems Science, Graduate School of Science, Hiroshima University

Seismic tomography revealed variable slab structure in the mantle transition zone (e.g., Fukao et al., 2009). Numerical modeling that incorporates phase transition at the 660 km depth and trench migration succeeded to simulate formation and avalanche of the stagnant slab (Christensen, 1996; Torii and Yoshioka, 2007; Nakakuki et al., 2010). Mechanisms to control structure of the transition zone slab are still controversial (e.g., Karato et al., 2001; King et al., 2015), since we have not sufficiently understood relationship of the slab dynamics with tectonic features in the actual subduction zone (cf. Billen, 2010). In this study, subduction zone geometric and kinematic observations are analyzed aiming to construct a scenario for the stagnant slab formation and evolution. The data, which are compiled by Lallemand et al. (2005) and (2008), include absolute motions of the subducting plate, the overriding plate and the trench based on a global plate motion model (SB04) constructed by Steinberger et al. (2004), slab depths, and slab dip angles.

Important observations connected to the stagnant slab dynamics are summarized as follows. (1) Most of upper mantle slabs are retreating with the velocity larger than 1 [cm/yr]. (2) In the subduction zone with the slab penetrating at the depth between 660 to 1200 km, trench motion is advance (e.g., Mariana, Java) or retreat with the slower velocity ( $< 1$  [cm/yr], e.g., north Kuril) than that in the subduction zone with the upper slab. (3) Back-arc extension often occurs in the subduction zone in which the maximum slab depth is 660 km (e.g., Tonga). (4) Dip angles of the upper and lower mantle slabs become steeper with the age. (5) Stagnant slabs are classified into 2 types. (i) The first type has young age ( $< 60$  [Ma]), slow subducting plate motion (the slab descending motion occur as if the plate peeled and fell off from the surface (e.g., New Hebrides)). (ii) The second type has old age ( $> 100$  [Ma]) and dip angles obviously shallower ( $< \sim 50$  [deg], e.g., south Kuril, Japan) than those of the slab penetrating into the lower mantle.

These results indicate important relationship of the slab dip angle and trench migration to the slab structure varying with the sinking depth. The following scenario emerges as a possible explanation for the stagnant slab mechanics. (1) A relatively young slab with shallow dip angle and trench retreat collides to the 660 km phase transition. (2) The slab start stagnates at 660 km, and the slab rollback is further enhanced. (3) When the dip angle of the stagnating slab is shallow, the slab retains the stagnation at the 660 km after the trench retreat is declined. (4) When the dip angle is steepened and/or the trench advance is generated owing to increase of subducting plate age or collision of the continent, the slab penetrates into the lower mantle.

キーワード：滞留スラブ、マントル遷移層、プレート運動、沈み込み帯

Keywords: stagnant slab, mantle transition zone, plate kinematics, subduction zone



## クロミタイトの微細構造に保存されたマントル遷移層における変形

## Messengers from the deep: Fossil wadsleyite-chromite microstructures from the Mantle Transition Zone

\*佐津川 貴子<sup>1</sup>、Griffin William<sup>2</sup>、Piazolo Sandra<sup>2</sup>、O'Reilly Suzanne<sup>2</sup>

\*Takako Satsukawa<sup>1</sup>, William L Griffin<sup>2</sup>, Sandra Piazolo<sup>2</sup>, Suzanne Y O'Reilly<sup>2</sup>

1.京都大学大学院理学研究科地球惑星科学専攻地球物理学教室、2.CCFS&GEMOC, Department of Earth and Planetary Sciences, Macquarie University

1.Department of Geophysics, Division of Earth and Planetary Sciences, Kyoto University,

2.CCFS&GEMOC, Department of Earth and Planetary Sciences, Macquarie University

Investigations of the Mantle Transition Zone (MTZ; 410-660 km deep) by deformation experiments and geophysical methods suggest that the MTZ has distinct rheological properties, but their exact cause is still unclear due to the lack of natural samples. Here we present the first direct evidence for crystal-plastic deformation by dislocation creep in the MTZ using a chromitite from the Luobusa peridotite (E. Tibet). Chromite grains show exsolution of diopside and SiO<sub>2</sub>, suggesting previous equilibration in the MTZ. Electron backscattered diffraction (EBSD) analysis reveals that olivine grains co-existing with exsolved phases inside chromite grains and occurring on chromite grain boundaries have a single pronounced crystallographic preferred orientation (CPO). This suggests that olivine preserves the CPO of a high-pressure polymorph (wadsleyite) before the high-pressure polymorph of chromite began to invert and exsolve. Chromite also shows a significant CPO. Thus, the fine-grained high-pressure phases were deformed by dislocation creep in the MTZ. Grain growth in inverted chromite produced an equilibrated microstructure during exhumation to the surface, masking at first sight its MTZ deformation history. These unique observations provide a window into the deep Earth, and constraints for interpreting geophysical signals and their geodynamic implications in a geologically robust context.

キーワード：マントル遷移層、クロミタイト、かんらん石、結晶方位定向配列、EBSD、微細構造

Keywords: Mantle transition zone, chromitite, olivine, CPO, EBSD, microstructure

Mantle transition zone beneath a normal seafloor in the northwestern Pacific: Electrical conductivity, seismic discontinuity, and water content

Mantle transition zone beneath a normal seafloor in the northwestern Pacific: Electrical conductivity, seismic discontinuity, and water content

\*松野 哲男<sup>1</sup>、末次 大輔<sup>2</sup>、歌田 久司<sup>1</sup>、馬場 聖至<sup>1</sup>、多田 訓子<sup>2</sup>、清水 久芳<sup>1</sup>、塩原 肇<sup>1</sup>、一瀬 建日<sup>1</sup>、杉岡 裕子<sup>3</sup>、伊藤 亜妃<sup>2</sup>

\*Tetsuo Matsuno<sup>1</sup>, Daisuke Suetsugu<sup>2</sup>, Hisashi Utada<sup>1</sup>, Kiyoshi Baba<sup>1</sup>, Noriko Tada<sup>2</sup>, Hisayoshi Shimizu<sup>1</sup>, Hajime Shiobara<sup>1</sup>, Takehi Isse<sup>1</sup>, Hiroko Sugioka<sup>3</sup>, Aki Ito<sup>2</sup>

1.東京大学地震研究所、2.海洋研究開発機構、3.神戸大学

1.Earthquake Research Institute, The University of Tokyo, 2.Japan Agency for Marine-Earth Science and Technology, 3.Kobe University

We conducted a joint electromagnetic (EM) and seismic experiment to probe the mantle structure below a normal seafloor around the Shatsky Rise in the northwestern Pacific. Specifically for the investigation of the mantle transition zone (MTZ) structure, we deployed our state-of-the-art instruments in two arrays to the north and south of the Shatsky Rise for 5 years from 2010 to 2015. Here we report the result of analyses of EM and seismic data obtained in the joint experiment as well as in previous studies. From the EM data analysis, a 1-D profile of electrical conductivity was obtained for two observational arrays. The thickness of the MTZ was also obtained by the P-wave receiver function analysis, from which temperature profiles in the MTZ below two arrays were then estimated. We found that the northern array provides EM and seismic data with much higher quality than the southern array, and therefore meaningful estimation of MTZ water content is possible for the northern array. We estimated water content based on profiles of electrical conductivity and temperature obtained by our geophysical observation and electrical conductivities of dry and wet MTZ minerals (wadsleyite and ringwoodite) obtained by mineral physics. The result of the forward modeling study indicated that the upper limit of water content below the northern array is 0.5 wt.% or 0.05 wt.%, depending on different results of laboratory experiments for water effects on electrical conductivities. The lower limit of water content was not constrained by our data.

## H-D inter-diffusion in Fe-free wadsleyite: implication for multiple hydrogen mechanism

\*Wei Sun<sup>1</sup>, Takashi Yoshino<sup>1</sup>, Naoya Sakamoto<sup>2</sup>, Hisayoshi Yurimoto<sup>3</sup>

1.Research Group of Physics of the Earth's Interior at the Institute for Study of the Earth's Interior, Okayama University, 2.Isotope Imaging Laboratory, Creative Research Institution, Hokkaido University, 3.Department of Natural History Sciences, Hokkaido University

It has been proposed that concentration and speciation of water-derived species in minerals may adjust in response to changes of point defect condition [Demouchy and Mackwell, 2006]. To lift the veil of Fe effect on hydrogen diffusion and isotopic differentiation, a sets of H-D inter-diffusion experiments were conducted in Fe-free wadsleyite single crystal couples at various temperatures, 16 GPa and compared with our former study on Fe-bearing wadsleyite. Distinguish with symmetric profiles in Fe-bearing condition, H-D inter-diffusion in Fe-free wadsleyite revealed evidently asymmetric properties and it indicates deuterium diffuses about 1 order faster than hydrogen in Fe-free wadsleyite. Both magnitude and anisotropy of H-D inter-diffusion in Fe-free condition are largely different with Fe-bearing condition, which strongly demonstrated a multiple hydrogen mechanism proposed by Karato (2013) association with free proton migration in interstitial sites. Simulation model suggests free proton migration in interstitial sites dominates the hydrogen diffusion in Fe-free condition and asymmetric properties might owe to the distinguished jumping probabilities from Mg sites to interstitial sites between hydrogen and deuterium.

Keywords: inter-diffusion, wadsleyite, asymmetric, multiple

High-pressure phase relations in the system  $\text{MgCr}_2\text{O}_4$ - $\text{Mg}_2\text{SiO}_4$  with implications to ultra-high pressure chromitites in ophiolites

High-pressure phase relations in the system  $\text{MgCr}_2\text{O}_4$ - $\text{Mg}_2\text{SiO}_4$  with implications to ultra-high pressure chromitites in ophiolites

\*赤荻 正樹<sup>1</sup>、姉川 由輝<sup>1</sup>、河原 愛理<sup>1</sup>、石井 貴之<sup>1</sup>、糀谷 浩<sup>1</sup>

\*Masaki Akaogi<sup>1</sup>, Yuki Anegawa<sup>1</sup>, Airi Kawahara<sup>1</sup>, Takayuki Ishii<sup>1</sup>, Hiroshi Kojitani<sup>1</sup>

1. 学習院大学理学部化学科

1. Department of Chemistry, Faculty of Science, Gakushuin University

Podiform chromitites which are enveloped by dunite are commonly found in harzburgite in ophiolites. It is interpreted that the podiform chromitites were formed through reaction between harzburgite and melt at a shallow upper mantle level. However, some podiform chromitites recently found in Luobusa ophiolites, Tibet, were interpreted to be derived from the deep mantle, because the chromitites contained diamond and coesite as inclusions. Yamamoto et al. (2009) and Arai (2013) suggested that the chromitites originally consisted of calcium-ferrite (CF) structured phase in the deep mantle and that they transformed to spinel-structured chromitites during upwelling.

We have recently determined phase relations in  $\text{MgCr}_2\text{O}_4$  at pressure up to 28 GPa and temperature to 1600 °C. Magnesium chromite ( $\text{MgCr}_2\text{O}_4$ ) with the spinel structure first transforms to a two-phase assemblage of modified ludwigite (mLd)-type  $\text{Mg}_2\text{Cr}_2\text{O}_5$  + corundum-type  $\text{Cr}_2\text{O}_3$  at 13-15 GPa, and the two phases combine into a calcium-titanate (CT) phase at 16-20 GPa (Ishii et al., 2015). Because chromite minerals in natural podiform chromitites are enveloped in olivine crystals, we have extended our experimental study to the system  $\text{MgCr}_2\text{O}_4$ - $\text{Mg}_2\text{SiO}_4$  to examine phase relations and mutual solubilities in the high-pressure phases.

High-pressure high-temperature experiments were performed in composition of 50mol% $\text{MgCr}_2\text{O}_4$ .50mol% $\text{Mg}_2\text{SiO}_4$  up to 26 GPa at 1600 °C using a Kawai-type multianvil apparatus. Phase identification of the synthesized samples were made using micro-focus and powder X-ray diffractometers. Composition analysis was performed using a scanning electron microscope with an energy-dispersive X-ray spectrometer. At pressure below about 14 GPa,  $\text{Mg}_2\text{SiO}_4$  forsterite coexists with  $\text{MgCr}_2\text{O}_4$ -rich spinel. At 14-19 GPa, a three-phase assemblage, mLd + garnet (Gt) + wadsleyite (Wd) becomes stable. This assemblage changes into  $\text{MgCr}_2\text{O}_4$ -rich CT + ringwoodite at about 20 GPa, which further changes into an assemblage of  $\text{MgSiO}_3$ -rich bridgmanite (perovskite) + CT + periclase at about 23 GPa. The mLd composition was close to  $\text{Mg}_2\text{Cr}_2\text{O}_5$ , while Gt has a middle composition between  $\text{Mg}_4\text{Si}_4\text{O}_{12}$  and  $\text{Mg}_3\text{Cr}_2\text{Si}_3\text{O}_{12}$ . Solubility of  $\text{Mg}_2\text{SiO}_4$  component in CT increases with pressure to about 20 mol% at 26 GPa. These results implies that, when podiform chromitites are subducted into the deep mantle,  $\text{MgCr}_2\text{O}_4$  spinel and forsterite react to form mLd + Gt at 14-19 GPa which corresponds to pressure range of the upper part of the mantle transition zone. Also, if CT phase containing substantial  $\text{Mg}_2\text{SiO}_4$  component goes upward from the deep mantle, it decomposes into the assemblage containing mLd + Gt at 14-19 GPa. To the best of our knowledge, mLd + Gt in association with chromite have not yet been found in the natural chromitites. Therefore, we suggest that possible occurrence of mLd and/or Gt coexisting with chromite is an important indicator which shows that the chromitites were derived from the depth at least upper-part of the transition zone.

キーワード : high pressure、chromitite、phase relation

Keywords: high pressure, chromitite, phase relation

## Boron doped diamond heater in the Kawai-type apparatus

\*Longjian Xie<sup>1</sup>, Akira Yoneda<sup>1</sup>, Eiji Ito<sup>1</sup>

1. Institute for Study of the Earth Interior, Okayama University

Boron doped diamond (BDD) heater has attracted special attention because of its high melting point and X-ray transparency. In the previous studies, graphite boron composite (GBC) was usually used as a precursor of BDD. In the present study, we firstly investigated the application of pre-synthesized BDD cylinder and BDD powder as cylindrical heaters in the Kawai-type apparatus at 15 GPa. BDD with 0.5 and 3.0 wt % was synthesized at 15 GPa and 2100 °C. The BDD powders were grinded from BDD blocks by using Nano Polycrystalline Diamond mortar at GRC, Ehime Univ.; SEM image showed that the grain size of diamond is about 1 μm. Both kinds of heaters showed good stability and high reproducibility. The BDD heater with 0.5 wt % boron (resistivity: ~0.001 Ω.m at 1500 °C) showed semi-conductive behavior, i.e. decreasing resistance with increasing temperature. However, the BDD heater with 3.0 wt % boron showed lower resistivity (about 0.00016 Ω.m at 1500 °C) and metallic behavior, i.e. increasing resistance with increasing temperature. This electrical characteristic enables us to adjust the boron concentration of BDD to get the desired resistivity. We succeeded to generate temperature about 3500 °C by using small heater (1.5 mm outer diameter, 1 mm inner diameter, 6 mm length) with TiC electrode. Temperature higher than 1800 °C was estimated by input power according to its temperature-power relationship. The pressure generation efficiency by assembly using BDD heater was checked by *in-situ* X ray experiments at SPring-8. Compared with assembly using GBC heater, the use of BDD heater has lower efficiency of pressure generation in the beginning stage of compression, while has higher efficiency in the later stage. It is obvious that the BDD heater is more advantageous than GBC heater, because it is free from the complicated temperature-power relationship and pressure drop associated with graphite to diamond conversion.

Keywords: Diamond synthesise, Boron doped diamond heater, Ultrahigh temperature, Multi-anvil apparatus

CO<sub>2</sub> レーザー加熱ダイヤモンドアンビルセルを用いた下部マントル条件下における MgO-MgSiO<sub>3</sub> 系の  
溶融関係

Melting relations in the MgO-MgSiO<sub>3</sub> system under the lower mantle condition using a CO<sub>2</sub>  
laser heated diamond anvil cell

\*大西 里佳<sup>1</sup>、桑山 靖弘<sup>2</sup>、井上 徹<sup>2</sup>

\*Satoka Ohnishi<sup>1</sup>, Yasuhiro Kuwayama<sup>2</sup>, Toru Inoue<sup>2</sup>

1.愛媛大学大学院理工学研究科、2.愛媛大学地球深部ダイナミクス研究センター

1.Graduate School of Science and Engineering, Ehime University, 2.Geodynamics Research Center,  
Ehime University

Seismological observations of the ultralow-velocity zones (ULVZs) suggest the presence of partial melts above the core-mantle boundary (CMB). Knowledge of the melting relations in the lower mantle is a key to understand the chemical differentiation at the base of the mantle. While melting relations of mantle materials at relatively low pressure (below 30 GPa) have been extensively studied using a multi-anvil apparatus (e.g. Ito *et al.*, 2004 *Phy. Earth Planet. Inter.*), the melting experiments at higher pressures are still limited. Only in a few model rock compositions, such as peridotite and mid-oceanic ridge basalt (MORB), the experiments were conducted under the CMB conditions using a laser-heated diamond anvil cell (LHDAC) (e.g. Fiquet *et al.*, 2010 *Science*, Andrault *et al.*, 2014 *Science*). Since chemical heterogeneity of both major elements (Mg, Si, Fe, Al...) and minor ones (e.g. alkalis and volatiles) should have a large effect on the melting behavior, the melting phase diagrams as a function of composition are fundamental to understand the nature of the ULVZs. For melting relations in a binary system MgO-MgSiO<sub>3</sub>, which is a major component in the lower mantle, the experiment up to only 26 GPa was performed (Liebske and Frost, 2012 *Earth Planet. Sci. Lett.*). So, further studies at higher pressure corresponding to the deep lower mantle condition are required. In this study, we determined the melting relations in the MgO-MgSiO<sub>3</sub> system above 30 GPa using a LHDAC. Glasses of several different compositions in the MgO-MgSiO<sub>3</sub> system (from 37 to 45 mol% SiO<sub>2</sub>) were used as the starting materials. A CO<sub>2</sub> laser heating system was used to heat the sample directly. The recovered samples were polished and analyzed by a dualbeam focused ion beam (FIB) and a field emission scanning electron microscope (FE-SEM), respectively. The eutectic compositions and the liquidus phases were determined on the basis of chemical and textural analysis of the quenched samples. Our results show that the eutectic composition at 30 GPa is about 44 mol% SiO<sub>2</sub> and it becomes about 40 mol% at 50 GPa. Above 50 GPa, it is predicted to become relatively constant, consistent with the previous result by Liebske and Frost (2012). From these results, MgO-rich melt layer may be generated by partial melting of the bulk mantle, such as pyrolite composition (i.e. 42 mol% SiO<sub>2</sub>), at the base of the mantle. The present result should provide basic information for better understanding on the melting relations at deep mantle conditions.

## X線非弾性散乱による下部マントル物質のフォノン物性

Phonon properties of the lower mantle minerals by inelastic x-ray scattering

\*福井 宏之<sup>1</sup>、米田 明<sup>2</sup>、芳野 極<sup>2</sup>、バロン アルフレッド<sup>3</sup>\*Hiroshi Fukui<sup>1</sup>, Akira Yoneda<sup>2</sup>, Takashi Yoshino<sup>2</sup>, Alfred Q.R. Baron<sup>3</sup>

1.兵庫県立大学大学院物質理学研究科、2.岡山大学地球物質科学研究センター、3.理化学研究所放射光科学総合研究センター

1.Graduate School of Material Science, University of Hyogo, 2.Institute for Study of the Earth's Interior, Okayama Univeristy, 3.RIKEN SPring-8 Center

Inelastic x-ray scattering is a useful technique to obtain phonon properties of lower mantle minerals with single crystal samples. We would like to discuss what kind of information can be derived from IXS spectra and how to do it with treating bridgmanite and ferropericalse as examples.

キーワード：下部マントル、フォノン、X線非弾性散乱

Keywords: lower mantle , phonon, inelastic x-ray scattering

## 大陸の離合集散を含む3次元マントル対流と物質循環

Three-dimensional mantle convection and material cycling with continental dispersal and coalescence

\*柳 竜之介<sup>1</sup>、吉田 晶樹<sup>2</sup>

\*Ryunosuke Yanagi<sup>1</sup>, Masaki Yoshida<sup>2</sup>

1.東京工業大学、2.JAMSTEC

1.Tokyo Institute of Technology, 2.JAMSTEC

The Earth is unique among the solar terrestrial planets, having the continents and the abundant liquid water on its surface. In this study, we have developed a three-dimensional mantle convection model that incorporates these two features into the numerical program, and investigate how the flow-temperature field and water distribution occur under the circumstances of continental dispersal and coalescence. In this numerical simulation of 3-D spherical mantle convection, supercontinent is introduced in the model set up in order to study how continental dispersal and coalescence happens and affect the structures of the interior. In addition, water is also introduced, considering water solubility of mantle rocks. The results show that water is transported to broader regions effectively once continental dispersal occurred, due to enhanced flow velocity associated with increased horizontal temperature gradient partly with the blanketing effect beneath the supercontinent. This process distributes the water in a spatially wide region as a "migrating entrance" of water into the mantle. Once the water subducts with the cold down-going flow and reaches CMB (core mantle boundary), then it horizontally spreads to be heated up. In some case, dehydration of Phase H may occur to generate a fluid that migrates upward and hydrate the overlying mantle, which occurs near the boundary between the cold continental domain and the warmer oceanic domain, yet within the hydrated cold domain beneath the (super)continent region. This mechanism, together with the near-surface process described above, may create a hydrous domain beneath the continental region where a rather uniform hydration may prevail. Therefore, the effective redistribution with a global scale structure can be created simultaneously. Such a mechanism could be important to account for the observed geochemical mantle hemispheres.

キーワード：大陸の離合集散、物質循環

Keywords: continental dispersal and coalescence, material cycling



2次元軸対称モデルによるスーパー地球マントル内上昇プルームの数値シミュレーション  
Numerical simulations on the upwelling plumes in the mantle of super-Earths in 2-D  
axisymmetric geometry

\*亀山 真典<sup>1</sup>

\*Masanori Kameyama<sup>1</sup>

1. 国立大学法人愛媛大学地球深部ダイナミクス研究センター

1. Geodynamics Research Center, Ehime University

We carried out numerical simulations on the upwelling plumes in the mantle of the Earth and that of super-Earths. The objects of our study are (1) to confirm the validity of our program which is newly developed for this study by comparing our numerical results with those of earlier studies on the upwelling plumes in the Earth's mantle and (2) to examine the effects of temperature-dependent viscosity and internal heating rate on upwelling plumes in the mantle of super-Earths. In this study, we consider a major upwelling plume at the center of 2-D axisymmetric model. We have carried out two series of numerical simulations. In a first series, we calculated thermal convection in an incompressible Boussinesq fluid under the conditions identical to those in earlier studies. In a second series, we performed simulations under the conditions for the mantle of super-Earths using the truncated anelastic liquid approximation (TALA).

In the first series of our simulations, we obtained the results similar to earlier ones, such as the increase in the heat flow at the top and bottom boundaries in proportion to  $Ra^{1/3}$ , demonstrating the validity of the numerical simulation in this study. Furthermore, the second series of our study of the mantle of super-Earths showed that (1) temperature-dependence of viscosity tends to reduce plume heat transport, and (2) the variation of compressibility does not affect the detail of the loss of plume heat flux during their ascent. We also found that heat can flow downward into the core when the chondritic rate of internal heating is present in the mantle of super-Earths, demonstrating significant effects of the internal heating rate on the convection and thermal state in the mantle of super-Earths.

キーワード：スーパー地球、マントル対流、内部発熱

Keywords: super-Earths, mantle convection, internal heating

## 第一原理計算による高圧下における固体鉄水素合金の研究

## First-principles study of solid iron-hydrogen alloys under high pressure

\*梅本 幸一郎<sup>1</sup>、廣瀬 敬<sup>1</sup>\*Koichiro Umemoto<sup>1</sup>, Kei Hirose<sup>1</sup>

1.東京工業大学地球生命研究所

1.Earth-Life Science Institute, Tokyo Institute of Technology

Hydrogen and iron are two of major constituents of the Earth and planetary interiors. The crystal structure of solid FeH<sub>x</sub> is one of the most fundamental information in order to understand properties of planetary cores. Recently, hydrogen-rich phases, FeH<sub>2</sub> and FeH<sub>3</sub>, were experimentally synthesized [1]. The crystal structure of FeH<sub>3</sub> was clarified by comparing experimental compression curve with calculated one. On the other hand, the structure of FeH<sub>2</sub> remains unclear. It is mainly because the hydrogen positions are quite difficult to be determined by x-ray diffraction measurements. Ref. 1 proposed the crystal structure of FeH<sub>2</sub>, but it is less consistent with its experimental compression curve. Here we report the results of first-principles calculations on FeH<sub>2</sub>. We find the new hydrogen positions which lead to more stable structure than proposed by Ref. 1 and reproduces experimental compression curve very well. Our new structure will be essential for constraining the amount of hydrogen in iron alloys.

[1] C. M. Pépin, A. Dewaele, G. Geneste, P. Loubeyre, and M. Mezouar, Phys. Rev. Lett. 113, 265504 (2014).

キーワード：鉄水素合金、第一原理計算、高圧における結晶構造

Keywords: Iron-hydrogen alloys, First principles, Crystal structure at high pressure

## マルチアンビル装置を用いた熱電対起電力への圧力効果の測定法

Determination of pressure effect on the thermocouple electromotive force using multi-anvil apparatus

\*西原 遊<sup>1</sup>、福家 一樹<sup>1</sup>、丹下 慶範<sup>2</sup>、肥後 祐司<sup>2</sup>

\*Yu Nishihara<sup>1</sup>, Kazuki Fuke<sup>1</sup>, Yoshinori Tange<sup>2</sup>, Yuji Higo<sup>2</sup>

1.愛媛大学地球深部ダイナミクス研究センター、2.高輝度光科学研究センター

1.Geodynamics Research Center Ehime University, 2.Japan Synchrotron Radiation Research Center

Our understanding of Earth's interior highly depends on physical and chemical properties of Earth materials which were determined based on high-pressure and high-temperature experiments. Temperature in high-pressure and high-temperature experiments is mostly determined using a thermocouple without any pressure correction. This may lead to erroneous results in estimated temperature and thus physical and chemical properties of Earth materials due to significant pressure effects of the thermocouple electromotive force (EMF). Getting and Kennedy (1970) determined the pressure effect on the EMF of two types of thermocouples up to 3.5 GPa and 1000°C. However, no new knowledge has been obtained on the absolute pressure effect on thermocouple EMF for more than 40 years.

In this study, we developed a method to determine the absolute pressure effect on thermocouple electromotive force, based on a single wire method using Kawai-type multi-anvil apparatus. By applying this method, single wire EMFs were measured up to a pressure of 7 GPa and temperature of 600°C for chromel and alumel, which passes through the entire length of the pressure medium. The chromel and alumel wires were contained in semi-sintered MgO and dense Al<sub>2</sub>O<sub>3</sub> insulating tubes, and the portions of the wires in MgO and Al<sub>2</sub>O<sub>3</sub> were subjected to higher and lower pressures, respectively. The temperature along the single wires was calibrated by separate experiments employing multiple thermocouples. Pressure conditions along the wires were evaluated based on in situ X-ray diffraction using synchrotron X-ray radiation and a thermal equation of state of Ni. The pressure effect of the Seebeck coefficients of chromel and alumel, determined by the analyses of single wire EMFs and pressure-temperature profiles along the wires, was virtually consistent with that of previous lower-pressure or lower-temperature studies. The difference between the nominal temperature by chromel-alumel thermocouple (type K) and the real temperature was calculated to be from 0 to -3°C in conditions up to 7 GPa and 600°C. Since the multi-anvil apparatus is capable of achieving much higher pressure and temperature, the method presented in this study promises to reveal absolute temperature correction for thermocouples over a wide range of pressure and temperature conditions.

キーワード：熱電対、高温高圧実験

Keywords: thermocouple, high-pressure and high-temperature experiments

## GHz-P波及びS波バッファロッド・トランスデューサーの開発

Development of P wave and S wave GHz buffer rod-transducer for DAC experiments

\*米田 明<sup>1</sup>\*Akira Yoneda<sup>1</sup>

1.岡山大学地球物質科学研究センター

1.Institute for Study of the Earth's Interior, Okayama University

P wave and S wave GHz buffer rods are now under developing. Both the transducers will be used for ultrasonic velocity measurement under high pressure in the combination of diamond anvil cell. The way of electrode for transducer is found to be critical for stable operation of buffer rod; I examined Au sputtering on the ZnO transducer, and got reasonable results. Simultaneously, I examined Sapphire, YAG, TiO<sub>2</sub>, Si, and SiO<sub>2</sub> crystals for feasibility of GHz buffer rod, and found substantial attenuation of GHz wave in Si and SiO<sub>2</sub> crystals. I confirmed the performance of Sapphire, YAG, and TiO<sub>2</sub> buffer rods with observing the reflection pulse from diamond plate (~2 mm thickness). I also detected a P-S conversion signal in YAG crystal; it is almost ready to assemble S wave GHz buffer rod made of YAG crystal. Furthermore I have a plan to make diamond buffer rod as well in near future.

キーワード：GHz、バッファロッド、弾性

Keywords: GHz, buffer rod, elasticity

## 下部マントル圧力下におけるブリッジマナイトの格子熱伝導率

Lattice thermal conductivity of bridgmanite at the lower mantle pressures

\*奥田 善之<sup>1</sup>、太田 健二<sup>1</sup>、八木 貴志<sup>2</sup>、新名 良介<sup>3</sup>、廣瀬 敬<sup>3</sup>\*Yoshiyuki Okuda<sup>1</sup>, Kenji Ohta<sup>1</sup>, Takashi Yagi<sup>2</sup>, Ryousuke Sinmyo<sup>3</sup>, Kei Hirose<sup>3</sup>

1.東京工業大学 理学院 地球惑星科学系 地球惑星科学コース、2.産業技術総合研究所 物質計測標準研究部門 熱物性標準研究グループ、3.東京工業大学 地球生命研究所

1.Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2.Research Institute for Material and Chemical Measurement, National Institute of Advanced Industrial Science and Technology, 3.Earth-Life Science Institute

The amount of heat flow from the Earth's core to mantle critically determines the thermo-chemical evolution both of the core and the mantle. Bridgmanite, iron and aluminum bearing  $\text{MgSiO}_3$  perovskite, is the most abundant mineral in the Earth's lower mantle, and thus its thermal transport property controls heat transport in the lower mantle. Here we measured lattice thermal diffusivity of bridgmanite with chemical composition of  $\text{Mg}_{0.832}\text{Fe}_{0.209}\text{Al}_{0.060}\text{Si}_{0.916}\text{O}_3$  up to 39 GPa at 300 K using the pulsed light heating thermoreflectance technique in a diamond anvil cell. The results indicate that the lattice thermal conductivity of the bridgmanite sample is slightly lower than that of iron and aluminum free bridgmantie determined by using the same experimental technique (Ohta et al., 2012). Our result exhibit insignificant effect of iron and aluminum incorporation into bridgmanite on its thermal conductivity, which imply temperature variation in the lower mantle is only factor to induce heterogeneity of thermal conductivity and core heat flux there that could drive large scale dynamics both in the core and mantle.

Reference: Ohta, K. et al. Lattice thermal conductivity of  $\text{MgSiO}_3$  perovskite and post-perovskite at the core-mantle boundary. *Earth Planet Sc Lett* 349-350, 109-115 (2012).

キーワード：熱伝導率、下部マントル、ブリッジマナイト

Keywords: thermal conductivity, lower mantle, bridgmanite

The effects of ferromagnetism and interstitial hydrogen on the equation of states of hcp  $\text{FeH}_x$

The effects of ferromagnetism and interstitial hydrogen on the equation of states of hcp  $\text{FeH}_x$

\*五味 斎<sup>1</sup>、Fei Yingwei<sup>2</sup>、芳野 極<sup>1</sup>

\*Hitoshi Gomi<sup>1</sup>, Yingwei Fei<sup>2</sup>, Takashi Yoshino<sup>1</sup>

1.岡山大学地球物質科学研究センター、2.Geophysical Laboratory, Carnegie Institution of Washington  
1.Institute for Study of the Earth's Interior, Okayama University, 2.Geophysical Laboratory, Carnegie Institution of Washington

Hydrogen is one of the important candidates of the light elements of the Earth's core, because only a small amount of interstitial hydrogens may drastically change the physical properties of compressed iron (e.g. melting temperature, density and elastic properties). Since the solubility of hydrogen has strong pressure dependence, hydrogen content must be determined by in-situ observations. Combined with X-ray diffraction measurements, the following relation is widely used for this purpose:  $x = (V_{\text{FeH}_x} - V_{\text{Fe}}) / V_{\text{H}}$ , where  $x$  is the hydrogen content,  $V_{\text{H}}$  is the volume expansion caused by unit concentration of hydrogen,  $V_{\text{FeH}_x}$  and  $V_{\text{Fe}}$  are volumes of  $\text{FeH}_x$  and pure iron, respectively (Fukai, 1992). Ambient pressure experiments on many face-centered cubic (fcc) metals suggest that  $V_{\text{H}}$  is insensitive to the hydrogen content  $x$ . However, it has not been confirmed for compressed iron. We computed the equation of states of hexagonal-closed pack (hcp) iron with interstitial hydrogen with  $x = 0.0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9$  and  $1.0$  by using the AkaiKKR firstprinciples package. Coherent potential approximation (CPA) was adopted to treat non-saturated hydrogen atoms, which randomly occupied the octahedral sites. The results of pure ( $x = 0.0$ ) and hydrogen saturated ( $x = 1.0$ ) iron are consistent with previous experiments (Hirao et al., 2004; Narygina et al., 2011). However, we found a discontinuous volume change as functions of hydrogen content for non-saturated  $\text{FeH}_x$  because of the ferromagnetic transition. We also found almost no  $x$  dependence on the volume in the ferromagnetic phase. Previous Mossbauer spectroscopy measurements suggest the ferromagnetic state is stable up to about 25 GPa for iron hydride (Mitsui et al., 2010; Narygina et al., 2011). This means that previous experiments possibly overestimate the hydrogen content of ferromagnetic  $\text{FeH}_x$ .

キーワード：FeH<sub>x</sub>、強磁性、状態方程式、KKR-CPA、第一原理計算

Keywords: FeH<sub>x</sub>, ferromagnetism, equation of states, KKR-CPA, firstprinciples calculation

## 地球核条件下にレーザー衝撃圧縮されたFeSの音速と密度

Sound velocity and density of liquid iron sulfide up to 600 GPa by laser-shock compression

尾方 優士<sup>1</sup>、\*境家 達弘<sup>1</sup>、近藤 忠<sup>1</sup>、寺崎 英紀<sup>1</sup>、重森 啓介<sup>2</sup>、弘中 陽一郎<sup>2</sup>Yushi Ogata<sup>1</sup>, \*Tatsuhiko Sakaiya<sup>1</sup>, Tadashi Kondo<sup>1</sup>, Hidenori Terasaki<sup>1</sup>, Keisuke Shigemori<sup>2</sup>, Yoichiro Hironaka<sup>2</sup>

1.大阪大学大学院理学研究科宇宙地球科学専攻、2.大阪大学レーザーエネルギー学研究センター

1.Graduate School of Science, Osaka University, 2.Institute of Laser Engineering, Osaka University

Sound velocity at Earth's core conditions are one of the most important physical properties in Earth science because it can be directly compared with the seismological Earth model (PREM: Preliminary Reference Earth Model) [1]. The composition of solid inner core is estimated from the comparison of the model [1] and the extrapolation of sound velocities as a function of density of iron and iron alloys obtained by the static compression experiment [2, 3]. Birch's law, a linear sound velocity-density relation [4], is used to extrapolate sound velocities to densities in the core condition. On the other hand, the composition of liquid outer core is estimated from the partitioning and solubility data in the inner core boundary condition for the composition of solid core. There has been some works for the sound velocity of iron and iron alloys on the Earth's core condition by dynamic techniques using explosive [5], gas gun [5-9], and laser [10-12]. Huang et al. estimated that the outer core composition is Fe with 0.5 wt.% O and 9.5 wt.% S by the comparison of PREM and sound velocities of Fe-S-O system [9]. In this study, we measure the sound velocity and density of liquid iron sulfide by shock-compression method using high-power laser.

We conducted shock-compression experiments using a High Intensity Plasma Experimental Research (HIPER) system at the GEKKO-XII laser irradiation facility [10] at the Institute of Laser Engineering, Osaka University. The sample is pyrrhotite ( $\text{Fe}_{0.9}\text{S}$ ). The sound velocities and densities of shock-compressed pyrrhotite using the high-power laser were measured by x-ray radiography [10-12] at pressures up to 600 GPa.

Part of this work was performed under the joint research project of the Institute of Laser Engineering, Osaka University.

## References

- [1] A. M. Dziewonski and D. L. Anderson, *Phys. Earth Planet. Inter.* 25, 297-356 (1981).
- [2] J. Badro et al., *Earth Planet. Sci. Lett.* 254, 233-238 (2007).
- [3] G. Fiquet et al., *Phys. Earth Planet. Inter.* 172, 125-129 (2009).
- [4] F. Birch, *Geophys. J. R. Astron. Soc.* 4, 295-311 (1961).
- [5] J. M. Brown and R. G. McQueen, *J. Geophys. Res.* 91, 7485-7494 (1986).
- [6] J. H. Nguyen and N. C. Holmes, *Nature* 427, 339-342 (2004).
- [7] J. W. Shaner et al., *Shock Waves in Condensed Matter*, 135-138 (1988).
- [8] H. Huang et al., *J. Geophys. Res.* 116, B04201 (2010).
- [9] H. Huang et al., *Nature* 479, 513-516 (2011).
- [10] K. Shigemori et al., *Eur. Phys. J. D* 44, 301-305 (2007).
- [11] K. Shigemori et al., *Rev. Sci. Instrum.* 83, 10E529 (2012).
- [12] T. Sakaiya et al., *Earth Planet. Sci. Lett.* 392, 80-85 (2014).
- [13] C. Yamanaka et al., *Nucl. Fusion* 27, 19-30 (1987).

キーワード：音速、レーザー、衝撃圧縮、FeS、地球核

Keywords: Sound velocity, Laser, Shock compression, Pyrrhotite, Earth's core



## 2観測点間の表面波位相・振幅データを用いた北米大陸上部マントルの3次元構造

3D shear wave structure in the North American upper mantle from interstation phase and amplitude data of surface waves

\*浜田 広太<sup>1</sup>、吉澤 和範<sup>2</sup>

\*Kouta Hamada<sup>1</sup>, Kazunori Yoshizawa<sup>2</sup>

1.北海道大学大学院理学院、2.北海道大学大学院理学研究院

1.Graduate School of Science, Hokkaido University, 2.Faculty of Science, Hokkaido University

The North American continent encompasses a variety of complex structural features, including regions with east-west extension, active volcanoes in the west, and stable cratons in the central and eastern areas. The high-quality broadband seismic data derived from the high-density Transportable Array (USArray) have facilitated many new tomographic studies in this region. We have recently developed a new technique to measure interstation phase speeds and amplitude ratios for fundamental-mode surface waves based on non-linear waveform fitting between two stations (Hamada and Yoshizawa, 2015, GJI). The amplitude anomalies of surface waves are affected by not only anelastic attenuation, but also elastic focusing/defocusing, which reflect the second derivatives of phase velocity across the ray path. Thus, the amplitude data are sensitive to shorter-wavelength structure than the conventional phase data.

The method has been applied to observed waveforms of USArray from 2007 to 2014, and we could achieve fairly uniform ray-path coverage across the U.S. We collect a large-number of phase speed and amplitude data for short interstation distances less than 1000 km, which can be helpful in enhancing the horizontal resolution of velocity models in North America. The measured interstation phase speed and amplitude ratios are inverted simultaneously for phase speed maps as well as local amplification factor at receiver locations. Our phase speed maps derived from both phase and amplitude data tend to exhibit better recovery of the strength of velocity perturbations than those from phase data only, with enhanced local-scale tectonic features characterized by strong velocity gradients. The results indicate that interstation amplitude data can be of help in reconstructing shorter-wavelength structures of the upper mantle.

Isotropic 3-D shear wave models in the depth range from 50 to 200 km are constructed from the phase speed maps of fundamental-mode Rayleigh waves. The isotropic S wave speed models show significant slow anomalies in Rio Grande Rift and Yellowstone in the western United States at shallower depth above 100 km. Smaller-scale tectonic features are also mapped clearly in the eastern areas; e.g., slow anomalies in New Madrid Seismic Zone as well as Great Meteor Hotspot Track, which can be found down to the depth of about 100 km.

キーワード：表面波、地震波トモグラフィー、北米大陸

Keywords: Surface waves, Seismic tomography, North America

Multi-scale heterogeneities in continental lithosphere and their implications for lithospheric discontinuities

Multi-scale heterogeneities in continental lithosphere and their implications for lithospheric discontinuities

\*吉澤 和範<sup>1</sup>、Kennett Brian L.N.<sup>2</sup>

\*Kazunori Yoshizawa<sup>1</sup>, Brian L.N. Kennett<sup>2</sup>

1.北海道大学大学院理学研究院、2.オーストラリア国立大学地球科学研究所

1.Faculty of Science, Hokkaido University, 2.Research School of Earth Sciences, Australian National University

Lines of evidence from recent seismological observations have revealed the nature of large-scale and fine-scale heterogeneities in the continental lithosphere. Large-scale 3-D images of the continental lithosphere have been constrained primarily from surface waves, which indicate the fast waves speed anomalies of the cratonic lithosphere with the typical lithospheric thickness of about 200 km. Surface wave observations in Australia also provide the evidence of strong radial anisotropy with faster SH wavespeed than SV, particularly at the shallower depth above 90 km as well as in the asthenosphere, while the strength of radial anisotropy tends to be weak in the middle to lower lithosphere (Yoshizawa, 2014, PEPI).

Discontinuities in the lithosphere have been well constrained by the body-wave receiver functions that indicate the wave conversions and reflections underneath seismic stations. One of the striking features in the S-wave receiver function analysis in Australian cratons is the clear signals of discontinuities in the mid-lithosphere at around 70-90 km (Ford et al., 2010, EPSL), which might indicate the rapid velocity drop or change in the character of radial anisotropy at the middle depth of the continental lithosphere where the wave speed is highest in general. This estimated depth of the enigmatic mid-lithosphere discontinuity (MLD) from receiver functions coincides well with the rapid change in the strength of radial anisotropy derived from surface waves (Yoshizawa & Kennett, 2015, GRL).

The latest observations from high-frequency P-wave reflectivity profiles derived from the auto-correlograms of ambient noises in the vertical component (Kennett, 2015, EPSL) provide us with an additional constraint on the nature of the lithospheric heterogeneity. The P reflectivity profiles have suggested vertical changes in the characters of fine-scale structures in the Australian continent, indicating stronger reflectivity in the crust and upper lithosphere underneath the cratons. Such observations support the existence of fine-scale laminated heterogeneity in the lithosphere, as suggested by the numerical simulations for high-frequency scattering of seismic waves for the paths in the cratonic areas (Kennett & Furumura, 2008, GJI; 2016, G-cubed). The existence of such horizontally laminated fine-scale heterogeneity in the upper lithosphere causes equivalent effects as the shape preferred orientation, which may eventually generate apparent change in the radial anisotropy as well as apparent discontinuities in the mid-lithosphere. Linking all these independent observation from a variety of seismic signals with different frequency band will be of help in the further understanding of the nature of lithospheric heterogeneity and anisotropy.

キーワード：リソスフェア、異方性、不均質性、不連続面、表面波

Keywords: Lithosphere, Anisotropy, Heterogeneity, Discontinuity, Surface Waves

Lowermost mantle dynamics driven by the plate subduction  
Lowermost mantle dynamics driven by the plate subduction

\*金子 岳郎<sup>1</sup>、中久喜 伴益<sup>1</sup>

\*Takeo Kaneko<sup>1</sup>, Tomoeki Nakakuki<sup>1</sup>

1.広島大学大学院理学研究科地球惑星システム学専攻

1.Department of Earth and Planetary Systems Science, Graduate School of Science, Hiroshima University

Subducted oceanic plates should be one of the significant factor deciding structures and the evolution of the Earth's lower mantle. The typical large and strong heterogeneity is large low shear velocity provinces on the lowermost mantle indicated by seismic tomography models. Seismic analyses indicate that these structures have prominent features, such as steep edges, a correlation between margins and hot spot locations, and the stability over a long term. We investigate that influence of the lower mantle properties on the mechanical interactions between the subducted plate and the compositionally dense layer, using mantle convection model in which plate-like motion is realized without any forces imposed on the surface plate.

Our 2-D Cartesian numerical model has chemically distinct materials with the density contrast of  $+77.3 \text{ kg m}^{-3}$  on the core mantle boundary (CMB) and the post-perovskite phase change with the Clapeyron slope of  $+8 \text{ MPa K}^{-1}$  and the density contrast  $+1.4\%$ . In this study, we focus on the following parameters that affect the slab subducting history and the evolution of the dense layer; depth dependence of the thermal expansivity, the depth profile of the background mantle viscosity and the yield strength of the slab. We also incorporate phase diagrams of hydrous minerals and hydration effects on the density and the viscosity.

In our result with the thermal expansivity decreasing with the depth, the dense material piles can remain on the CMB during the calculation. Moreover, the thermal expansivity depending on the depth makes the slab velocity slow down, even when the viscosity is small. This effect leads to the convection layered at the boundary between the regular mantle and dense segments, such that the slow convection is generated in the regular mantle layer and the active convection is in the dense layer. Steep difference in the temperature overlaps at the chemical boundary between dense materials and the surrounding mantle. Upwelling plumes are generated on the top of the dense layer. These plumes entrain only small amount of the dense material. On the other hand, the constant thermal expansivity destabilizes the dense layer. The dense segments deform strongly and rise off the CMB by the subducted slab plunging into the dense material piles. The subducted slab deformation near the 660 km depth discontinuity occurs except when the thermal expansivity is constant and the yield strength is 300 MPa. The amplitude and the wavelength of the slab buckling become larger with lower mantle viscosity. This deformation influences contact area between the dense structure and the slab, and spatial distribution of water in the lower mantle. We did not find significant effects of the slab buckling on the evolution of the lowermost mantle structures in the long term.

キーワード: subducted slab、LLSVPs、thermal expansivity

Keywords: subducted slab, LLSVPs, thermal expansivity

Effects on onset and stability of plate-like behavior in the global mantle water cycle  
Effects on onset and stability of plate-like behavior in the global mantle water cycle

中川 貴司<sup>1</sup>、\*中久喜 伴益<sup>2</sup>

Takashi Nakagawa<sup>1</sup>, \*Tomoeki Nakakuki<sup>2</sup>

1.海洋研究開発機構数理科学・先端技術研究分野、2.広島大学大学院理学研究科地球惑星システム学専攻  
1.JAMSTEC/MAT, 2.Hiroshima Univ.

In our previous study, the rheological properties of hydrous mantle minerals would have the negative thermal feedback (high heat flow across the oceanic lithosphere and colder mantle temperature when the rheological strength of hydrous mantle minerals is strong enough) [Nakagawa et al., 2015]. However, the yield strength in that study was not consistent with that found from deformation experiments [Kohlstedt et al., 1995]. Moreover, the hydrous oceanic crust might occur to the lubrication that could enhance the activity of plate-like behavior caused by mantle volatiles [Cramer et al., 2012; Bercovici, 1998]. Here we examine to find the stable plate tectonics-like behavior from weak to strong oceanic lithosphere that seems to be consistent with the laboratory experiments in numerical mantle convection simulations with water migration [Nakagawa and Speigleman, to be submitted]. Two water cycle regime with the stable plate-like behavior would be found as a function of ductile yield strength and friction coefficient of brittle deformation, which would be 'regassing dominated cycle' for weaker oceanic lithosphere and 'balanced water cycle' for stronger oceanic lithosphere because the faster plate velocity could be expected for the weaker oceanic lithosphere than for the stronger oceanic lithosphere. With the constraint of mantle water content [Hirschmann, 2006; Hirschmann and Kohlstedt, 2012], the balanced water cycle regime would be preferable for reconciling the mantle water cycle. The onset timing of plate-like behavior would be much earlier than the dry mantle convection case with pseudo-plastic yielding rheology.

キーワード：水、マントル対流、プレートテクトニクス、降伏応力

Keywords: water, mantle convection, plate tectonics, yield strength

## 中心核形成における内部不均質形成仮説の数値シミュレーション

Numerical tests on hypothetical primitive origins in a growing planet with core formation

\*古市 幹人<sup>1</sup>、中川 貴司<sup>1</sup>\*Mikito Furuichi<sup>1</sup>, Takashi Nakagawa<sup>1</sup>

## 1. 海洋研究開発機構

1. Japan Agency for Marine-Earth Science and Technology

The presence of large-scale heterogeneity in the deep mantle has been proposed on the basis of the interpretation of strong seismic anomalies in the present Earth's mantle. We hypothesize that the origin is in the protoplanetary material that survived the late accretion stage of planetary formation. In order to examine this hypothesis, we performed the 3-D Stokes flow simulation under a self-gravitating field with a free-surface treatment. The numerical calculation starts with a Mars-sized protoplanet (PP) which may serve as the sources of primitive reservoir by assuming that it has the undifferentiated material or primitive mantle differentiated in an early oxidized state. From this Mars-sized initial embryo, the planet grows under continuous impacts which create local magma ponds which eliminate the chemical signature of PP. We found that generations of such melt ponds may not completely remove the PP material from the growing planet. We suggest that such remaining PP material may be the origin of the primitive reservoir in the deep mantle inferred from geochemical analyses. Our analysis may contribute to the understanding of the thermochemical structure of the early Earth.

キーワード：コア形成、不均質領域、ストークス流れ

Keywords: core formation, primitive reservoir, Stokes flow