

Advent of Continent: Evidence from the Nishinoshima Volcano in the Ogasawara Arc

*Yoshihiko Tamura¹, Osamu Ishizuka^{2,1}, Tomoki Sato¹, Alexander Nichols¹

1.R & D Center for Ocean Drilling Science, Japan Agency for Marine-Earth Science and Technology,
2.Geological Survey of Japan/AIST

Nishinoshima, one of the submarine volcanoes in the Ogasawara Arc, ~1,000 km south of Tokyo, Japan, suddenly erupted in November 2013, after 40 years of dormancy. The Nishinoshima volcano might represent the missing link between the mantle and the continental crust because (1) Nishinoshima, whose underlying crust is only 21 km thick, is one of the world's closest volcanoes to the mantle, and (2) the lavas have been andesites and were similar in composition to the continental crust. Here we report the scientific results of our endeavours to collect lavas from the currently erupting lava flows on the surface to the submarine lavas of the Nishinoshima volcano. Using olivine-bearing phenocryst-poor andesite samples, we developed a mantle-derived andesite model for the genesis of the Nishinoshima volcano. Shallow and hydrous mantle melting is necessary to produce primary andesite magmas, and thus it is only achieved beneath Nishinoshima and submarine volcanoes in the Ogasawara arc, where the crust is thin. We also show that the primary magma composition change from basalt produced at a considerable depth beneath the old thick lithosphere to andesite produced beneath the present crust corresponds to the thermal evolution of the mantle wedge in the Ogasawara arc.

Keywords: andesite, continental crust, oceanic arc



Izu-Bonin-Mariana Arc basement from IODP Exp. 351(Amami Sankaku Basin)

*Osamu Ishizuka¹, Rosemary Hickey-Vargas², Gene Yogodzinski³, Masafumi Sudo⁴, Anders McCarthy⁵, Ivan Savov⁶, Yuki Kusano¹, Richard Arculus⁷, Kara Bogus⁸

1.Geological Survey of Japan, AIST, 2.Florida International University, 3.University of South Carolina, 4.Universitaet Potsdam, 5.University of Lausanne, 6.University of Leeds, 7.Australian National University, 8.Texas A&M University

IODP (International Ocean Discovery Program) Exp.351 targeted, in particular, evidence for the earliest evolution of the Izu-Bonin-Mariana (IBM) arc system following inception. This can be obtained by identification and exploration of regions adjacent to an arc, where unequivocally pre-arc crust (basement) overlain by undisturbed arc-derived materials exists. The drill site (U1438) is located in Amami Sankaku Basin (ASB), west of the Kyushu-Palau Ridge (KPR), i.e., paleo-IBM arc. Exp.351 successfully recovered pre-Izu-Bonin-Mariana arc basement, a volcanic and geologic record spanning pre-Arc, arc initiation to remnant arc stages. In this contribution, age and geochemical characteristics of arc basement and its tectonic significance will be discussed. At U1438, 1611m thick section in total was cored, comprising 1461m of sediment section and 150m of igneous basement. Numerous flow contacts were identified in the basement, several chilled margins could be identified but low recovery in basement rock makes their interpretation in terms of volcanic stratigraphy difficult. Paleomagnetic measurements show consistent shallow plunges of maximum anisotropy axes, and that implies these basalts were emplaced as sheet flows rather than pillow lavas.

Basalts have diverse textures that include aphyric to subophitic/ophitic. Phenocrysts are present in approximately half of the basalts and consist of plagioclase, clinopyroxene, titanomagnetite, spinel and minor olivine.

Age interpretations based on biostratigraphy (Arculus et al., Nat Geosci, 2015) determined that the basement section is between 64 and 51 Ma, and direct age determination of basement basalt by Ar/Ar dating is under way.

Whole rock geochemical analysis of the basement basalt shows that these basalts are low K tholeiitic basalts. They mostly have high-MgO (generally >8 wt%), low-TiO₂ (0.6-1.1 wt%), Ti/V, low-Zr (mostly <50 ppm). On Ti/V plot, the U1438 basalts and forearc basalt (FAB) which is supposed to be associated with subduction initiation from IBM arc are similar to each other and clearly distinct from Philippine Sea backarc basin basalts and normal MORB by having low Ti/V.

One prominent characteristic of the basalts is their depletion of immobile highly incompatible elements compared with MORB, e.g., they are strikingly light REE (LREE) depleted. But La/Nd ratio and Th/LREE increases upcore in the uppermost part of the basement, and these trace element variations with depth is accompanied by variation of major element composition such as Si and Ti. Hf-Nd isotopes for the basement basalts show a significant range of compositions, and relatively radiogenic Hf compared to Nd indicates an Indian Ocean-type MORB source. However, the dominant signature, with $e_{\text{Hf}} > 16.5$, is more radiogenic than most Indian MORB.

Preliminary data suggests that the basement basalts are relatively primitive melts and likely derived from Indian Ocean-type MORB sources that are more strongly depleted in terms of incompatible trace elements than typical MORB or backarc basin basalt in the Philippine Sea. Geochemical variation in the uppermost part of the core might imply variability of slab-derived enrichment and/or fertility of mantle at the onset of subduction.

Geochemical characteristics of arc basement from Exp.351 imply that depleted MORB-like basalt ("FAB") appears to have formed in wider area than previously thought, i.e., including "reararc" side

of ancient IBM arc. However, critical assessment of genetic relation between FAB and basement basalt of ASB requires precise age determination to constrain tectonic setting where the ASB basalts formed.

Keywords: arc basement, Izu-Bonin-Mariana arc, geochemical characteristics, Ar/Ar age

Physical properties of fore-arc basalt and boninite recovered by IODP EXP352 and its significance for the seismic velocity structure in the oceanic crust

*Katsuyoshi Michibayashi¹, Mutsumi Honda², Yuzuru Yamamoto³, Nana Kamiya⁴, Tetsuya Sakuyama⁵, Tohru Watanabe⁶

1.Institute of Geosciences, Shizuoka University, 2.Graduate School of Integrated Science and Technology, Shizuoka University, 3.Department of Mathematical Science and Advanced Technology, JAMSTEC, 4.Graduate School of Integrated Sciences, Nihon University, 5.Faculty of Science, Osaka City University, 6.Graduate School of Science and Engineering, University of Toyama

Physical properties of the cores recovered by IODP EXP352 were characterized through a set of measurements on whole core sections and discrete samples. Gamma ray attenuation density (GRA), magnetic susceptibility (MS) and P-wave velocity (PWL) of the cores were obtained using the Whole-Round Multisensor Logger (WRMSL). Natural gamma radiation (NGR) was measured by the Natural Gamma Radiation Logger (NGRL). Point magnetic susceptibility (PMS), reflectance spectroscopy and colorimetry (RCS) data were acquired using the SHMSL. Thermal conductivity (TCON) and moisture and density (MAD) were obtained on sections and discrete samples, respectively. All raw data were subsequently "filtered" to remove spurious points that correspond to empty intervals in the liner or broken pieces. We have mainly studied the relationships among P-wave velocity, density, porosity and magnetic susceptibility. We show that the physical properties are useful dataset for the interpretation of the layer 2 (i.e. volcanic rocks) in the seismic velocity structure of the oceanic crust.

Keywords: IODP EXP352, Physical property

Evolution process of volcano-bounded basin revealed by mapping of seismic reflectors associated with geological boundary from drilling results in Izu rear arc

*Mikiya Yamashita¹, Yoshihiko Tamura², Narumi Takahashi¹, Seiichi Miura¹, Shuichi Kodaira¹

1.Japan Agency for Marine-Earth Science and Technology, 2.ODS, Japan Agency for Marine-Earth Science and Technology

International Ocean Discovery Program (IODP) has successfully conducted the first rear-arc drilling in the Izu-Ogasawara (Bonin) intra-oceanic arc at Site U1437 in 2014. The drilling purpose of Site U1437 is to reveal the formation of oceanic arc crust and its evolution into continental crust about the history of "the missing half" of the subduction factory. Site U1437 is located at the volcano-bounded basin between the Manji and Enpo backarc seamount chains in the Izu rear arc and ~90 km west of the arc-front volcanoes Myojinsho and Myojin Knoll, at 2117 m below sea level. Site U1437 had excellent core recovery in Holes U1437B and U1437D, and we succeeded in hanging the longest casing ever in the history of R/V *JOIDES Resolution* scientific drilling (1085.6 m) in Hole U1437E and cored to 1806.5 mbsf.

In order to evaluate the crustal structure of this proposed site before the IODP drilling, Japan Agency for Marine-Earth Science and Technology carried out many seismic reflection and refraction surveys using R/V *Kaiyo* and *Kairei* in the Izu rear arc during 2006 to 2008. Five clear seismic reflection profiles consisting of three kinds of survey environments and one seismic velocity image by seismic refraction survey are obtained across the drilling site U1437. Five sedimentary units consisting of volcanoclastics are identified from our interpretations around the drilling site in the seismic reflection profiles over the 5 km/s and 6 km/s iso-contours of P-wave velocity obtained by the velocity image of seismic refraction survey in order to evaluate structure of the drilling location before drilling. However, some unit boundary is not recognized from the drilling core. It means the difficulties for identification the geological target from only seismic images in volcanic regions. According to the drilling results, the acquired geological core is consisted of seven lithological units (I, II, III, IV, V, VI and VII). Units I to V was produced at age of 0-9 Ma. The Unit VI and VII, 1320-1806.5 mbsl, have the ages ranging from 9 to ~14 Ma. P-wave velocity calculated from obtained core samples increases downhole from ~1500 to ~4500 m/s which agree with the range of our velocity analysis. Tops of unit II, V and VII correspond to the strong reflector of seismic profiles. We interpreted and mapped using grid survey data around the drilling site. The top of unit II which corresponds to the volcanoclastics has south dipping trend. The top of unit V which corresponds to the mudstone with volcanoclastics layer has south dipping trend. The top of unit VII which corresponds to volcanoclastics estimated by near-vent deposit has southeast dipping trend. Each reflector is interrupted by igneous basement near seamount chains. We also calculate the thickness from unit VII to V and unit II to seafloor from mapped reflectors. These features suggest the different activity of various directions in the Izu rear arc. We will discuss about the evolution process in Izu rear arc deduced from our mapping results.

Keywords: site survey, volcano-bounded basin, rear arc

Preliminary reports on the Nature of the Lower Crust and Moho at Slower Spreading Ridges (SloMo)

Henry J.B. Dick², Christopher J. MacLeod³, Peter Blum⁴, *Tomoaki Morishita¹, Natsue Abe⁵, Toshio Nozaka⁸, Alessio Sanfilippo^{1,6}, Scientific Party Expedition 360⁷

1.Kanazawa University, 2.Woods Hole Oceanographic Institute, 3.Cardiff University, 4.Texas A & M University, 5.JAMSTEC, 6.University of Pavia, 7.Expedition 360 Scientific Party, 8.Okayama University

IODP Expedition 360, Phase I of the Nature of the Lower Crust and Moho at Slower Spreading Ridges (SloMo) project of a Multi-Leg Drilling Project, was carried out from 1 Dec., 2015 to 31 Jan, 2016, at Atlantis Bank that is a 5 km local high along the eastern wall of the Atlantis II Transform of the Southwest Indian Ridge. Atlantis Bank has been interpreted as an oceanic core complex, where intact lower crust and the uppermost mantle are tectonically exposed by a long-lived detachment fault. We conducted all drilling operations at a single site in a single Hole U1473A and drilled 789.7 m through gabbros. This is the deepest single-leg hard-rock drilling hole in ocean crust. Expedition 360 Hole U1473 is located at 2.2 km Northeast of 1.5 km deep Hole 735B and at 1.4 km north of 158 m deep Hole 1105A. This provides us, for the first time, a unique opportunity to explore three dimensional lower crustal characteristics beneath the slow-spreading ridges. Phase II of the SloMo has proposed to drill 6 km through MOHO by the CHIKYU. In the meeting, we will introduce preliminary results of IODP Expedition 360 and the future perspective leading to Phase II of the SloMo, a mantle drilling into ultraslow-spreading ridges.

Expedition 360 Scientific Party: Donna K. Blackman, Julie A. Bowles, Michael J. Cheadle, Kyungo Cho, Jakub Ciazela, Jeremy R. Deans, Virginia P. Edgcomb, Carlotta Ferrando, Lyderic France, Biswajit Ghosh, Benoit M. Ildefonse, Mark, A. Kendrick, Juergen H. Koepke, James A.M. Leong, Chuanzhou Liu, Qiang Ma, Antony Morris, James H. Natland, Oliver Pluemper, Jason B. Sylvan, Maurice A. Tivey, Riccardo Tribuzio, Gustavo Viegas

Keywords: Ultraslow-spreading ridge, Oceanic Core complex, Oceanic Lower Crust

Lord Howe Rise Drilling: Deep stratigraphic record for the Cretaceous eastern Gondwana margin

*Saneatsu Saito¹, Ron Hackney², Yasuhiro Yamada¹, Kliti Grice³, Junichiro Kuroda¹, Jessica Whiteside⁴, Marco Coolen³, Fumio Inagaki¹, Richard Arculus⁵, Dietmar Müller⁶

1.Japan Agency for Marine-Earth Science and Technology, 2.Geoscience Australia, 3.Curtin University, 4.University of Southampton, 5.Australian National University, 6.University of Sydney

The Lord Howe Rise (LHR) is an elongate ribbon of submerged and extended continental crust that separated from Australia during the Late Cretaceous. Current knowledge of the LHR is based only on widely-distributed marine and satellite geophysical data, limited dredge samples, and sparse shallow drilling into Cenozoic pelagic sediments. Building more detailed knowledge of LHR geology and the evolution of the eastern Gondwana margin requires drilling into rocks that record the tectonic and climatic history of the region.

Geoscience Australia and JAMSTEC are leading an international effort to promote an IODP project (871-CPP) to drill a deep stratigraphic hole through a LHR rift basin using D/V CHIKYU that will recover Cretaceous and older sediments and basement rocks. The objectives of this proposal are to: 1) define the role and importance of continental crustal ribbons, like the LHR, in plate tectonic cycles and continental evolution; 2) recover new high-latitude data in the southwest Pacific to better constrain Cretaceous paleoclimate and linked changes in ocean biogeochemistry; and 3) test fundamental evolutionary concepts for sub-seafloor microbial life over a 100-million-year timeframe.

Keywords: IODP, Lord Howe Rise, Chikyu, Cretaceous, Gondwana, Zealandia

Seismological structure of oceanic lithosphere inferred from Po/So waves

*Azusa Shito¹, Daisuke Suetsugu², Takashi Furumura³

1.Institute of Seismology and Volcanology, Kyushu University, 2.Department of Deep Earth Structure and Dynamics Research, Japan Agency for Marine-Earth Science and Technology, 3.Earthquake Research Institute, University of Tokyo

Oceanic lithosphere, comprising oceanic crust and upper mantle, plays an important role in plate tectonics. Knowledge of the detailed structure of oceanic lithosphere is key to understanding its origin.

The propagation of Po/So waves over large distances across oceanic lithosphere provides information about lithospheric structure. Po/So waves, characterized by a high frequency, large amplitude, and long duration, were identified as early as 1935 [Linehan, 1940]. Many previous studies have attempted to quantitatively explain the generation and propagation mode of Po/So phases. Although it is generally accepted that Po/So phases are guided waves traveling very efficiently throughout the oceanic lithosphere, the generation and propagation processes of Po/So phases remain unclear.

In the past two decades, there have been great technological advances in computer simulations of high-frequency seismic waves in heterogeneous structures and in broadband seismic observations on the seafloor. These gains motivated us to further investigate the processes relating to guided waves in oceanic lithosphere and the generation of Po/So waves, using broadband seismic data and the numerical Finite Difference Method (FDM) to simulate high-frequency seismic waves. In this presentation, we outline recent progress in the study of Po/So waves.

Shito et al. [2013] reported that Po/So waves are generated by multiple forward scattering of P- and S-waves due to small-scale heterogeneities in oceanic lithosphere. The laterally elongated heterogeneities are described by a von Karman distribution function with a correlation length of 10 km in the horizontal and 0.5 km in thickness, with a velocity perturbation of 2%.

Kennett and Furumura [2013; 2014] and Shito et al. [2015] found that the propagation efficiency of Po/So waves depends on the age of oceanic lithosphere, and that this relationship can be qualitatively explained by thickening of oceanic lithosphere that contains small-scale heterogeneities and a reduction in intrinsic attenuation. Recently, Kennett and Furumura [2015] proposed a new model that the amplitude of such heterogeneities increases with depth to the bottom of the lithosphere. These results suggest that small-scale heterogeneities may form continuously in oceanic lithosphere, from the time of its formation at a spreading ridge, via the solidification of melts in the asthenosphere.

The petrological and mineralogical processes that cause the small-scale heterogeneities remain poorly known. Future studies that combine seismological observations with petrological analyses will yield a greater understanding of the origin of oceanic lithosphere.

Keywords: Po/So waves, oceanic lithosphere, small-scale heterogeneities

The effect of melt density, assimilation, and eruption location on volatile and halogen contents in MORB glasses: an example from IODP Site 1256

*Takashi Sano¹, Shigeru Yamashita²

1.Department of Geology and Paleontology, National Museum of Nature and Science, 2.Institute for Study of the Earth's Interior, Okayama University

A continuous section of ~700 m-thick lava pile at Integrated Ocean Drilling Program Site 1256 provides important depth variation of volatile (H₂O, CO₂, S) and halogen (Cl) contents in fresh glasses to understand style of eruption, condition of magma storage, and degree of hydrothermal assimilation. The lava pile is divided into two groups based on eruption location: off-axis basalts of upper portion (250–534 m beneath seafloor, mbsf) and on-axis basalts of lower portion (534–941 mbsf). Majority of the lava pile is composed of pillow, sheet, and massive lavas, but both groups have short (1–2 m) intervals of hyaloclastite layers (e.g., Wilson et al., 2006, Science). H₂O contents of hyaloclastite samples are distinctly higher than those of lavas, suggesting that the high H₂O would induce explosive eruption to produce the hyaloclastite materials. CO₂/Nb and S/Dy in most fresh glasses are smaller than degassed ratios (e.g., Saal et al., 2002, Nature), and these facts indicate shallow degassing and CO₂ and S losses during transport to the seafloor. Saturation pressures calculated by dissolved H₂O and CO₂ contents are wide range from pressure of seafloor (~25 MPa) to pressure of magma chamber (~60 MPa). The saturation pressure positively correlates with melt density, but obvious differences between off-axis and on-axis samples are not identified. These observations may imply that the melt density is important factor to estimate degassed pressure. When the melt density is low, the melt can ascent from melt lens (top of magma chamber) to shallow place within oceanic crust and highly degassed before reach to seafloor. Conversely, dense melt would not ascent to the shallow place and less degassed at deep level near the melt lens.

The most distinctive character for the Site 1256 glasses is higher Cl/Nb and Cl/K than any other MORB glasses. The strong Cl enrichment is explained by assimilation of highly hydrothermally influenced crust (e.g., Sano et al., 2008, 2011, G-cubed). Beneath the Site 1256, melt lens was very shallow (<1.5 km) and hydrothermal circulation of high-salinity brines would easily reach to roof crust of the melt lens.

Keywords: IODP, volatile, melt density

A seamount on top of Ontong Java Plateau was created by remelting of plateau lithosphere by plate flexure

*Takeshi Hanyu¹, Maria Luisa Tejada¹, Kenji Shimizu¹, Osamu Ishizuka², Jun-Ichi Kimura¹, Qing CHANG¹, Ryoko Senda¹, Takashi Miyazaki¹, Kosuke T. Goto², Akira Ishikawa³

1.Japan Agency for Marine-Earth Science and Technology, 2.Geological Survey of Japan, AIST, 3.The University of Tokyo

The Ontong Java Plateau (OJP) was formed by a main volcanism occurred at ca. 120 Ma, followed by several pulses of the late-stage volcanism on and along the margins of the plateau. The origin of the late-stage volcanism is poorly understood because of limited rock sampling. We present the age and geochemical composition of basalts dredged from Nuugurigia Atoll that stands on the plateau basement, possibly erupted in the late-stage volcanism. The $^{40}\text{Ar}/^{39}\text{Ar}$ age of dredged basalts is ca. 20 Ma, younger than any other known late-stage volcanism on OJP (down to 34 Ma), and is coincided with the collision of the plateau with the proto-Solomon arc. These basalts have enriched isotopic signatures pointing towards EM1 distinct from any other rocks thus far collected from OJP. Moreover, they show unusual trace element composition with Sr enrichment and Zr-Hf depletion relative to the elements with similar incompatibility. Such isotopic and trace elemental feature are shared with quartz-bearing garnet pyroxenite xenoliths rarely found from Solomon Islands (Ishikawa et al., 2007). Ishikawa et al. inferred that such pyroxenite was derived from delaminated granulitic lower crust and was part of the OJP lithosphere underplated beneath the plateau lithosphere via the mantle upwelling responsible for the main plateau-forming volcanism. We suggest that the basalts in the late-stage volcanism at Nuugurigia were products of rejuvenated melting of such pyroxenite. Melt transport through the lithosphere may have been facilitated by plate flexure occurred just south of the atoll when the plateau collided with the proto-Solomon arc around 20 Ma.

Keywords: oceanic plateau, lithosphere, recycled material

Understanding continental crust emplacement: A continental scientific drilling perspective

*Thomas Wiersberg¹, Ulrich Harms¹

1.Helmholtz Centre Potsdam GFZ German Research Centre for Geosciences

Scientific drilling is a critical tool for Earth science to understand crustal evolution and processes. In the recent past, the International Continental Scientific Drilling Program, ICDP, supported several drilling operations in hard rock addressing the formation and emplacement of continental crust from Archean volcanism (Peering into the Cradle of Life, Barberton, South Africa), Paleoproterozoic volcanism (FAR-DEEP, Baltic Shield, Russia), rift volcanism (Krafla, Iceland), hot spot volcanism (Hawaii, USA, Snake River Plain, Idaho, USA) and Himalyan-style thrust sheet emplacement (COSC, Sweden). The results of these studies have significantly improved our understanding in past and present formation of continental crust.

ICDP supports scientific drilling operations that facilitate outstanding science at globally important sites. ICDP brings together scientists and stakeholders from 24 nations to work together at the highest scientific and technical level. Since its founding in 1996, more than 40 drilling projects and 75 planning workshops have been supported by ICDP worldwide. The outcomes of scientific drilling operations supported by ICDP cover the full range of the Earth sciences from climate change, natural hazards and earth resources to the origins and evolution of life on Earth, effectively addressing the needs of our growing population for energy, sustenance, and quality of life.

Forthcoming ICDP drilling into the Samail ophiolite complex in Oman will provide key data on melt extraction processes from the mantle, igneous accretion of oceanic crust, and hydrothermal modification of that crust. Drilling will also investigate present day alteration processes of mantle peridotites and their relationship to the deep biosphere.

The Surtsey volcano drilling in fall 2016 aims to investigating processes of rift zone volcanism, hydrothermal alteration and biological colonization of basaltic tephra. It will refine our understanding of seawater- interactions with magma and rock that influence diagenetic and microbial alteration of tephra to produce lithified tuff and shed new light on how rift zone volcanic islands form, lithify, and are ultimately destroyed.

ICDP is also funding drilling for Reservoir Triggered Seismicity near the Koyna dam located close to the west coast of India. The basement rocks of the Koyna-Warna region consists of Precambrian basement overlain by a more than 1 km thick Deccan Trap cover that erupted about 65 Ma ago. Two pilot boreholes will provide critical information on the in-situ stress regime, pore fluid pressure, fluid/gas properties and hydrological parameters of basement rocks, and the geothermal regime. These 3 km deep wells will lay the ground for two deep main boreholes for penetrating into the source of induced seismicity at more than 5 km depth.

Keywords: ICDP, Scientific Drilling, Continental Crust emplacement

From Birth to Death: Fate of The Extremely High-T Subduction Zone of The Oman Ophiolite

*Susumu Umino¹, Yuki Kusano², Keitaro Kitamura³, Kyoko Kanayama⁴, Kazuyha Nagaishi⁵, Tsuyoshi Ishikawa⁶, Ryo Otsuka¹

1.Department of Earth Sciences, Kanazawa University, 2.Geological Survey of Japan, AIST, 3.Asia Air Survey Co., Ltd., 4.Tottori Prefectural Museum, 5.Marine Works Japan Ltd., 6.Kochi Institute for Core Sample Research, JAMSTEC

The Oman Ophiolite preserves the entire geological records of intra-oceanic subduction zone formation and arc evolution, where 98-96 Ma MORB-like basalt magmatism (V1) was followed by 96-94 Ma arc tholeiitic and low-silica boninitic volcanism (V2) [1,2,3,4]. The remnants of the subducted slab are now preserved as the high-grade metamorphic soles beneath the ophiolite sheets [5]. After a quiescence period, alkali basalt flows (V3) were extruded at about 90 Ma, which has an intermediate geochemical characteristics between OIB and EMORB [6]. The V3 trace element compositions can be reproduced by pooled partial melts in the stability field of garnet and spinel lherzolite. Through the V2 magmatism, the source mantle shows progressive depletion by stepwise melt extraction, as shown by the lower Nb/Ta ratios for the younger volcanic rocks (V2 boninite < V2 tholeiite < V1) [2] with identical $\epsilon_{\text{Hf}}(t)$ values. V2 glasses have higher B, Pb, and LILEs with age, indicating an increasing contribution of slab fluids from earlier arc tholeiite to later boninite. Boninite magma was generated with the supply of high-T hydrous fluid and sedimentary melt liberated from the metamorphic sole as demonstrated by the Sr-Nd isotopic compositions of the amphibolite and metachert in the sole and clinopyroxene separates from boninites [1,2,7]. Therefore, the metamorphic sole beneath the ophiolite sheets are responsible for the generation of V2 arc magmas. Melt inclusions in Cr spinel derived from boninite comprise homogeneous glass of mostly low-Si boninitic [3,4] and slightly differentiated composition, with SiO₂ ranging in 52-62 wt% and MgO up to 16 wt%. The primary boninite magma assumed as the most magnesian melt inclusion can coexist with mantle olivine and orthopyroxene [9] at 0.4-0.6 GPa and 1350°C. This T-P conditions indicate a segregation depth of ~17 km from the mantle with a potential T of 1400°C. Meanwhile, the peak metamorphic conditions for the subducted slab that liberated high-T fluids to form boninite and arc tholeiite magmas are 770-900°C and 1.1-1.3 GPa [1,6,7].

The keys of the Oman subduction zone are 1) the preservation of diapiric structures in the mantle [10], 2) short time interval of the V1 and V2 magmatism <2 m.y., 3) T-P conditions for a primary low-Si boninite (Umino et al., 2014); and 4) extremely high T & low-P metamorphic conditions for the sole that liberated the fluids generating the V2 magmas [1,5]. These lines of evidence are most readily explained by intraoceanic thrusting initiated near the ridge axis that developed into a shallow and hot subduction zone [1,2,8,10].

Forced subduction of such an extremely high-T, buoyant slab suppressed convection in the mantle wedge, resulted in the progressive depletion of the source mantle through the V2 arc magmatism. Numerical modeling suggests that melting of the slab and mantle wedge occurs only in the early stage and ceases as the mantle wedge cools because of the absence of convection [10]. Consequently, the Oman arc volcanism terminated in only a few million years. After several million years, parts of the subducted slab delaminated and induced upwelling and adiabatic melting of DMM asthenosphere, resulted in the generation of alkalic magmas of V3 extruded onto thick pelagic sediments before colliding onto the Arabian Peninsula.

[1] Ishikawa et al., doi:10.1016/j.epsl.2005.09.049 [2] Kusano, Y. et al.

<http://dx.doi.org/10.1144/SP392.9> [3] Kanayama et al., doi: 10.1111/iar.12000 [4] Umino et al.,

doi: 10.1130/G36191.1 [5] Cowan et al., <http://dx.doi.org/10.1144/SP392.8> [6] Umino,

doi:10.1029/2012JB009198 [7] Kusano et al., 2015. Goldschmidt, Prague, p. 1725 [8] Putirka, doi: 10.2138/rmg.2008.69.3. 2008 [9] Nicolas, 1989. Kluwer Academic Publication [10] Gutcher et al., doi:10.1130/0091-7613(2000)28<535:CSMBCB>2.0.CO;2

Keywords: Oman Ophiolite, boninite, high-T subduction zone, metamorphic sole, subduction initiation, obduction

The redox state in the Fizh mantle section, the northern Oman ophiolite as an analog of mantle wedge in subduction zone

Masafumi Tani¹, *Eiichi TAKAZAWA^{1,2}, Madhusoodhan Satish-Kumar¹, Masahide Akasaka³, Yoshihiko Tamura²

1.Department of Geology, Faculty of Science, Niigata University, 2.R & D Center for Ocean Drilling Science, Japan Agency for Marine-Earth Science and Technology, 3.Department of Geoscience, Shimane University

The redox state of the upper mantle is important because it controls phase equilibrium and partitioning of some trace elements. Previous studies of mantle peridotites show mantle xenoliths in arc lavas are more oxidized than abyssal peridotites on the ocean floor because subducting slab is dehydrated to supply 'oxidized fluid' such as water and carbon dioxide to the mantle wedge in hanging wall (e.g., Parkinson and Pearce, 1998).

The mantle section of the Oman ophiolite is a good analog for a mantle wedge because it experienced supra-subduction zone setting during oceanic thrusting prior to the obduction. In this study, we focus on the Fizh mantle section in the north Oman ophiolite as a proxy of mantle wedge in a subduction zone and evaluate the redox state recorded in the peridotites. We calculated $\Delta \log f_{O_2}$ (FMQ) using equation of Ballhaus et al. 1991, chemical composition of olivine and spinel with ferric and ferrous iron ratio assuming stoichiometry. The spatial distribution of $\Delta \log f_{O_2}$ (FMQ) in the Fizh mantle section indicates that the basal part of the mantle section is more reduced whereas the upper part of the mantle section beneath the Moho is more oxidized. This may indicate that the basal part of mantle wedge above the contact to subducting slab is most reduced.

We compared ferric and ferrous iron ratio calculated from spinel composition assuming perfect stoichiometry to those directly obtained by Mössbauer spectrometry. The results show that the difference is minor and does not significantly affect the tendency mentioned above. As independent approach, we used abundance ratio of vanadium against scandium in whole rock peridotites. Vanadium partitioning between peridotite and melt is sensitive to the change in oxygen fugacity whereas scandium doesn't. Thus the ratio of vanadium against scandium is a good proxy for estimating oxygen fugacity. As a result, $\Delta \log f_{O_2}$ (FMQ) shows negative correlation with the ratios of vanadium against scandium for whole rock peridotites. This result confirms that the basal part of the Fizh mantle section is more reduced than the upper part of the mantle section.

One possible factor for the reduced signature in the basal part of Fizh mantle section is a reaction of peridotite with a reduced melt produced by melting of sediment with reduced carbon subducted beneath the Fizh mantle section. This hypothesis is supported by high Th/Ce ratio in whole rock peridotites and clinopyroxenes with low $\Delta \log f_{O_2}$ (FMQ). The high Th/Ce ratio is one of the features which indicate contribution of oceanic sediments and suggests those influence. The Raman peak of methane has also been detected in the fluid inclusions in the peridotite from the basal part. On the other hand, a peak of methane has not been detected from the fluid inclusion in the peridotites beneath the Moho Transition Zone. The presence of methane in the fluid inclusions from the basal part of mantle section also supports reduced condition at the base. We consider that the methane was derived from reduced carbon in the subducted oceanic sediment during oceanic thrusting of the ophiolite.

Keywords: Oman ophiolite, Redox state, mantle peridotite, mantle wedge, oxygen fugacity, subduction zone