

The Great Sanriku-oki earthquake of 2 March 1933: Source Characterization Based Regional and Far Field Information and Implications for Off-Trench Normal Faulting Tsunami Sources in Japan and Elsewhere

*Stephen Kirby¹, Naoki Uchida², Emile Okal³, Ryota Hino², Norihito Umino², Azusa Nishizawa⁴

1.U.S. Geological Survey, Menlo Park, 2.Faculty of Science, Tohoku Univ., 3.Northwestern Univ., 4.Japan Coast Guard

Three investigations of the great off-trench earthquake of 2 March 1933 have been completed (Uchida and five others GJI Submitted 2015; Okal, Kirby, and Kaligaris, GJI Submitted, 2016; Kirby, Hino, and Nishizawa, in preparation). Analyses of contemporary data from the early 1930's have revealed new details of this seismic source that caused the devastating tsunami waves that struck the Sanriku coast that led to more than 3,000 dead or missing. The source region of this remarkable event was in the off-trench of a sector of the Japan Trench characterized by large off-trench gravity and bathymetric anomalies and relatively long and straight normal fault scarps parallel to the relevant trench. These conditions are satisfied in Japan in the Hokkaido-Kurile Trench system south of Hokkaido Island and the southern Kurile Islands and in some sectors of the Bonin and Marianas subduction systems. Analysis of fault scarps and curvature changes in the off-trench seafloor and plate convergence rates suggest that recurrence times for great off-trench normal-faulting range from about 1,000 to 10,000 years, roughly 10 times those for interplate thrust earthquake sources.

Keywords: Japan Trench, Normal fault earthquakes, Tsunami sources

Seismicity and its relation to the bending-related faults in the incoming Pacific Plate along the Japan Trench

Seismicity and its relation to the bending-related faults in the incoming Pacific Plate along the Japan Trench

*尾鼻 浩一郎¹、藤江 剛¹、小平 秀一¹、高橋 努¹、山本 揚二郎¹、佐藤 壮¹、山下 幹也¹、海宝 由佳¹、中村 恭之¹、三浦 誠一¹

*Koichiro Obana¹, Gou Fujie¹, Shuichi Kodaira¹, Tsutomu Takahashi¹, Yojiro Yamamoto¹, Takeshi Sato¹, Mikiya Yamashita¹, Yuka Kaiho¹, Yasuyuki Nakamura¹, Seiichi Miura¹

1. 海洋研究開発機構 地震津波海域観測研究開発センター

1. Research and Development Center for Earthquake and Tsunami, Japan Agency for Marine-Earth Science and Technology

After the 2011 Tohoku-Oki earthquake (Mw 9.0), many M7-class normal-faulting earthquakes occurred in the trench-outer rise region seaward of the largest co-seismic slip area during the 2011 Tohoku-Oki earthquake. Outer-rise normal-faulting earthquakes likely relate to the normal-faults, which cut the oceanic crust and form the horst and graben structures. However, relationships between the outer-rise seismicity and crustal structures are not clearly understood from the observations using land seismic network since the outer-rise area is far away from the coast. After the 2011 Tohoku-Oki earthquake, we have conducted repeated ocean bottom seismograph (OBS) observations in the trench-outer rise region along the Japan Trench. Based on the OBS observations, shallow seismicity within the oceanic crust coincides with the seafloor topographic lineation. Their orientations are both parallel and oblique to the trench axis. The trench-parallel topographic lineation is related to the horst and graben structures formed in the trench outer-rise region prior to the subduction. On the other hand, the topographic lineation oblique to the trench axis is almost parallel to the magnetic anomalies. The seismicity along the topographic lineation oblique to the trench axis suggests reactivation of the pre-existing structures formed at the mid oceanic ridge. The results of the OBS observations indicate both pre-existing and newly created faults within the oceanic plate have actively deformed in the trench-outer rise region. Furthermore, close-up OBS observations in a westward dipping normal fault detected concentrated seismicity west of the fault escarpment on the seafloor. Although we could not obtain a clear image of the normal fault within the oceanic crust in the outer-rise of the Japan Trench through seismic reflection surveys, the seismicity could provide some information of the fault geometry within the crust.

キーワード: normal faults、trench outer rise、horst and graben

Keywords: normal faults, trench outer rise, horst and graben

Local heat flow variations seaward of the Japan Trench: Implications for development of fractures in the oceanic crust

Local heat flow variations seaward of the Japan Trench: Implications for development of fractures in the oceanic crust

*山野 誠¹、川田 佳史²、濱元 栄起³、後藤 秀作⁴

*Makoto Yamano¹, Yoshifumi Kawada², Hideki Hamamoto³, Shusaku Goto⁴

1.東京大学地震研究所、2.東北大学災害科学国際研究所、3.埼玉県環境科学国際センター、4.産業技術総合研究所

1.Earthq. Res. Inst., Univ. Tokyo, 2.IRIDEs, Tohoku Univ., 3.Center Environ. Sci. Saitama, 4.Geol. Surv. Japan, AIST

Anomalous heat flow values, higher than that expected for the seafloor age, are pervasively observed on the seaward side of the Japan Trench (Yamano et al., 2014). Appreciably high values were obtained within about 150 km of the trench axis, indicating that the anomaly is related to deformation of the incoming Pacific plate associated with subduction. The broad high heat flow zone seaward of the trench can be attributed to efficient vertical heat transport by pore fluid circulation in a permeable layer in the oceanic crust which thickens toward the trench through fracturing due to plate bending (Kawada et al., 2014). Overlapping the broad anomaly, local variations at a scale of a few kilometers were detected through concentrated measurements at some sites. Such short-wavelength anomalies cannot have their origin deep in the plate and may arise from heterogeneity of the oceanic crust.

For investigation of the nature and the origin of the local anomalies, we conducted dense heat flow surveys around 39°N in 2014 and 2015. Most of the measurements were made along an E-W pre-existing multichannel seismic survey line (JAMSTEC SR101), perpendicular to the trench. A detailed heat flow profile was obtained around 60 to 80 km from the trench axis, in the region where immature horst and graben structures are found with no significant surface displacement. Heat flow values along this 20 km transect range from 60 to 110 mW/m² and show prominent sawtooth-like variations at a scale of 3 to 5 km. Heat flow variation in the N-S direction, parallel to the trench, was also examined by closely-spaced measurements along a short line crossing the E-W transect at one of the heat flow peaks. The values obtained along the 5 km N-S line has a very high average, about 100 mW/m², with a lateral variation as large as the variations along the E-W line. It suggests that high anomalies in this region have elongated shapes extending in the direction parallel to the trench.

The local heat flow variations may have resulted from heterogeneous development of fractures in the oceanic crust. We conducted numerical modeling of heat transport by pore fluid circulation and found that local development of fractures, i.e. sharp lateral variation in the thickness of the permeable layer, would yield heat flow anomalies at the scale of km, with elevated heat flow above more developed fractures. The observed heat flow distribution may therefore indicate that there exist a series of N-S extending well-fractured zones with spacing of several km. Two-dimensional survey of the heat flow distribution on the outer rise would provide information on fracturing process in the oceanic crust.

キーワード：日本海溝、太平洋プレート、熱流量、海洋地殻、亀裂、流体循環

Keywords: Japan Trench, Pacific plate, heat flow, oceanic crust, fracture, fluid circulation

Visco-elasto-plastic deformation and hydration of oceanic plates at trench-rise systems

*Manuele Faccenda¹

1.Dipartimento di Geoscienze, University of Padova

Over the last decade, bending-related normal faulting and hydration of the incoming oceanic plates has been widely observed at trench-rise systems by geophysical surveys. Along with these observations, 2D numerical models of spontaneously bending oceanic plates have contributed to investigate the dynamical evolution of visco-elasto-plastic deformation patterns and the mechanisms of plate hydration by downward seawater percolation along normal faults. In this contribution, I will discuss the main tectonic processes occurring at the trench-rise system within the incoming plate on the basis of the principal geophysical and numerical experiments conducted so far.

Keywords: slab hydration, plate bending, numerical modelling

蛇紋岩の浸透率に基づくアウターライズ断層沿いの海洋マンツルの加水作用

Mantle hydration along the outer rise faults inferred from permeability of serpentinites

*畠山 航平¹、片山 郁夫¹*Kohei Hatakeyama¹, Ikuo Katayama¹

1. 広島大学大学院理学研究科地球惑星システム学専攻

1. Hiroshima university Department of earth and planetary systems science

近年の地震波探査では地震波速度が遅い領域が沈み込み帯のアウターライズ領域で観測されており、海洋マンツルの含水化（蛇紋岩化）が示唆されている（Ranero et al., 2003; Fujie et al., 2013など）。海洋マンツルの含水プロセスについては、アウターライズ断層が海洋リソスフェアを断裂させ、海洋マンツルにおいて断層沿いに蛇紋岩化が進行していると述べられている（Ranero et al., 2003など）。海洋マンツルが従来よりも多く含水しているのならば沈み込む水の総量は多くなることが予想される（Peacock, 2001）。また、海洋スラブ内で発生する稍深発地震は海洋マンツルの脱水と関連していると報告されている（Seno and Yamanaka, 1996）。このように海洋マンツルの蛇紋岩化は地震発生メカニズムと沈み込む水の収支に重要な役割を担っている。

海洋マンツルでの蛇紋岩の広がりについては、蛇紋岩化の反応速度が比較的速いため（Martin and Fyfe, 1970）、マンツル反応前線への水の供給速度に律速されると考えられている（Macdonald and Fyfe, 1985）。マンツル反応前線に供給される水は既存の蛇紋岩を通過する水であるため、蛇紋岩の浸透率が重要となる。そこで本研究では、低温型蛇紋岩の浸透率を測定し、水理拡散率を計算することでアウターライズ断層沿いの蛇紋岩化の広がりを検証した。試料は嶺岡帯に産出する低温型蛇紋岩とパレスベラ海盆、南マリアナ海溝、トンガ海溝からドレッチした試料を用いた。実験は広島大学設置の容器内変形透水試験機を使用し、間隙流体として窒素ガスを用いた定差圧流量法によってダルシーの法則から気体浸透率を測定した。なお、間隙流体に気体を用いるため、気体浸透率からクリンケンベルグ効果を用いて固有浸透率を求めた。

浸透率は封圧5~100MPaで 10^{-16} ~ 10^{-21} m²と封圧増加に伴い低下した。得られた浸透率から、水理拡散率を計算するとトンガ海溝産試料において封圧100MPaで 3.8×10^{-5} m²/secとなる。流体が断層沿いに供給される時間を100万年とすると海洋マンツル最上部で流体は断層沿い両側で17km広がり、深くなるにつれて狭くなっていく。蛇紋岩化作用の反応速度は地質学的にかなり速いため流体の移動距離を蛇紋岩化の広がりともみなすことができる。アウターライズ断層沿いの蛇紋岩化から海洋マンツルが地球内部へ輸送する水の総量を計算すると 2.2×10^{12} kg/yearとなり、海洋地殻が輸送する水の総量（ 1.3×10^{12} kg/year）よりも多いことが予想される。そして、火成活動による地球内部からの水輸送量よりも多いため、現在の海水は減少しているのかもしれない。

キーワード：浸透率、蛇紋岩、アウターライズ断層、水理拡散率、水輸送

Keywords: permeability, serpentinite, outer rise fault, hydraulic diffusivity, water transport

海洋プレート沈み込み直前の加水作用解明に向けた掘削計画概要と展望

An outline of Bend-Fault Hydrology in the Old Incoming Plate (H-ODIN) project and its perspective

*森下 知晃¹、藤江 剛²、小野 重明²、山野 誠³、小平 秀一²、木村 純一²、斎藤 実篤²、阿部 なつ江²、守岩 慈英尊⁴

*Tomoaki Morishita¹, Gou Fujie², Shigeaki Ono², Makoto Yamano³, Shuichi Kodaira², Jun-Ichi Kimura², Saneatsu Saito², Natsue Abe², Jason Morgan⁴

1.金沢大学、2.海洋研究開発機構、3.東京大学、4.ロンドン大学

1.Kanazawa University, 2.JAMSTEC, 3.University of Tokyo, 4.University of London

Crustal hydration at the mid-ocean ridge by hydrothermal circulation has been considered to be the first-order control on the degree of the oceanic plate hydration. Previous ocean drilling projects have revealed hydration processes and their extent of oceanic crust at spreading centers. Recently, hydration due to plate bending-induced normal faults (bend-fault hereafter) in incoming plate just prior to subduction has drawn considerable attention (e.g., Ranero et al., 2003 Nature). In the last decade, a number of geophysical structure studies have been conducted to test the hypothesis that significant hydration is due to plate bending-induced normal faults just prior to subduction (e.g., Fujie et al., 2013 Geophys. Res. Lett.). However, we really do not know what is the bending-induced fault zone. Bend-fault hydration processes depend on various conditions, such as thermal conditions and stress state. Ideally, comparing subduction zones with several different basic states (e.g. Old cold plate vs Young hot plate) will be the most effective approach to expand our knowledge of bend-fault hydration processes. Two new IODP proposals on hydration in incoming plate of middle America site (Morgan et al., 2014, Pre-876: Bend-Fault Serpentinization (BFS): Oceanic Crust and Mantle Evolution from Ridge through Trench) and northwest Pacific site (Morishita et al., 2015 Pre-886: Bend-Fault Hydrology in the Old Incoming Plate) have been submitted. The world's largest dense onland seismic observation network is in NE Japan. In addition, a large number of onland/offshore reflection and refraction seismic surveys have been conducted here, both before and after the 2011 Tohoku earthquake. This network and surveys provides an invaluable data set to study how various subduction zone processes shape subduction inputs, tectonics, and volcanism. The northwestern Pacific is one of the best targets because here horst-and-graben bend-fault structures are the best developed in the world (Nakanishi, 2011, Springer), and this is also one of the world oldest, thus coldest, subducting oceanic plates, hence likely to be associated with the deepest extent of bend-fault serpentinization. The maximum penetration depth of seawater, and thus degree of serpentinization is thought to be inversely proportional to the temperature of the incoming plate, i.e., deeper in cold plate than in hot plate. In the presentation, I introduce the contents and perspective of the new IODP proposals using CHIKYU on hydration of incoming plate prior to subduction.

キーワード：沈み込む海洋プレート、プレート屈曲断層、加水過程

Keywords: Incoming oceanic plate, Bending fault, Hydrology

Regional variations in the nature of the incoming oceanic plate in the NW Pacific margin
Regional variations in the nature of the incoming oceanic plate in the NW Pacific margin

*藤江 剛¹、小平 秀一¹、Grevemeyer Ingo³、Morgan Jason⁴、山田 知朗²

*Gou Fujie¹, Shuichi Kodaira¹, Ingo Grevemeyer³, Jason Morgan⁴, Tomoaki Yamada²

1.海洋研究開発機構、2.東京大学地震研究所、3.GEOMAR、4.Royal Holloway, University of London

1.Japan Agency for Marine-Earth Science and Technology, 2.Earthquake Research Institute, University of Tokyo, 3.GEOMAR, 4.Royal Holloway, University of London

Various subduction zone processes, such as arc magmatism and generation of earthquakes, are directly or indirectly related to the nature of the subducting oceanic plate, indicating that the regional variations in the nature of the incoming oceanic plate may cause the regional variation of these subduction zone processes.

In the northwestern Pacific margin, the old oceanic Pacific plate (120Ma-130Ma) subducts beneath the northeast Japan arc. Many large interplate earthquakes has occurred in this subduction zone but their distribution are not uniform, suggesting that the interplate seismic coupling is not uniform along this subduction zone as pointed out by the geodetic studies.

The interplate seismic coupling is considered to be dependent on the materials that exist at the plate interface. For example, an ocean drilling after the 2011 Tohoku-oki earthquake (JFAST) revealed that the existence of the pelagic clay layer at the plate interface was a key to the large coseismic slip to the trench (Chester et al., 2013).

Several previous seismic structure studies on the forearc region of this region pointed out large interplate earthquakes occurred outside of areas with low seismic velocities and high Poisson's ratio along the plate interface (Wang and Zhao, 2006; Zhao et al., 2011; Fujie et al., 2013).

Similarly, a thin layer with low seismic velocities identified at the depth of the plate interface in a region of low interplate microseismicity (Fujie et al., 2002; Mochizuki et al., 2005). These previous studies suggested that the nature of the incoming Pacific plate, especially its water content, is a key to understanding this regional variation in interplate seismicity.

Since 2009, to investigate structural variations in the incoming Pacific plate, focusing on the amount of water, we conducted extensive active source seismic surveys on the outer rise of the northwestern Pacific margin. We confirmed that seismic velocities gradually reduce and V_p/V_s ratio gradually increase toward the trench axis accompanied by the development of bend faults, suggesting the water penetration into the oceanic plate through the bend faults. In addition, we observed remarkable regional variations in the sediment thickness, crustal thickness, and seismic velocities within the crust and mantle before the area of bend faults.

In this paper, we are going to show these structural variations in the incoming plate and discuss the origin of the variations and possible effects on the subduction zone processes.

キーワード : incoming plate、OBS、petit spot

Keywords: seismic structure, bending-related faults, subduction zone

Evaluation of coseismic physicochemical processes in fault zones based on geochemical analyses of fault rocks

Evaluation of coseismic physicochemical processes in fault zones based on geochemical analyses of fault rocks

*石川 剛志¹

*Tsuyoshi Ishikawa¹

1. 海洋研究開発機構高知コア研究所

1. Kochi Institute for Core Sample Research, Japan Agency for Marine-Earth Science and Technology

Trace element and isotope compositions of rocks record processes associated with their petrogenesis and later modifications. Interactions with fluids and melts are major factors to control chemical characteristics of the rocks. Geochemical analyses of rocks thus can be useful means for understanding nature of such fluid- and melt-related interactions, constituting essential part of broad physicochemical processes occurring in the earth's interior. Here I focus on an attempt to evaluate coseismic physicochemical processes in fault zones based on rock geochemistry.

Frictional heating in a fault zone during earthquake slip affects the slip behavior itself.

Increased temperature on a fault can induce dynamic fault weakening by processes such as pressurization of interstitial fluid by thermal expansion, known as thermal pressurization, and melt lubrication. Recent studies revealed that trace element and isotope analyses of fault rocks combined with geological, mineralogical, structural, and geophysical observations are useful for elucidating such slip weakening processes.

In the Chelungpu fault in Taiwan, which slipped during the 1999 Mw 7.6 Chi-Chi earthquake, the slip-zone rocks showed clear geochemical anomalies, including decreases of Li, Rb, Cs and $87\text{Sr}/86\text{Sr}$ and an increase of Sr relative to adjacent host sedimentary rocks (Ishikawa et al., 2008). Model calculations revealed that these anomalies were produced by coseismic fluid-rock interactions at $>350^\circ\text{C}$, which may have caused a dynamic decrease of friction along the fault through thermal pressurization. Ancient slip zone rocks from a major reverse fault in the Boso Emi accretionary complex at 1–2 km depth (Hamada et al., 2011) and from a decollement of the Kodiak accretionary complex at seismogenic depth (Yamaguchi et al., 2014) showed similar evidence for coseismic fluid-rock interactions at high temperatures. The $87\text{Sr}/86\text{Sr}$ ratios of the Kodiak slip zone rocks also revealed involvement of fluids derived from subducted oceanic crust during the earthquake slip. Recent high-velocity frictional experiments under wet condition demonstrated that detectable fluid-induced geochemical anomaly can be produced by a single earthquake event (Tanikawa et al., 2015). For the slip zone rocks from the Shimanto accretionary complex in Kure area, which represent rocks of ancient megaspray fault at 2.5–5.5 km depth, geochemical signals derived from high-temperature fluids overlap with those from melting, indicating coseismic fluid-rock interactions followed by disequilibrium frictional melting (Honda et al., 2011). These results demonstrate that geochemical characteristics of the fault rocks are useful indicators of such physicochemical events. However, further studies are required especially for fault system developed within basaltic crust, for which geochemical characteristics of fault rocks have been poorly understood.

Refs: Ishikawa et al. (2008) *Nature Geosci.* 1, 679–683; Hamada et al. (2011) *JGR* 116, B01302; Yamaguchi et al. (2014) *EPS* 66:58; Tanikawa et al. (2015) *GRL* 42, 10.1002/2015GL063195; Honda et al. (2011) *GRL* 38, L06310.

キーワード：地球化学、断層岩、地震、流体岩石相互作用

Keywords: geochemistry, fault rocks, earthquakes, fluid-rock interactions

プレート収束域における深部メタンとヘリウムの放出

Deep methane and helium emissions at convergent plate boundaries

*佐野 有司¹、鹿児島 涉悟¹、木下 尚也¹、高畑 直人¹、坂田 将²、ラン テファン³、ウェン シンイ⁴、チン アーチ⁴、トモナガ ヤマ⁵

*Yuji Sano¹, Takanori Kagoshima¹, Naoya Kinoshita¹, Naoto Takahata¹, Susumu Sakata², Tefang Lan³, Hsinyi Wen⁴, Ai-Ti Chen⁴, Yama Tomonaga⁵

1. 東京大学大気海洋研究所海洋地球システム研究系、2. 産業技術総合研究所地質調査総合センター、3. 台湾中央研究院地球科学研究所、4. 国立台湾大学地質科学系、5. スイス連邦水圏科学技術研究所

1. Division of Ocean and Earth Systems, Atmosphere and Ocean Research Institute, University of Tokyo, 2. Geological Survey of Japan, AIST, 3. Institute of Earth Sciences, Academia Sinica, 4. Department of Geosciences, National Taiwan University, 5. Swiss Federal Institute of Aquatic Science and Technology

Methane plays an important role in the greenhouse effect of the atmosphere and in the chemistry of ozone reduction. Major methane emissions into the atmosphere originate from the biosphere (e.g., wetlands, rice paddies and animals), the geosphere (e.g., hydrocarbon basins and geothermal areas) and anthropogenic activity (e.g., natural gas production and distribution, coal mining). Natural methane emission from the geosphere is generally characterized by a radiocarbon-free signature (age > 50 kyrs) and may preserve information on the deep fluid in the Earth's crust. In this work, the origin of methane in forearc and volcanic arc regions at subduction zones, and collision zone at convergent plate boundaries is discussed using the carbon isotope composition of methane together with the abundances and isotope signatures of associated volatile elements such as helium, argon and nitrogen. We collected methane-rich natural gas samples in the South Kanto gas field in Japan and from mud volcanoes in South Taiwan. For comparison, we acquired natural gases in the Akita and Niigata gas fields located in the volcanic arc region of Japan. Chemical composition (CH_4 , C_2H_6 , C_3H_8 , CO_2 , N_2 , O_2 , Ar, and He abundances), carbon and nitrogen isotope signatures ($^{13}\text{C}/^{12}\text{C}$ of CH_4 , $^{15}\text{N}/^{14}\text{N}$), noble-gas isotope ratios ($^3\text{He}/^4\text{He}$, $^4\text{He}/^{20}\text{Ne}$, and $^{40}\text{Ar}/^{36}\text{Ar}$) were measured using a quadrupole mass spectrometer, a continuous flow GC-IRMS system, and a noble gas mass spectrometer, respectively. The methane-rich gas in the South Kanto gas field shows a typical microbial signature characterized by light carbon isotopes, high $\text{CH}_4/(\text{C}_2\text{H}_6 + \text{C}_3\text{H}_8)$ and low $^3\text{He}/^4\text{He}$ ratios, while the natural gases in the Akita and Niigata region show a thermogenic signature with prevalence of heavy carbon isotopes, low $\text{CH}_4/(\text{C}_2\text{H}_6 + \text{C}_3\text{H}_8)$ and high $^3\text{He}/^4\text{He}$ ratios. These observations are consistent with those reported in the literature. On the other hand, methane-rich gases from mud volcanoes in South Taiwan show heavy carbon isotopes, $\text{CH}_4/(\text{C}_2\text{H}_6 + \text{C}_3\text{H}_8)$ ratios between microbial and thermogenic signatures and variable $^3\text{He}/^4\text{He}$ ratios, a part of which cannot be explained by a simple binary mixing of microbial and thermogenic methane. We also measured helium and carbon isotopes of submarine hydrothermal systems close to the Tokara Islands and within Kagoshima Bay. Gas geochemistry of the collected seawater and porewater samples is compared with those from the Japan Trench and the Nankai Trough. Finally, all methane-rich gases from submarine and on-land expressions of fluid emission are discussed within the frame work of the respective geotectonic settings.

キーワード：メタンガス、ヘリウム同位体、プレート境界

Keywords: methane gas, helium isotope, plate boundary

Lithostratigraphic Evolution of the Shikoku Basin, Inputs to the Nankai Trough Subduction Zone

*Michael Underwood¹

1. New Mexico Institute of Mining and Technology

The Shikoku Basin (Philippine Sea plate) hosts inputs of sediments and igneous crust to the Nankai Trough subduction zone of SW Japan. After five decades of scientific ocean drilling (DSDP, ODP, IODP), and three successful transects across the margin (Ashizuri, Muroto, Kumano), we are beginning to recognize and understand the complexities and idiosyncrasies of the basin's evolution. This presentation will focus on three elements of that evolution: (a) time-transgressive facies changes; (b) an enigmatic interval of anomalously high porosity; and (c) systematic, gradual changes in detrital clay mineral assemblages over time.

Perhaps the most noteworthy of several time-transgressive facies changes occurs at the base of the uppermost lithostratigraphic unit (hemipelagic-pyroclastic facies). The base of that unit (otherwise known as Upper Shikoku Basin facies) coincides with the deepest discrete layer of volcanic ash. The age of the unit boundary changes considerably along strike, from 3.9 Ma at ODP Site 1177, to 3.3 Ma at ODP Site 1173, to 7.6-7.8 Ma at IODP Sites C0011 and C0012. Those temporal-spatial differences in the onset of substantial pyroclastic influx might be related to a NE-directed migration of the triple junction that joins the Japan, Izu-Bonin, and Nankai subduction boundaries; that migration should have triggered along-strike changes in explosive arc volcanism. Mudstones within the zones anomalously high-porosity contain unusually large proportions of dispersed volcanic ash (glass shards and pumice). Temperatures at the base of the zones vary from site to site, so thermally driven diagenesis alone cannot account for the anomaly. Instead, dispersed ash affects mudstone microstructure by forming cohesive aggregates that inhibit collapse of intergranular pore space. Partial dissolution of glass shards also contributes silica for precipitation of weak cement. Deeper stratigraphic intervals with smaller amounts of dispersed ash show no such effects, regardless of temperature or depth.

The clay mineral assemblages of Shikoku Basin show consistent temporal changes, particularly over the last 10 Myr, with gradual reductions of detrital smectite and gradual increases in illite and chlorite (moving up-section). At IODP Sites C0011 and C0012, percentages of smectite within bulk mudstones decrease by roughly 3 wt-% for every 1-million-year reduction in age. Causes of this trend are multi-faceted but probably include: (1) intensification of the Kuroshio Current after closure of the Isthmus of Panama (at about 3 Ma); (2) blockage of transport routes from the East China Sea by rifting of the Okinawa Trough and subaerial buildup of the Ryukyu Arc (peaking at about 7 Ma); and (3) progressive uplift and denudation of detrital source rocks in the Outer Zone of Japan, which gradually stripped away the near-surface products of anomalous near-trench magmatism and exposed deeper plutonic roots and surrounding meta-sedimentary strata of the Shimanto Belt. These systematic changes in clay mineral assemblages are important because they modulate friction properties, consolidation behavior, and fluid production from dehydration reactions as the strata move deeper into the subduction zone.

Keywords: Shikoku Basin, clay minerals, anomalous porosity, time-transgressive facies

Variation in Porosity of the Nankai Trough Incoming Sediments off the Kii Peninsula,
Southwest Japan

Variation in Porosity of the Nankai Trough Incoming Sediments off the Kii Peninsula,
Southwest Japan

*朴 進午¹

*Jin-Oh Park¹

1. 東京大学 大気海洋研究所 海洋底科学部門

1. Department of Ocean Floor Geoscience, Atmosphere and Ocean Research Institute, The University of Tokyo

The Nankai Trough is one of the best subduction-zone to study megathrust earthquake fault. Historic, great megathrust earthquakes with a recurrence interval of 100-200 yr have generated strong motion and large tsunamis along the Nankai Trough subduction zone. At the Nankai Trough margin, the Philippine Sea Plate (PSP) is being subducted beneath the Eurasian Plate to the northwest at a convergence rate ~4 cm/yr. The Shikoku Basin, the northern part of the PSP, is estimated to have opened between 25 and 15 Ma by backarc spreading of the Izu-Bonin arc. The >100-km-wide Nankai accretionary wedge, which has developed landward of the trench since the Miocene, mainly consists of offscraped and underplated materials from the trough-fill turbidites and the Shikoku Basin hemipelagic sediments. Particularly, physical properties of the incoming sediments to the Nankai Trough may be critical for seismogenic behavior of the megathrust fault. We have carried out core-log-seismic integration (CLSI) to estimate 3D acoustic impedance and porosity for the incoming sediments to the Nankai Trough off the Kii Peninsula, southwest Japan. For the CLSI, we used 3D seismic reflection, P-wave velocity, density, and porosity data obtained during IODP (Integrated Ocean Drilling Program) Expeditions 322 and 333. We computed acoustic impedance depth profiles for the IODP drilling sites from P-wave velocity and density data. We constructed seismic convolution models with the acoustic impedance profiles and a source wavelet which is extracted from the seismic data, adjusting the seismic models to observed seismic traces with inversion method. As a result, we have successfully obtained 3D acoustic impedance volume. With the 3D acoustic impedance volume and the porosity data at the IODP sites, we have performed multi-attribute transform that allows us to predict rock properties beyond the well location from seismic attributes calibrated with well-log data. The seismic attributes can be calculated internally, or provided as external attributes. The analysis was carried out in several stages: (1) to examine the log and seismic data at well locations to determine which set of attributes is appropriate; (2) to derive a relationship using multi-linear regression or Neural Networks; (3) to apply the derived relationship to a 3D SEG-Y volume to create a volume of the desired log property. In general, the 3D porosities show decrease with depth. We found a porosity anomaly zone with alteration of high and low porosities seaward of the trough axis. In this talk, we will show detailed 3D porosity of the incoming sediments, and present implications of the porosity anomaly zone for the megathrust fault behavior.

キーワード: Nankai Trough、3D porosity、Core-log-seismic integration

Keywords: Nankai Trough, 3D porosity, Core-log-seismic integration

Role of the subducting plate on the ETS-type phenomena around the mantle wedge corner

*Xiang Gao^{1,2}, Kelin Wang³, Jiangheng He³

1.Institute of Oceanology, Chinese Academy of Sciences, 2.Qingdao National Laboratory for Marine Geology, 3.Pacific Geoscience Centre, Geological Survey of Canada, Natural Resources Canada

Episodic tremor and slip (ETS) and related phenomena around the mantle wedge corner usually occur in young-slab subduction zones and are most abundant in Nankai, Cascadia, and Mexico. They are rare or absent in old-slab subduction zones such as Japan Trench. The relationship of these ETS-type phenomena with the subducting plate is a question of fundamental importance. In this work, we address this question through thermal modelling using heat flow data as model constraints. Our model employs the hypothesis that extremely high fluid pressure exists around the mantle wedge corner, consistent with relevant seismic and other observations. In young-slab subduction zones Nankai, Cascadia, and Mexico, the subduction fault has two frictional segments, with the first one being shallower than the mantle wedge corner because the viscous strength of the subduction faults is relatively small. Around the mantle wedge corner, the second frictional segment appears, but with abnormally low frictional strength due to the high fluid pressure and possibly the presence of hydrous minerals. We propose this is where the ETS occurs. Between these two frictional segments, the fault exhibits semi-frictional and/or viscous behaviour, which may facilitate long-term slow slip events and/or aseismic creep. For most old-slab subduction zones such as Japan Trench, there is only one frictional segment which extends to deeper than the mantle wedge corner because of the greater viscous strength of its megathrust than in young-slab subduction zones. We think the lack of a second frictional segment retards the development of ETS. However, two frictional segments are present in the old-slab Northern Hikurangi subduction zone because the greater frictional strength of its megathrust causes the termination of the frictional segment to occur shallower than the mantle wedge corner, explaining why ETS-type phenomena are observed here. The greater frictional strength is attributed to the roughness of the subducting seafloor. Our results indicate that both the age and roughness of the subducting plate play a key role on fault rheology which may control the occurrence of the ETS-type phenomena.

Keywords: subducting plate, episodic tremor and slip, high fluid pressure, heat flow

海洋リソスフェア内地震のb値の歪み速度依存性

Strain rate dependency of b-values of oceanic intraplate earthquakes

*篠島 僚平¹、伊藤 武男¹

*Ryohei Sasajima¹, Takeo Ito¹

1.名古屋大学大学院 環境学研究科

1.Graduate School of Environmental Studies, Nagoya University

Oceanic intraplate earthquakes (OCEQs), which are intraplate earthquakes occurred in the oceanic lithosphere (from oceanic ridge to before outer-rise), provide information regarding stress state in the oceanic lithosphere. We modeled stress evolution in the oceanic lithosphere and revealed that observed focal mechanism of OCEQs could be basically explained by thermal stress [Sasajima and Ito, 2015, SSJ fall meeting]. Also decreasing seismic moment release by OCEQs with age of the oceanic lithosphere could be well explained by age dependent thermal stress generation rate in young age oceanic lithosphere (<30Ma) quantitatively. However, seismic moment release by OCEQs in old age oceanic lithosphere (>45Ma) was only approximately 10% of expected one from model. This gap was significant even if we consider the uncertainties due to shortage of observation period. In this presentation, we report candidate of explanation of this gap and remarkable results. We revealed that b-values of OCEQs show clear age dependency. OCEQ b-values in young oceanic lithosphere (<15Ma) are around 1.0, those in middle age (15-30Ma) are around 1.4, and those in old age (>30Ma) are around 1.7, which is significantly higher than b-values in the world. If b-value is greater than 1.5, contribution of small earthquakes on total seismic moment release becomes significant. Thus, we suggest that missing seismic moment release in old age oceanic lithosphere is released by unobservable ($M_w < 5.0$) small magnitude earthquakes.

We also found that b-value of intraplate earthquakes in the Ninety East-Sumatra orogen, where oceanic lithosphere has higher strain rate than it of normal oceanic lithosphere, is 0.93, which is significantly smaller than OCEQ b-value (1.7) at the same age. Thus, we conclude that OCEQ b-value age dependency relates to the age dependent slow strain rate of the oceanic lithosphere and is not caused by age dependent thermal structure.

キーワード：海洋リソスフェア内地震、b値、海洋リソスフェア

Keywords: Oceanic intraplate earthquakes, b-value, Oceanic lithosphere

地震波干渉法を適用したエアガン-海底地震計探査記録に基づく日本海溝北西太平洋の海洋プレート反射構造

Seismic structure of northwest of the Pacific Plate by applying seismic interferometry to airgun-OBS data

*大友 周平¹、日野 亮太¹、東 龍介¹、藤江 剛²、小平 秀一²

*Shuhei Otomo¹, Ryota Hino¹, Ryosuke Azuma¹, Gou Fujie², Shuichi Kodaira²

1.東北大学大学院理学研究科地震・噴火予知研究観測センター、2.海洋研究開発機構

1.Research Center for Prediction of Earthquakes and Volcanic Eruptions, Graduate School of Science, Tohoku Univ., 2.Institute for Research on Earth Evolution (IFREE), Japan Agency for Marine-Earth Science and Technology (JAMSTEC)

海洋性地殻の地震波速度構造は、比較的均質性が高いと考えられるが、海洋性地殻下部から最上部マントルに複雑な構造をもった反射面が認められることがある。こうした地殻深部・マントルにみられる不均質構造は、海洋プレートの形成・発達の過程を記録したものと考えられ、たとえば、千島海溝沖の北西太平洋の最上部マントルに認められる反射構造は、プレート運動に伴って形成されたリーデル剪断と解釈されている。本研究では、日本海溝沖におけるエアガン-海底地震計(OBS)構造探査で取得したエアガン波形記録を解析し、地殻・マントル内の地震波反射構造からこの領域の海洋プレートの形成・発達過程を解明することをめざす。従来、大震災距離での観測に特徴付けられるOBSデータの解析手法は、エアガン信号の走時情報から速度場を推定する走時解析が一般的であったが、近年では、OBSデータに反射法的な解析を施すことで、深部構造のより高分解能なイメージングが可能となってきた。OBSデータに地震波干渉法を適用すると、OBS観測点間隔より稠密な仮想受信点配置の地震波波形を得ることができるため、反射法的解析の結果を改善できることが期待できる。そこで、本研究ではOBSデータに地震波干渉法を適用して海面発震-海面受信の稠密な合成波形データを作成し、これに反射法的処理を施して海洋プレート内反射構造のイメージングを行った。今回使用した探査データは、2011年に北西太平洋において実施されたエアガン-OBS探査記録である。本測線周辺の海底下構造は、水平成層に近い構造を持っている。OBSは6 km間隔で23台設置されており、エアガン発振は0.2 km間隔で1199回行った。地震波干渉法を適用することで、エアガン発振間隔の0.2 km間隔で仮想的に受信点を配置したときの、稠密な合成波形が得られることが期待される。解析では、まず地震波干渉法を適用し、合成波形を作成した。地震波干渉法では、1つの受信点で観測した、異なる2つの震源からの波形の相互相関をとることで、一方を震源、もう一方を受振点としたときの波形記録を合成できる。例えば、海底反射波と多重反射波の相互相関を取ることで、モホ面反射波の合成波形を得ることができる。次に、震源と受信点の midpoint が一致するトレースを並べ、CMPアンサンブルを作成して重合を行った。RMS速度を1.5 km/sとしてNMO補正を行って重合を行った反射断面には、垂直往復走時7.3秒付近に連続性のあるフェーズがみとめられ、水深5,500 m程度にある海底が、地震波干渉法と反射法的処理によりイメージングされていることが確かめられた。海底下については、重合速度を1.75 km/sから3.75 km/sの範囲で変化させて、重合記録を検討したところ、モホ面に対応すると考えられる連続性の良い反射面が、3.50 km/sでNMO補正を行ったときにイメージされた。現状では、モホ面反射波のS/N比は低いことから、地震波干渉法適用前の元波形に適切なフィルタを適用するなどすることにより、反射断面の品位向上を図り、地殻・最上部マントル内部の詳細な構造イメージングを進める予定である。

北西太平洋海溝海側斜面に発達するプレート屈曲起源の地形的特徴

Bending-related topographic structures of the subducting Pacific plate in the Northwestern Pacific Ocean

*中西 正男¹*Masao Nakanishi¹

1. 千葉大学大学院理学研究科 地球生命圏科学専攻

1. Geosystem and Biosystem Sciences Division Graduate School of Science Chiba University

Prolonged topographic structures associated with bending of the subducting Pacific plate along the trenches of the western margin were investigated using multibeam bathymetric data, accompanying reidentification of magnetic anomaly lineations on the Pacific plate to reveal controlling factors for strikes of bending-related topographic structures. The newly compiled bathymetric map demonstrates that most of bending-related topographic structures exist in the oceanward trench slopes within 100 km from trench axes. The bending-related topographic structures are developed parallel to the trench axis or inherited seafloor spreading fabric. Seafloor spreading fabric were reactivated instead of forming new trench parallel structures in the area where it strikes at an angle less than 30 to a trench axis. The topographic expression of bending-related structures is classified into two types according to whether new structures develop parallel to the trench axis or inherited seafloor spreading fabric reactivates. That of reactivated structures is characterized by half-graben or narrow ridge topography. That of newly formed structures is characterized by a horst and graben topography.

キーワード：プレート屈曲起源の地形、海底拡大構造、太平洋プレート

Keywords: bending-related topographic structures, seafloor spreading fabric, Pacific Plate

Marine magnetic anomaly and magnetization of the subducting Pacific Plate seaward of the Japan Trench

Marine magnetic anomaly and magnetization of the subducting Pacific Plate seaward of the Japan Trench

*富士原 敏也¹、木戸 ゆかり²

*Toshiya Fujiwara¹, Yukari Kido²

1. 海洋研究開発機構地震津波海域観測研究開発センター、2. 海洋研究開発機構地球深部探査センター

1. Research and Development Center for Earthquake and Tsunami, Japan Agency for Marine-Earth Science and Technology, 2. Center for Deep Earth Exploration, Japan Agency for Marine-Earth Science and Technology

We studied marine magnetic anomaly and magnetization of the subducting Pacific Plate seaward of the Japan Trench. Magnetization is one of the indicators to understand physical and chemical conditions of the oceanic plate, and also tectonic processes the plate has experienced. The Pacific Plate on the seaward slope of the Japan Trench is characterized by a series of parallel magnetic anomalies (Japanese Lineation) during M14-M7 (140-132 Ma). Half spreading rates when the plate was formed were estimated to be 4-6 cm/yr which is categorized as fast-spreading. It was considered that the plate was situated at low latitudes in the southern hemisphere, and was drifted to the present place. We made the magnetic anomaly dataset by compiling our JAMSTEC data, as well as data from Japan Oceanographic Data Center (JODC), National Geophysical Data Center (NGDC), and Geological Survey of Japan (GSJ). The magnetic anomalies are well lineated and have high-amplitudes. Meanwhile, the Pacific Plate is being subducted beneath the Tohoku Arc at the Japan Trench. The amplitudes of the anomalies gradually decrease toward the land from the trench axis. The decrease of amplitudes is mainly caused by increasing depth of the oceanic plate associated with subduction. In order to correct the effects caused by increasing depth of the subducting plate, we calculated magnetization from the magnetic anomaly. We used a three-dimensional inversion method for the calculation of magnetization. Upper surface of the magnetic layer was assumed to be the surface of oceanic crust. In the seaward slope, the surface was defined by subtracting 500 m as the thickness of sedimentary layer from the bathymetry. The surface of the subducting oceanic crust in the landward of the trench was determined using seismic refraction and reflection profiles. Densely distributed profiles of seismic survey in the study area enabled us to constrain the depth of the plate. The declination and inclination of magnetization were set to be several directions around -30° to $+30^\circ$, -20° to $+20^\circ$, respectively, in reference to ocean drilling rock magnetic measurements, skewness of the magnetic anomaly, and seamount's magnetization. On the seaward slope of the Japan Trench, fracture zones, which are originated from transform faults and are large offsets in the oceanic plate, are not identified. However, the existence of non-transform discontinuities (NTD) is probable. The NTDs are important to better understand the physical conditions of the oceanic plate, because they may act as some kind of weak zones. However, the NTDs are difficult to be identified, because sedimentary layers cover the old seafloor and conceal the abyssal hill fabrics. In this situation, the magnetization is useful for identifying the NTDs. Magnetic lineations located in 37° - 39° N adjacent to the trench show highly oblique and largely discontinuous. This disorganized structure was implicated in the past ridge propagation. In other places, higher magnetization or small dislocation linearly aligned sub-perpendicular to the strike of magnetic lineations could suggest small offset NTDs. Low magnetization appears on the seaward slope near the trench, and the magnetization intensity gradually decreased as the plate subduction proceeded. On the seaward slope, horst-graben structure has been developed and large steps have

grown associated with plate bending and normal faulting. Tectonic phenomena such as formation of the horst-graben structure and subsequent plate subduction would cause destruction and disorganization of the magnetic layer by faulting, which implies mechanical demagnetization. Possible cause of chemical demagnetization in the oceanic crust is low-temperature oxidation due to hydrothermal circulation along the faults. Serpentinization of the uppermost mantle may modify the magnetization.

キーワード : Magnetic Anomaly、Magnetization、Pacific Plate、Japan Trench、Horst-Graben、Subduction
Keywords: Magnetic Anomaly, Magnetization, Pacific Plate, Japan Trench, Horst-Graben, Subduction

人工電流源電磁探査による、沈み込み直前の太平洋プレート地殻比抵抗構造調査

Crustal resistivity Structure of Pacific Plate just before Subduction using Marine Controlled-Source Electromagnetic Surveys

*後藤 忠徳¹、笠谷 貴史²、川田 佳史³、市原 寛⁴、山野 誠⁵

*Tada-nori Goto¹, Takafumi Kasaya², Yoshifumi Kawada³, Hiroshi Ichihara⁴, Makoto Yamano⁵

1.京都大学大学院工学研究科、2.海洋研究開発機構、3.東北大学災害科学国際研究所、4.神戸大学大学院海事科学研究科、5.東京大学地震研究所

1.Graduate School of Engineering, Kyoto University, 2.Japan Agency for Marine-Earth Science and Technology, 3.International Research Institute of Disaster Science, Tohoku University, 4.Graduate School of Maritime Sciences, Kobe University, 5.Earthquake Research Institute, University of Tokyo

In the Japan Trench subduction zone, an old oceanic plate with an age of over 100 m.y., the Pacific plate, is subducting beneath the northeast Japan arc. However, recent heat flow surveys have revealed that the Pacific plate just before subduction may not be uniformly cold contrary to its old age. Localized high heat flow anomalies suggest that thermal structure of the Pacific plate in this area is not a typical one for old oceanic lithosphere, at least at shallow depths. To evaluate the factors controlling the temperature structure (e.g., fractures, fluid etc., together with heat sources), we conducted dense heat flow surveys and electromagnetic soundings on the seaward slope of the Japan Trench. In this presentation, we focus on the preliminary result obtained by the electromagnetic soundings.

The offshore experiment was done on July, 2014 at the KY14-10 cruise (R/V Kaiyo, JAMSTEC). The water depth in the target area (on the Pacific plate, far off Tohoku region, Japan) is about 5300-5600 m. We used a newly-developed controlled-source electromagnetic (CSEM) survey system for this research. It can be applied to exploration of seafloor resistivity, originally designed for AUV but available with a deep-tow system (DT). The power for transmitter is supplied from the DT. Three pressure cases (transmitter, controller and power converter) are mounted on the DT frame, and a long cable with length of 350m is towed behind the DT. Two source electrodes attached at the both end of cable are used for transmitting the artificial electric current into the seawater. The pulse width for artificial current was 2 seconds, and the amplitude of current was approximately 20-30 A. Two CSEM profiles were conducted in this cruise. The altitude of DT (and the source dipole) was about 50-100m with towing speed of about 1 knot or less.

The preliminary results indicate the low resistivity of shallow layers of the oceanic crust of Pacific plate. The amplitude of received electric field recorded by an OBEM was about 0.04 nV/(Am²) when the source-receiver separation was about 1km. The received signal can be clearly recognized within the source-receiver separations of about 4km. All of the amplitude of electric field is smaller than the predicted ones from a numerical simulation, with assuming the uniform sub-seafloor structure with 1 Ohm-m (and a sea water layer of 0.3 Ohm-m). This implies that the shallow part of oceanic crust (possibly less than 1km) have very low resistivity than 1 Ohm-m. The anomalous feature may be related to the buried fractures in the top of oceanic crust, and could be related to the fluid circulations, a candidate of factors to explain the high heat flow.

キーワード：比抵抗、人工電流源電磁探査、太平洋プレート

Keywords: resistivity, CSEM, Pacific Plate

日本海溝のホルスト・グラーベン構造を被覆する堆積物表層の岩相と物性：新青丸KS-15-3航海コア
解析速報

Lithology and physical property of sediments covering horst-graben structures of the Japan
Trench: Preliminary results of KS-15-3 sediment core analysis

*山口 飛鳥¹、新井 和乃²、池原 研³、金松 敏也²、福地 里菜¹、中村 恭之²、宇佐見 和子³、奥津 なつみ¹、清
家 弘治¹、芦 寿一郎¹

*Asuka Yamaguchi¹, Kazuno Arai², Ken Ikehara³, Toshiya Kanamatsu², Rina Fukuchi¹, Yasuyuki Nakamura²
, Kazuko Usami³, Natsumi Okutsu¹, Koji Seike¹, Juichiro Ashi¹

1.東京大学大気海洋研究所、2.海洋研究開発機構地震津波海域観測研究開発センター、3.産業技術総合研究所
地質情報研究部門

1.Atmosphere and Ocean Research Institute, The University of Tokyo, 2.R&D Center for Earthquake
and Tsunami, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 3.Research Institute
of Geology and Geoinformation, National Institute of Advanced Industrial Science and Technology
(AIST)

Large coseismic slip along the shallow portion of the plate boundary fault beneath the Japan Trench
caused the catastrophic tsunami of the 2011 Tohoku Earthquake (e.g. Fujiwara et al., 2011; Ito et
al., 2011; Kodaira et al., 2012). Researches on core samples taken by D/V Chikyu (IODP Expedition
343, JFAST) revealed that the plate boundary fault at the IODP Site C0019 of the Japan Trench is
composed of pelagic clay layer which contains large fraction of smectite, and low coseismic
friction of smectite would contributed to the coseismic slip propagation along the shallow portion
of the plate boundary fault (Ujiie et al., 2013; Kameda et al., 2015; Moore et al., 2015). However,
high-resolution seismic profiles indicate heterogeneous distribution of pelagic clay layer, and
there are several portions where the thicknesses of the incoming sediments are very thin (less than
50 m). To reconcile the nature of the "thin" incoming sediments, we sampled seven piston cores from
horst-graben structures of the Japan Trench during the KS-15-3 cruise (R/V Shinsei Maru, May 3-19,
2015).

Cores were taken from four different settings: PC01 from graben, PC02 and PC07 from graben edge,
PC03 and PC04 from horst, PC05 and PC06 from seaward trench slope, respectively. We present
preliminary results of core analysis including visual core descriptions, X-ray CT images and
successive density and magnetic susceptibility values measured by multi-sensor core logger (MSCL),
and discuss sedimentation process of horst-graben structures that causes the diversity of incoming
sediments.

摩擦発熱から推定する日本海溝東側斜面における表層堆積物の力学的性質

Mechanical properties of surface sediments estimated from frictional heating on east slopes in the Japan Trench

*吉本 佳太²、山野 誠⁴、笠谷 貴史³、川村 喜一郎¹

*Keita Yoshimoto², Makoto Yamano⁴, Takafumi Kasaya³, Kiichiro Kawamura¹

1.山口大学・JAMSTEC、2.山口大学、3.海洋研究開発機構、4.東京大学地震研究所

1.Yamaguchi University/JAMSTEC, 2.Yamaguchi University, 3.JAMSTEC, 4.ERI, University of Tokyo

1. Introduction

Marine heat flow measurements have been conducted on the northwest Pacific plate off the Japan Trench since the 1960s, and a large amount of data was accumulated by recent intensive surveys (Yamano et al, 2014). Generally, the heat flow is obtained in the following way. 1) a heat flow probe (hereafter probe) of several meters in length attached with several temperature sensors is penetrated into sediments in order to measure the geothermal gradient, 2) thermal conductivity of the sediments collected at or around the penetration point is measured, 3) the heat flow at the point is calculated from the geothermal gradient and the thermal conductivity. The geothermal gradient can be calculated from the temperature data obtained with the sensors attached along the probe. In general, right after the probe penetration into the seafloor, the sensor temperature increases rapidly by about 1 K due to frictional heating and then gradually approaches the sediment temperature at the sensor depth. But, in the area off the Japan Trench, we found that there are horizons with extraordinarily large frictional heating, where the temperature rise by frictional heating exceed 5 K (hereafter anomalous heating horizon). The anomalous heating horizons might correspond to some hard and/or consolidated layers. The purpose of this study is to understand the relationship between the anomalous heating horizons and the mechanical properties of sediments.

2. Survey area and the research cruises

Heat flow was measured at 69 points using a 3 m long probe during the cruises YK14-21, KS-14-17 and KS-15-16. Sediment samples were obtained at two sites in the survey area (KR09-16 HFPC01 and KR10-12 HFPC01) using a piston corer with temperature sensors attached along the core barrel. In this study, we used four measurement devices attached to or in combination with the probe in order to obtain information on the penetration process; 1) Temperature sensors (sampling rate: 10 sec, except for eight sites with a rate of 1 sec). 2) Tiltmeter (30 sec). 3) Wire tension meter (1/10 sec). 4) Accelerometer (1/100 sec). Because we cannot directly observe the penetration process, we should indirectly understand the penetration process (e.g., penetration start time and end time) using above four data.

3. Result

The core sediments, KR09-16 HFPC01 (274.5 cm in length) and KR10-12 HFPC01 (181.0 cm in length), are mainly composed of silty clay with diatoms. Tephra layers were observed at 120 - 145 cm and 155 - 180 cm in KR09-16 HFPC01 and at 100 - 102 cm and 140 - 148 cm in KR10-12 HFPC01.

Anomalous heating horizons were detected in a depth range from 150 cm below seafloor (hereafter cm-bsf) to 300 cm-bsf. The frictional heat generated at the horizon tends to be higher at eastern and northern parts in the survey area.

The results of this study indicate that the anomalous heating horizon continuously extends sub-parallel to the seafloor, and there is a regional variation in the amount of heat generation. We suppose that the anomalous heating horizons result from a hard layer with lateral variation of the thickness, corresponding to the tephra layers observed in the core samples. However, the subbottom depths of the anomalous heating horizons, ranging from 150 to 300 cm-bsf, are several

tens of centimeters deeper than those of the tephra layers in the samples. We intend to make further investigation on the problem of this depth difference.

キーワード：太平洋プレート、熱流量、テフラ層

Keywords: Pacific plate, Heat flow, Tephra layer

南海トラフ室戸沖から熊野沖にかけて観測された熱流量変化のモデル化

Modeling the observed heat flow variation along the Nankai Trough between the Muroto and Kumano transects

*川田 佳史¹、山野 誠²*Yoshifumi Kawada¹, Makoto Yamano²

1.東北大学災害科学国際研究所、2.東京大学地震研究所

1.International Research Institute of Disaster Science, Tohoku University, 2.Earthquake Research Institute, University of Tokyo

Gradual decrease in heat flow with a spatial scale of several tens of kilometers is observed between the Muroto and Kumano transects of the Nankai Trough, which are located 150 km apart [Yamano et al., 2013, JpGU]. The average heat flow on the trough floor is twice as that of model predicted value around the Muroto transect, whereas that is comparable to model predicted value around the Kumano transect. In this study, we construct a series of numerical models including hydrothermal heat transport within the oceanic crust in order to explain the observed gradually decreasing heat flow.

The first model considers a three-dimensional domain using enhanced thermal conduction within the aquifer [Spinelli and Wang, 2008, Geology], which mimics hydrothermal heat transport by a simplest manner. With a gradual decrease of basal heat flow along the trench axis corresponding to the age difference between the Muroto and Kumano transects, the resulting heat flow gradually varies along the trench axis but is too gradual to account for the observation. With a sharp contrast of the aquifer permeability, i.e., high permeability at the Muroto transect and low permeability at the Kumano transect, hydrothermal heat transport at the Muroto transect does not affect the Kumano transect, and this is also inconsistent with the observation. We expect that the difference in the aquifer permeability as well as the difference in the basal heat input is required to account for the observed heat flow contrast between the Muroto and Kumano transects.

The second model considers the effect of trench-parallel heat and fluid transport, using a two-dimensional computational domain parallel to the trench axis covering the area between the trough floor of the Muroto and Kumano transects. Referring to the result of the first model, in which hydrothermal heat transport occurs along the trench axis, we include an internal heating in a half of the aquifer corresponding to the Muroto transect. In cases of no or weak hydrothermal circulation, for example the permeability is of the order of 10⁻¹² m², square-sized convection cells are formed within a 500 m thick aquifer, which transport heat vertically through the aquifer and result in a sinusoidal heat flow pattern. The increase in the aquifer permeability results in an elongated convection cell across the heating and non-heating areas. Heat flow above the elongated cell gradually decreases with the spatial scale of the cell. The spatial scale is several tens of kilometers using the permeability of the order of 10⁻¹⁰ m². This high permeability is consistent with the condition for along-aquifer hydrothermal heat transport to occur.

We construct two end-member models that can qualitatively explain the observed heat flow: (1) three-dimensional heat upwelling involving in the Muroto and Kumano transects and (2) two-dimensional heat upwelling along the Muroto transect and along-trench heat transport of the upwelled heat below the trough floor from the Muroto to Kumano areas. The first model assumes that heat transport pervasively occurs throughout the aquifer between the two transects. The second model, in contrast, assumes that the heat is mined below the Muroto transect, and a part of which is transported along-trench direction below the trough floor. The area involved and the intensity of heat upwelling are different between these models. More heat flow observations, especially on

the landward crust between the Muroto and Kumano transects, are required to distinguish these two end-member models.

キーワード：地殻熱流量、数値モデリング、間隙水循環、南海トラフ、室戸沖、熊野沖

Keywords: heat flow, numerical modeling, pore fluid flow, Nankai Trough, Muroto transect, Kumano transect

Structural variation of the incoming Philippine Sea plate along the Nankai Trough off Shikoku

Structural variation of the incoming Philippine Sea plate along the Nankai Trough off Shikoku

*仲西 理子¹、山本 揚二郎¹、山下 幹也¹、藤江 剛¹、三浦 誠一¹、小平 秀一¹、金田 義行^{1,2}

*Ayako Nakanishi¹, Yojiro Yamamoto¹, Mikiya Yamashita¹, Gou Fujie¹, Seiichi Miura¹, Shuichi Kodaira¹, Yoshiyuki Kaneda^{1,2}

1.海洋研究開発機構、2.名古屋大学減災連携研究センター

1.Japan Agency for Marine-Earth Science and Technology, 2.Disaster Mitigation Research Center, NAGOYA University

The next large-thrust earthquake along the Nankai Trough, southwest Japan is concerned to occur within this century. Nonvolcanic deep low-frequency tremors and very low-frequency (VLF) earthquakes considered as one of indicators of the future large-thrust earthquakes are observed around the down-dip limit of the coseismic rupture zone of the last Tonankai and Nankai earthquakes [Obara, 2002]. However the activity of the nonvolcanic deep low-frequency tremors and VLF earthquakes is not homogeneous, and the belt-like tremor zone is divided into several segments bounded by gaps [Obara, 2010]. One of the causes of these low-frequency seismic phenomena is considered to be fluid generated by dehydration processes from the subducting slab. To investigate the relation between the formation process of the Philippine Sea plate and the occurrence of the low-frequency seismic phenomena as well as large-thrust earthquakes, it is important to investigate structural variation in the incoming Philippine Sea plate, including its fluid content.

In 2014, we conducted the seismic refraction and reflection survey in the northern margin of the Shikoku Basin, where the Philippine Sea plate is subducting beneath the Eurasia plate at the Nankai Trough. We conducted a 360km long seismic profile about 50-60km seaward of the deformation front along the Nankai Trough. 35 OBSs were deployed along the profile with the interval of 10km. A tuned airgun array shot with a total volume of 7800 cu. in. every 200m. Moreover, high-resolution multichannel seismic reflection survey is conducted along the same profile by using a 192-channel, 1.2km-long hydrophone streamer and an airgun array of 380 cu. in. volume shot by every 37.5m.

In the time-migrated reflection section, variation in the configuration of the basement reflection can be recognized as the structural boundary off the cape Muroto. The result of first-arrival tomography based on the wide-angle OBS data shows dramatic change in P-wave velocity just beneath the basement corresponding with the structural boundary observed along the reflection section as mentioned above. The structural boundary may be related to the boundary of the plate age proposed by magnetic lineation [Okino, 2015]. From the northeast of the structural boundary, it seems that the undulation cycle of the basement configuration is remarkably similar to the cycle pattern of frequency distribution of tremor along the tremor belt-like zone beneath the Shikoku Island shown by Obara [2010]. Similar undulation cycle of the basement configuration is also recognized in the reflection section previously acquired along trough axis just landward of the seismic profile of this study. Such incoming Philippine Sea plate with the dramatic structural variation is subducting beneath the southwestern Japan, and may have the possibility of any effects on the occurrence of the low-frequency seismic phenomena.

We will show the structural variation in the V_p/V_s ratio that is a sensitive to the presence of fluid by using PS converted wave observed by OBSs.

This study is part of 'Research project for compound disaster mitigation on the great, earthquakes and tsunamis around the Nankai Trough region' funded by MEXT, Japan.

Subduction of oceanic plate irregularities in Mexico and Japan and the influence on large megathrust earthquakes

*Marina Manea¹, Vlad Constantin Manea^{1,2}, Shoichi Yoshioka²

1.Computational Geodynamics Laboratory, Geosciences Center, National Autonomous University of Mexico, 2.Research Center for Urban Safety and Security, Kobe University, Japan

It is known that oceanic plates morphology is not a simple one, but rather complicated by a series of irregularities as seamounts, fracture zones and mid-ocean ridges. These features present on the oceanic floor form part of the fabric of oceanic crust, and once formed they move together with the oceanic plates until eventually enter a subduction zone. Some improvements have been done in the last decades in understanding the influence of seamount subduction on the forearc morphology and also on occurrence of large subduction earthquakes. Although the Cocos plate offshore Mexico present a series of multiple small height seamount chains that enter into subduction as well as large fracture zones (i.e. Tehuantepec fracture zone), the understanding of the influence of these oceanic features that might play in geohazards (i.e. the generation of large earthquakes) in Mexico is not yet well studied. Previous studies along the Mexican trench reveal that only a little portion of the oceanic sedimentary blanket is accreted and this margin seems to be marked by erosional processes. In the particular case of seamounts, it is not well understood for example whether the Cocos plate seamount chains are accreted to the forearc or carried down into the subduction zone, and how the forearc morphology is affected. Whereas offshore Mexico the Cocos plate seafloor is littered with small but numerous seamounts and seamount chains, the Pacific plate offshore Japan is marked by relatively few but large irregularities (i.e. seamounts). However, both subduction zones are very active and prone to produce large and devastating earthquakes. In this study we use for the ocean bottom topography SRTM30+ datasets which provide global topography and bathymetry at 30" per-cell resolution (~ 1 km). In order to better identify shallow features as seamounts, we use a median-depth grid (using a median filtering technique) which is applied to the original bathymetric data. Here we present some preliminary results where we investigate in a comparative manner the shape and structure of medium-size to large seamounts and fracture zones in the vicinity of the trench in Mexico and Japan, and examine the possible links that might exist between these features and the rupture history of large subduction zone earthquakes.

Keywords: seamounts, fracture zones, Japan subduction zone, Mexican subduction zone

プレートテクトニクス沈込スラブへの拡張

Extension of Plate Tectonics to subducting Slab

*新妻 信明¹*Nobuaki Niitsuma¹

1. 静岡大学理学部地球科学教室、仙台

1. Institute of Geosciences, Shizuoka University, Sendai

半世紀前に確立したプレートテクトニクスは、中央海嶺における海洋プレートの拡大を定量的に記述することに成功した。しかし、拡大した海洋プレートの沈込については定性的な推定の域（「海洋底拡大説」の域）を脱するに到っていない。変形しなければ海洋プレートは沈込めないで、プレートテクトニクスの中心教義「変形しないプレート」を放棄しなければならない。海溝周辺の活発な地震活動は、海溝に沿って海洋プレートが沈込していることに対応している。東日本大震災以後の地震活動は、太平洋プレートが海溝に沿って同心円状屈曲して沈込み、平面化して和達の深発地震面に連続していることを示している。

プレートテクトニクスを海洋プレートに連続するスラブへ拡張するために、以下の仮定に限定して中心教義「変形しないプレート」を解除し、定量化する。

1) 海溝軸は屈曲しており、日本海溝については北から襟裳・最上・鹿島と名付けた小円に沿う円弧をなしている。

2) オイラー回転する海洋プレートに連続するスラブ上の点は地心から見て同一オイラー緯線に沿って移動する。

3) オイラー回転によるスラブ上面に沿うの移動距離は海洋プレートの移動距離に等しく、地心から見たスラブ上の点の移動速度はスラブ傾斜に応じて海洋プレート上の点よりも遅い。

4) スラブ上面深度は海溝軸小円心からの距離によって決定される。小円心はスラブ上面深度断面の回転対称軸になる。

5) 海洋プレートは海溝に沿って同心円状屈曲し、平面化角 A_p に達すると平面化して深発地震面に接続する（付図）。各小円についての同心円状屈曲半径や平面化角などの係数は地震活動に基づき決定する。

これらの仮定のもとに、日本海溝に沿って沈込む太平洋スラブの運動を約10km間隔で0.125my毎に算出し、気象庁の初動発震機構解とCMT発震機構解を比較解析した。気象庁の地震計網から外れている日本海溝域の震源を海底地震計によって決定された震源（Shinohara et al., 2011, 2012; Obana et al., 2011, 2012, 2013）と比較したところ、気象庁のCMT解の初動震源深度が深目に出ているが、震央分布に問題ないことが確認された。

深海底面上の点と海底面下5kmの点との距離は、海洋プレートが移動しても5kmと一定であるが、海溝に沿って同心円状屈曲すると屈曲半径の小さい深度5kmの点が先行し、距離が増大する。5%の5.25kmに達する位置は日本海溝側の海岸線に沿っており、屈曲スラブの平面化に伴う地震活動と対応している。スラブが日本列島下を通過して日本海側に到るとその距離は10%以上に増大し、5.5km以上になる。スラブ表層5kmでもこれだけ大きな変形をもたらす沈込は、日本列島下のマントルへ更に大きな影響を与え、日本列島の大地形形成に関与していることを示唆している。

海洋プレート運動方向のオイラー緯線に直交するオイラー経線に沿って並ぶ点の間の併進距離は、海洋プレート上では一定であるが、スラブ沈込に伴うスラブ上面深度差によって変動する。小円心が島弧側に位置する場合には併進距離が増大、海洋側に位置する場合には減少する。変動の細部は小円心の位置とプレート運動方位によって支配され、併進距離変動は最大±1%に達する。小円心が島弧側の最上小円区で引張応力による正断層型発震機構解が優勢で、海洋側の襟裳小円区・鹿島小円区で圧縮応力による逆断層型発震機構解が優勢であることと良く対応している。震源分布も併進距離変動と対応している。

深度200km以深CMT発震機構解の地心三次元座標系における最小自乗法によって算出された襟裳・最上・鹿島小円区からの平面（Vlad面）と、日本海溝に沿うスラブ上面との交線を算出した。この交線に沿って平面化による逆断層型震源が配列することと、その日本海側でCMT震源数が急減することは、海溝に沿うスラブの同心円状屈曲とVlad面への平面化が進行していることを示している。海溝から遠くの深い位置で交わる最上小円区から沈込だVlad面にはCMT震源が分布せず、海溝近くの浅い位置で交わる襟裳・鹿島小円区から沈込だVlad面に

CMT震源が分布していることは、深発地震発生過程について示唆を与える。

キーワード：プレートテクトニクス、沈込スラブ、オイラー回転、同心円状屈曲、平面化、三次元距離変化
 Keywords: Plate Tectonics, subducting slab, Euler rotation, concentric bending, unbending, changes in 3d distance

