

## モンゴル・アルタイ地域に認められる異なる変成履歴

Contrasting pressure-temperature records from the Altai Range, Mongolia; constraints from multiple growth of garnet, aluminosilicates and monazite

\*中野 伸彦<sup>1</sup>、小山内 康人<sup>1</sup>、大和田 正明<sup>2</sup>、Satish-Kumar Madhusoodhan<sup>3</sup>、足立 達朗<sup>4,1</sup>、Jargalan Sereenen<sup>5</sup>、吉本 紋<sup>1</sup>、Syeryekhan Kundyž<sup>5</sup>、Boldbaatar Chimedtseie<sup>6</sup>、Puntsagdamba Nomintsetseg<sup>7</sup>、Boldbaatar Dolzodmaa<sup>8</sup>

\*Nobuhiko Nakano<sup>1</sup>、Yasuhito Osanai<sup>1</sup>、Masaaki Owada<sup>2</sup>、Madhusoodhan Satish-Kumar<sup>3</sup>、Tatsuro Adachi<sup>4,1</sup>、Sereenen Jargalan<sup>5</sup>、Aya Yoshimoto<sup>1</sup>、Kundyž Syeryekhan<sup>5</sup>、Chimedtseie Boldbaatar<sup>6</sup>、Nomintsetseg Puntsagdamba<sup>7</sup>、Dolzodmaa Boldbaatar<sup>8</sup>

1.九州大学大学院比較社会文化研究院地球変動講座、2.山口大学大学院理工学研究科、3.新潟大学理学部、4.九州大学アジア埋蔵文化財研究センター、5.Department of Mineral Exploration, Mongolian University of Science and Technology、6.Mongolian Exploration Partners, LLC、7.九州大学大学院比較社会文化学府、8.九州大学大学院地球社会統合科学府

1.Division of Earth Sciences, Faculty of Social and Cultural Studies, Kyushu University, 2.Department of Earth Sciences, Yamaguchi University, 3.Department of Geology, Niigata University, 4.Kyushu University Advanced Asian Archaeological Research Center, 5.Department of Mineral Exploration, Mongolian University of Science and Technology, 6.Mongolian Exploration Partners, LLC, 7.Graduate School of Social and Cultural Studies, Kyushu University, 8.Graduate School of Integrated Sciences for Global Society, Kyushu University

Pelitic gneisses from the western Mongolian Altai Range of the Central Asian Orogenic Belt exhibit multistage aluminosilicate formation and various chemical zoning patterns in garnet. The studied pelitic rocks are divided into four types; garnet-kyanite-biotite gneiss, garnet-cordierite-biotite gneiss, garnet-staurolite-sillimanite-biotite gneiss, and garnet-staurolite-kyanite-biotite gneiss. The former two gneisses contain kyanite in the matrix and sillimanite inclusions in garnet. The Ca concentrations in garnet from the garnet-kyanite-biotite gneiss increase and those from garnet-cordierite-biotite gneiss decrease from centre to inner rim in the core. In garnet-staurolite-sillimanite-biotite gneiss, and garnet-staurolite-kyanite-biotite gneiss, sillimanite or kyanite occurs in the matrix, respectively, and both have kyanite inclusions in garnet. The garnet has homogeneous high-Ca core part, and the mantles are characterized by low-Ca. Monazite U-Th-Pb dating for the studied samples shows the bimodal ages; c. 360 Ma and c. 260 Ma. Combining the microstructural information, thermodynamic calculations, and geochronology suggests all rock types experienced compression at c. 360 Ma, but this compression occurred at different crustal levels. The garnet-kyanite-biotite gneiss and garnet core in garnet-cordierite-biotite gneiss represent compression at low-pressure conditions (~5.2 to 7.2 kbar) under moderate-temperature conditions (~620-660 C). In contrast, the garnet cores in garnet-staurolite-sillimanite-biotite and garnet-staurolite-kyanite-biotite gneisses were formed during compression at higher pressure conditions (~7.0 to 8.9 kbar at ~600-640 C). It is still obscure why the different thermal gradients existed during the compression but it is likely due to presence of ridge subduction in the Altai Range reported by several workers. The subducting ridge could supply heat to the accretionary wedge and produced new geotherm. The difference in thermal gradients observed in this study might be due to variations in thermal regimes from the subducting ridge, which has an important role in developing the variations in prograde pressure-temperature paths during burial of the accretionary wedge. Garnet mantles and the matrix mineral assemblages in three rock types except garnet-kyanite-biotite gneiss were recrystallized during the c. 260 Ma event characterized by amphibolite-facies metamorphism with a metamorphic thermal gradient of ~25

C/km, which might be caused by collision-related granitic activity and re-equilibrium at middle crustal depths. This study also suggests monazite U-Th-Pb dating combined with the occurrence, texture, and chemistry of associated mineral phases in amphibolite-facies rocks will allow us to recognize multiple events and the equilibrium phases during each event.

キーワード：ザクロ石、アルミノ珪酸塩鉱物、モナズ石、変成履歴、アルタイ、モンゴル

Keywords: garnet, aluminosilicate, monazite, pressure-temperature path, Altai, Mongolia

## スリランカ超高温変成岩の岩石学・年代学

Petrological and chronological constraints of UHT granulites from Highland Complex, Sri Lanka: implications on Gondwana correlation

\*小山内 康人<sup>1</sup>、Krishnan Sajeev<sup>2</sup>、中野 伸彦<sup>1</sup>、北野 一平<sup>3</sup>、Kehelpannala Wilbert<sup>4</sup>、加藤 涼介<sup>3</sup>、足立 達郎<sup>1</sup>、Malaviarachchi Sanjeewa<sup>5</sup>

\*Yasuhito Osanai<sup>1</sup>、Sajeev Krishnan<sup>2</sup>、Nobuhiko Nakano<sup>1</sup>、Ipppei Kitano<sup>3</sup>、Wilbert Kehelpannala<sup>4</sup>、Ryosuke Kato<sup>3</sup>、Tatsuro Adachi<sup>1</sup>、Sanjeewa Malaviarachchi<sup>5</sup>

1.九州大学大学院比較社会文化研究院地球変動講座、2.インド科学大学、3.九州大学大学院地球社会統合科学府、4.ボツワナ大学、5.ペラデニヤ大学

1.Division of Evolution of Earth Environments, Faculty of Social and Cultural Studies, Kyushu University, 2.Centre for Earth Sciences, Indian Institute of Science, 3.Graduate School of Integrated Sciences for Global Society, Kyushu University, 4.Department of Geology, University of Botswana, 5.Department of Geology, Faculty of Science, University of Peradeniya

The high-grade ultrahigh-temperature regional metamorphism of Sri Lanka has a significant role in understanding the tectonics and formation of Gondwana super-continent. Sri Lanka in specific is important because of its position in Gondwana assembly, placed close to southern India, Madagascar and eastern Antarctica. The central Highland complex in Sri Lanka consists of UHT granulites intercalated with pelitic and mafic granulites along with calc-silicates, which preserves textural evidence for UHT metamorphism. The U-Pb zircon dating of sapphirine bearing granulites yielded two major age populations 565-620 Ma and 525-563 Ma without older zircon cores. Another granulite sample with Grt-Sil-Spl-Crd assemblage has similar metamorphic age ranging *ca.* 526, 563 and 602 Ma concordant clusters, with inherited zircon core ages from 1600 to 2040 Ma. The Fe-Al rich pelitic granulites also had undergone bimodal metamorphism (520-565 Ma and 590-622 Ma). Some of these samples have inherited zircon cores ranging from with 760 to 3060 Ma. The zircons in mafic granulite samples have age range of 533-566 Ma and 578-620 Ma. The scapolite bearing calc-silicate sample also has similar age populations at *ca.* 533 Ma and 560-570 Ma. The compilation of the new results and available ages from Sri Lanka and its adjacent continental fragments of the Gondwana, we propose the Northern Madurai Block, Southern Madurai Block, Wannai Complex and Highland Complex are alternate older and younger terranes formed in a major convergent regime, metamorphosed during Ediacaran-Cambrian regional UHT event. The petrological and geochronological similarities indicate that the Highland Complex, Sri Lanka, Madurai Block, southern India and Lützow-Holm Complex, Antarctica and central Madagascar should have been together during the closure of the Mozambique Ocean as part of the Gondwana amalgamation.

キーワード：ジルコン年代、原生代後期、超高温変成岩、スリランカ・ハイランド岩体

Keywords: U-Pb zircon LA-ICP-MS dating, Late Neoproterozoic, ultrahigh-temperature granulite, Highland Complex, Sri Lanka

## スリランカHighland Complexに産する変成炭酸塩岩の堆積年代

## Depositional ages for metacarbonate rocks from the Highland Complex, Sri Lanka

\*白川 桃子<sup>1</sup>、Satish-Kumar Madhusoodhan<sup>2</sup>、Malaviarachchi Sanjeewa P.K.<sup>3</sup>\*Momoko Shirakawa<sup>1</sup>, Madhusoodhan Satish-Kumar<sup>2</sup>, Sanjeewa P.K. Malaviarachchi<sup>3</sup>

1.新潟大学大学院自然科学研究科、2.新潟大学理学部地質科学科、3.ペラデニア大学

1.Graduate School of Science and Technology, Niigata University, Japan, 2.Department of Geology, Faculty of Science, Niigata University, Japan, 3.Department of Geology, Faculty of Science, University of Peradeniya, Peradeniya, Sri Lanka

Sri Lanka is an integral part of the Late Proterozoic to Early Cambrian collision zone of the so-called East African-Antarctic Orogen (EAAO) that includes Mozambique, Madagascar, southern India and the Dronning Maud Land in East Antarctica. The Mozambique Ocean is supposed to have existed between the different continents between East and West Gondwana, before its final amalgamation. The carbonate rocks in the EAAO are considered to have chemically deposited. Therefore, these rocks are ideal for obtaining the geochemical information of the Mozambique Ocean. The 540-600 Ma metasedimentary rocks in the Highland Complex (HC) is the only unit in Sri Lankan basement containing metacarbonate rocks. In this study, we attempt to apply the strontium (Sr) isotope chemostratigraphy to estimate the sedimentation age of the Highland Complex. In order to estimate the depositional age, it is necessary to consider the influence on carbonate rocks by metamorphism. For example, the oxygen isotope records fluid-rock interaction, while carbon isotopes alter with lithification or rare earth element + yttrium pattern changes by the contamination with the mixing of continental rocks. In addition, it is necessary to consider the effect of radioactive decay of rubidium to strontium. Thus in this study, taking advantage of the characteristics of metacarbonate rocks, samples that were least influenced by alteration were selected by chemical screening. For this purpose oxygen and carbon isotopic composition and trace and rare earth element pattern and Sr isotope ratio were used.  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values of the samples range between 19 to 25 ‰ and -2.0 to 1.5 ‰. The results indicate that they preserve the primary depositional marine chemical composition. REE + yttrium patterns had oceanic characteristics as well. However, the Sr initial ratios have regional differences,  $^{86}\text{Sr}/^{84}\text{Sr}$  (850 Ma) values ranges from 0.70431 to 0.70711. These values when correlated with the late Proterozoic Sr isotope chemostratigraphic curve of Halverson et al. (2010) suggest the depositional age in between 900 to 660 Ma.

The depositional ages are consistent with the inference from the youngest detrital ages of  $834 \pm 12$  Ma and  $722 \pm 14$  Ma for zircons in the metapelitic rocks, though there is considerable regional differences. Our estimate of the depositional age are also comparable with the depositional age of metacarbonate rocks from the Sør Rondane Mountains (SRM), East Antarctica (880-850 Ma and 820-790 Ma; Otsuji et al., 2013), and those of Mozambique belt (890-860 Ma; Melezhik et al., 2008). The results indicate that the carbonate sedimentation in the Mozambique Ocean persisted extensively from southeastern Africa to east Antarctica, through southern India and Sri Lanka during the late-Tonian to early-Cryogenian periods. The result also places important constraints on the temporal and spatial extent of the Mozambique Ocean, which may lead to the understanding of the processes and timing of Gondwana formation. Further detailed analytical studies are being carried out, to understand the reliability and regional extent of depositional ages for the metacarbonate rocks in the Highland Complex, Sri Lanka.

References; Dharmapriya et al. (2015), Precam. Res. 771, 311-333., Halverson et al. (2010), Precam. Res. 182, 337-350., Melezhik et al. (2008), Precam. Res. 162, 540-558., Otsuji et al. (2013), Precam. Res. 243, 257-278.

キーワード：変成炭酸塩岩、スリランカ、化学的層序学

Keywords: metacarbonate rocks, Sri Lanka, Chemostratigraphy

Neoproterozoic (ca. 860 Ma) A-1 type volcanic activity of the Gyemyeongsan metavolcanics in the Okcheon Metamorphic Belt, Korea: a prelude to the Rodinia breakup?

\*Kye-Hun Park<sup>1</sup>, Myoung-Jung Kim<sup>1</sup>

1.Dept Earth Environmental Sci, Pukyong National Univ, Korea

The Gyemyeongsan Formation is composed of mainly metavolcanic rocks and is distributed over the northeastern part of the Okcheon Metamorphic Belt. In general, constituent rocks show very high abundances of rare earth elements (REE) and high field strength elements (HFSE). In part, contents of magnetite and/or rare earth minerals approach to ore grade. Lack of noticeable Nb negative anomaly on spider diagram indicates negligible contribution of crustal material in their generation and also excludes the arc environment for their tectonic environment. If we apply the tectonic discrimination diagram utilizing immobile HFSE Nb and Y, the metatrachytes of the Gyemyeongsan Formation are categorized as within-plate products. They also show the typical features of A-type magma, such as noticeably high contents of REE, HFSE and Ga. Their geochemical characteristic can be classified further as A-1 type that is one of the distinctive features of the continental rifting. Hafnium isotopic composition of zircons yields positive initial epsilon values about 5, indicating their derivation from slightly enriched mantle. SHRIMP U-Pb zircon age determination of the metavolcanics yields eruption ages of ca. 860 Ma in the Neoproterozoic. The age of metavolcanics of the Gyemyeongsan Formation is somewhat earlier than generally accepted initiation of breakup of the supercontinent Rodinia, i.e. ca. 830 Ma based on evidences from the South China Block. However, considering the facts that the metavolcanics of the Gyemyeongsan Formation have A-1 type geochemical characteristics and hafnium isotopic signature of mantle, it would be natural to seek possible link with a continental rifting event. Probably it was a prelude to the breakup of supercontinent Rodinia.

Keywords: A-type, U-Pb zircon age, breakup, Rodinia

中部ベトナム、コンツム地塊に産するプルーム起源マグマと火成・変成作用

Late Permian plume-related magmatism and tectonothermal events in the Kontum Massif, central Vietnam

\*大和田 正明<sup>1</sup>、小山内 康人<sup>2</sup>、中野 伸彦<sup>2</sup>、足立 達朗<sup>2</sup>、北野 一平<sup>2</sup>、チャン チ<sup>3</sup>、加々美 寛雄<sup>4</sup>

\*Masaaki Owada<sup>1</sup>, Yasuhito Osanai<sup>2</sup>, Nobuhiko Nakano<sup>2</sup>, Tatsuro Adachi<sup>2</sup>, Ippei Kitano<sup>2</sup>, Tri Van Tran<sup>3</sup>, Hire Kigali<sup>4</sup>

1.山口大学大学院理工学研究科、2.九州大学比較文化研究院、3.ベトナム地質調査所、4.新潟大学自然科学研究科

1.Graduate School of Science and Engineering, Yamaguchi University, 2.Department of Earth Sciences, Kyushu University, 3.General Department of Geology and Minerals of Vietnam, 4. Graduate School of Science and Technology, Niigata University

The Kontum Massif is situated in the southern part of Trance Vietnam Orogenic Belt (TVOB), central Vietnam. The massif consists of various kinds of metamorphic rocks including ultrahigh-pressure/ultrahigh-temperature metamorphic rocks and intrusive rocks. While the geochronological data indicates two tectonothermal events (480 -420 Ma, 270 -240 Ma), the intense metamorphic and magmatic activities occurred during Late Permian -Early Triassic as a result of the continental collision between South China and Indochina cratons. In this study, geochronological analyses for three samples (2 metagabbros and 1 charnockite) were conducted with zircon U-Pb LA-ICP-MS dating. The zircon dating gave a magmatic age range of 260 -250 Ma for the three samples and an inherited age of ~1400 Ma for the charnockite. The magmatic ages were found to be identical with those of peak metamorphic periods in the Kontum Massif. These results combined with Nd isotopic data for the granitic rocks and pelitic gneisses from the Kontum Massif suggest that the massif may have been derived from a reworked continental crust. Geochemical features of the metagabbros reveal that the parental basaltic magma corresponds to the Song Da igneous suite situated in the northern part of TVOB, and has been assimilated by crustal materials. The Song Da igneous suite is a member of Emeishan large igneous province, and has been derived from the Late Permian mantle plume. It is concluded that the plume-related magma has intruded into the deeper part of Kontum Massif, and led the ultrahigh-temperature metamorphism by acting as a heat source.

キーワード：変はんれい岩、マントルプルーム、ジルコンU-Pb年代、ペルム紀後期、コンツム地塊

Keywords: metagabbro, mantle plume, zircon U-Pb dating, Late Permian, Kontum Massif

インド西ダルワールクラトンのチドラドゥルガ周辺にみられる古始生代TTGの火成活動  
Igneous activity of Paleoproterozoic TTGs around Chitradurga, western Dharwar craton, India

\*亀井 淳志<sup>1</sup>、外田 智千<sup>2</sup>、Satish-Kumar Madhusoodhan<sup>3</sup>、豊島 剛志<sup>3</sup>、三島 郁<sup>4</sup>、上野 雄一郎<sup>4</sup>、片岡 香子<sup>5</sup>  
\*Atsushi Kamei<sup>1</sup>, Tomokazu Hokada<sup>2</sup>, Madhusoodhan Satish-Kumar<sup>3</sup>, Tsuyoshi Toyoshima<sup>3</sup>, Kaoru Mishima<sup>4</sup>, Yuichiro Ueno<sup>4</sup>, Kyoko Kataoka<sup>5</sup>

1.島根大学総合理工学部、2.国立極地研究所、3.新潟大学理学部、4.東京工業大学理学部、5.新潟大学災害復興科学

1.Department of Geoscience, Interdisciplinary Faculty of Science and Engineering, Shimane University, 2.National Institute of Polar Research, 10-3, Midori-cho, Tachikawa-shi, Tokyo 190-8518, Japan, 3.Department of Geology, Niigata University, 2-8050 Ikarashi, Nishi-ku, Niigata 950-2181, Japan, 4.Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan, 5.Research Institute for Natural Hazards and Disaster Recovery, Niigata University, 2-8050 Ikarashi, Nishi-ku, Niigata 950-2181, Japan

The western Dharwar craton (WDC) comprises mainly of TTG-type peninsular gneiss, greenstone belts, and high-potassium granites (e.g. Chadwick et al., 2000; Jayananda et al., 2013). The TTGs are widely distributed in the WDC as 3.4-3.2 Ga basement rock and occasionally contain more older sediments and volcanics (i.e. the Sargur Group) (e.g., Peucat et al., 1993). The greenstone belts, named the Bababudan Group and the Chitradurga Group, unconformably overlies the TTGs as 2.9-2.6 Ga sedimentary covers (Hokada et al., 2013). These groups consist mainly of conglomerate/quartzite, BIFs, and mafic to felsic volcanics (e.g. Chadwick et al., 1981). The high-potassium granites crop out as several isolated intrusions in both the TTGs and the greenstone belts. The intrusive ages of the granites are mostly around 2.61 Ga (e.g. Jayananda et al., 2006). The aim of this study is to discuss the igneous activity in the Paleoproterozoic TTGs based on the petrographic and geochemical characteristics of the peninsular gneiss around Chitradurga in the WDC, southern India.

Although the TTGs are generally composed of granitic to tonalitic gneisses, we can group them into two types in correspondence with their petrographic features; K-feldspar (Kfs)-poor meta-trondhjemite and Kfs-rich meta-granite. The Kfs-poor meta-trondhjemite has suitable petrographic feature of slab-derived TTGs. On the other hand, the Kfs-rich meta-granite contains abundant K-feldspar, a mineral phase that is not common in the TTGs.

The Kfs-poor meta-trondhjemite (SiO<sub>2</sub>: 65-75 wt%) have typical TTGs in composition, with high Al<sub>2</sub>O<sub>3</sub> (14.7-19.1 wt%) and Na<sub>2</sub>O contents (3.6-5.2 wt%), and correlated low K<sub>2</sub>O/Na<sub>2</sub>O (0.27-0.74) as well as moderately high Sr (195-527 ppm) and Sr/Y (21-109), although the REE patterns have fairly variable from flat to fractionated ((La/Yb)<sub>n</sub>=1.2-26), with no Eu anomaly. However, the Kfs-rich meta-granite (SiO<sub>2</sub>: 69-78 wt%) shows different geochemical characteristics than that of TTGs. In particular, their Al<sub>2</sub>O<sub>3</sub> (13.4-15.7 wt%) and Sr (68-310 ppm) contents are not enriched, and their K<sub>2</sub>O/Na<sub>2</sub>O (0.5-1.9) ratios are higher. The REE patterns are largely sub-divided into two types, one is fractionated ((La/Yb)<sub>n</sub>=10-64) with no Eu anomaly, the other is moderately fractionated ((La/Yb)<sub>n</sub>=1.8-21) associated with significant negative Eu anomaly. Probably, the Kfs-rich meta-granite had been formed by several different types of magmatic activities. In Paleoproterozoic, the western Dharwar craton was intruded by many slab-derived TTGs. In addition, it is probable that the various granitic activities were derived from strong magma differentiation and/or crustal reworking operated simultaneously to form a stable continental crust.

References: Chadwick et al., 1981, Precambrian Research 16, 31-54, Chadwick et al., 2000, Precambrian Research 99, 91-101, Jayananda et al., 2006, Precambrian Research 150, 1-26, Jayananda et al., 2013, Precambrian Research 227, 55-76, Hokada et al., 2013, Precambrian Research 227,



99-119

キーワード：トーンナル岩 トロニウム岩 花崗閃緑岩、ダルワール クラトン、インド  
Keywords: TTG, Dharwar Craton, India

Timing of melt metasomatic event at arc deep crust beneath Ichinomegata crater (Northeast Japan) constrained by LA-ICP-MS U-Pb geochronology of zircon from the mafic xenoliths  
Timing of melt metasomatic event at arc deep crust beneath Ichinomegata crater (Northeast Japan) constrained by LA-ICP-MS U-Pb geochronology of zircon from the mafic xenoliths

\*柳田 泰宏<sup>1</sup>、谷 健一郎<sup>2</sup>、中村 美千彦<sup>1</sup>、吉田 武義<sup>1</sup>

\*Yasuhiro Yanagida<sup>1</sup>, Kenichiro Tani<sup>2</sup>, Michihiko Nakamura<sup>1</sup>, Takeyoshi Yoshida<sup>1</sup>

1.東北大・大学院・地学専攻、2.国立科学博物館

1.Department of Earth Science, Graduate School of Science, Tohoku University, 2.Department of Geology and Paleontology, National Museum of Nature and Science

In the Ichinomegata crater, Oga peninsula in the Northeast Japan, abundant amphibole-bearing ultramafic to mafic xenoliths occurred with andesitic magma erupted 0.06–0.08 Ma. The previous studies on their texture and isotope composition suggest that they were metasomatized by aqueous fluid or silicate melt at deep crustal levels. In order to constrain timing of the metasomatic event, we obtained <sup>206</sup>U–<sup>238</sup>Pb ages of zircon from three biotite-bearing hornblende gabbro xenoliths consisting of amphibole ( $Mg/Mg+Fe_{total} = 0.60-0.65$ ), plagioclase ( $An_{53-86}$ ), biotite ( $Mg/Mg+Fe_{total} = 0.62-0.68$ ), magnetite, apatite and zircon. Zircon grains collected from these biotite-bearing gabbros displayed sub-rounded morphology and complex zoning in cathodoluminescence and BSE images. The textural variation of zircon grains was divided into three types: (1) heterogeneous zircon, which is characterized by heterogeneous Th concentration domains with thorite and rhyolitic melt inclusion, (2) magmatic zircon characterized by oscillatory zoning, and (3) mixed zircon characterized by the “heterogeneous zircon” mantled by the “magmatic zircon”. All the zircon grains exhibited sub-rounded to rounded form. The microstructure of the heterogeneous and mixed zircons suggests that those zircon grains were subjected to recrystallization coupled with dissolution and reprecipitation (CDR) mediated by felsic melt. The rounded shape of most zircon grains suggests that they experienced dissolution in the last stage although magmatic growth followed the CDR for the mixed zircon. The spot <sup>206</sup>U–<sup>238</sup>Pb dating of heterogeneous and magmatic zircon, conducted with a LA-ICP-MS at National Museum of Nature and Science, revealed that concentrations of radioactive Pb derived from U decays in zircon were very low, close to or lower than the detection limit. This means that the recrystallization ages were significantly young (< 1.0 Ma). This is consistent with the observation that the metasomatic agent (silicate melt) has similar isotopic signature to the andesitic host magma (Yamamoto et al. 2013). The recrystallization of these zircons is likely to have occurred in the pre-eruptive activity on the felsic magmatism at a deep crust beneath the Ichinomegata crater.

Reference

Yamamoto et al. (2013) *Lithos*, 162–163, 88–106

キーワード：一ノ目瀉マール、下部地殻、苦鉄質捕獲岩、ジルコン年代、交代作用

Keywords: Ichinomegata crater, Lower crust, Mafic xenolith, Zircon age, Metasomatism

インド南部太古代ダールワール岩体チトラドゥルガ片岩帯における炭素質物質の石墨化について  
Graphitization of carbonaceous materials from the Archaean Chitradurga schist belt, Dharwar  
Craton, Southern India

\*村松 樹<sup>1</sup>、中村 佳博<sup>1</sup>、Satish-Kumar Madhusoodhan<sup>2</sup>

\*Itsuki Muramatsu<sup>1</sup>, Yoshihiro Nakamura<sup>1</sup>, Madhusoodhan Satish-Kumar<sup>2</sup>

1.新潟大学自然科学研究科、2.新潟大学地質科学科

1.Graduate School of Science & Technology, Niigata University, 2.Department of Geology, Faculty of  
Science, Niigata University

Sedimentary rocks often include small amounts of organic material as an accessory phase, called as "Carbonaceous Material" (CM), during sedimentation. In this study, we report that the structural change and the isotopic composition of CMs within the carbonate rocks from the Chitradurga schist belt, Dharwar craton, southern India. Because CM (RSCM) geothermometer have been mainly developed by using pelitic rocks, this study verifies whether graphitization of CM in carbonate rocks is suitable as a geothermometer. The carbonate rocks from this region can be divided into two types by its occurrence and mineral assemblage Type-I is partly exposed at the southern part of the study area, and it is characterized by the lens-shaped limestones in dolostones. XRD patterns, Raman spectroscopy along with SEM observations indicate that the CM contained in Type-I carbonate rocks has been metamorphosed to about 450 ~ 500°C, and the graphite show a hexagonal platy structure. In addition,  $\delta^{13}\text{C}$  value of CM ranged between -11 and -7.8 permil. On the other hand, Type-II is exposed at the northeastern part of the schist belt and is characterized by laminated dolomite which is cut across by numerous quartz and carbonate veins. The CM in Type-2 carbonate rocks has been affected by the metamorphism at about 350 ~ 400°C, and the graphite was found to be characterized by sheet-like and two-dimensional crystal structure (2H) based on the analysis with XRD and Raman spectroscopy and the observation with SEM. In addition, the  $\delta^{13}\text{C}$  value ranged between -16.1 and -10.9 permil, which is suggesting a temperature dependent equilibrium carbon isotopic fractionation between graphite and carbonate. On the other hand, CM of pelitic rocks and CM in stromatolites from this region retain original  $\delta^{13}\text{C}$  value (-29 ~ -22.9 permil) of organic matter origin. However, some samples of Type-II carbonates showed anomalous  $\delta^{13}\text{C}$  values (-16.1 permil), and the morphology of graphite differs from the Type-I. They have globule forms and secondary overgrowth was observed on the sheet-like graphite. We consider these graphite to have recrystallized under the presence of a carbon-bearing fluid. The fluid crystallized graphite was found to have globular form, heterogeneous crystallinity and negative  $\delta^{13}\text{C}$  values. Furthermore, it was found that metamorphic temperature estimated using CM in carbonate rocks have larger error even at high metamorphic grades as compared with those of the pelitic rocks. We discuss the reasons for this difference and whether the graphitization process has been affected by the difference in fluid composition, abundance of  $\text{CO}_2$  and recrystallization and growth rate of the host mineral. The results of this study might put forward questions on the suitability of RSCM geothermometer in low to medium grade metacarbonate rocks and suggests the necessity of selecting samples which are suitable for analysis through the comprehensive evaluation.

キーワード：石墨、ラマンスペクトル、炭素同位体比、X線回折

Keywords: Graphite, Raman spectra, carbon isotopes, X-ray diffraction

南インド、Chitradurga 片岩帯のIngaldhal地域 における始生代の緑色岩類の地球化学的研究  
Geochemical study of Archean greenstones of Ingaldhal Formation in the Chitradurga schist belt, Southern India

\*塩谷 佳大<sup>1</sup>、三島 郁<sup>2</sup>、高橋 俊郎<sup>3</sup>、Satish-Kumar Madhusoodhan<sup>3</sup>、豊島 剛志<sup>3</sup>

\*Yoshihiro Enya<sup>1</sup>, Kaoru Mishima<sup>2</sup>, Toshiro Takahashi<sup>3</sup>, Madhusoodhan Satish-Kumar<sup>3</sup>, Tsuyoshi Toyoshima<sup>3</sup>

1.新潟大学大学院自然科学研究科環境科学専攻、2.東京工業大学大学院地球惑星科学専攻、3.新潟大学理学部地質科学科

1.Graduate school of Science and Technology, Niigata University., 2.Department of Earth & Planetary Science, Tokyo Institute of Technology, 3.Department of Geology, Faculty of Science, Niigata University

Archean greenstone and TTG gneiss are widely distributed over the western Dharwar craton in southern India. Especially, the komatiite occurrences in the Paleoproterozoic greenstone belts have been geochemically studied well and the influence of Archean volcanic activity and mantle geodynamics are often discussed from their magma genesis.

In this study, we focused on the greenstones in the Chitradurga schist belt, western Dharwar craton, in southern India. There are many occurrences of volcanic rocks in this belt, all of which were altered by low- to medium-grade metamorphism. After carrying out a detailed thin section petrography, we performed various geochemical analyses on whole rock samples. Based on the results of petrography and geochemical analyses, the greenstones were divided into 4 units. The oldest unit A is affected by amphibolite facies metamorphism, whereas the units B and C are greenschist facies metabasalts and they usually have dendritic texture made of amphibole or pyroxene similar to the texture shown by komatiites. Rarely, pillow lava structures were observed and it indicates that the volcanic activity has occurred under subaqueous (marine?) conditions. The unit D is the youngest unit and most altered one among the 4 units studied. These rocks have high calcite content and the LOI was very high (>12wt.%). Because of this reason, we decided to select the samples that contain LOI less than 5 wt.% in order to restrict the effect of alteration in the geochemical data. The major and trace element compositions of the samples from three units can be grouped into 2 types. The first type of rocks occurring in the unit A and unit C are characterized by flat REE pattern and spider diagram. The second group of rocks, occurring in the unit B, have enriched compositions of LIL, LREE and slightly depleted HREE than the first type. In addition, Nd isotope ratio was different for these 2 types, the first group have near zero to positive Nd values compared to negative values for the second group.

Three possibilities were considered based on the geochemical differences observed: the difference in the degree of partial melting, the effect of crustal contamination and the changes in tectonic setting. The Nd isotopic data and various discrimination diagrams indicate that the source mantle was different. So it is difficult that this difference occurred by the partial melting. Secondly regarding the crustal contamination, we calculated AFC model, but in this model the trace elements didn't follow the AFC curves. In addition, it was recognized that if the difference had been generated by contamination it is necessary to link it with a felsic contaminant such as continental crust. However, the volcanic activity at this time has occurred in a marine environment where a corresponding contaminant is lacking. Considering these points, it was concluded that the compositional variations observed between the two types of volcanic rocks were caused by the difference in source magma genesis due to a difference in tectonic settings. The units A and C have likely oceanic ridge type characteristics as evidenced by the REE patterns and Nd isotopic ratio

related to a possible upwelling mantle plume. On the other hand, the unit B volcanism can be related to an arc setting possibly caused by subduction. However, some features of geochemical discrimination diagrams suggested that units A, B and C have formed under different settings, which needs further evaluation. The difference in tectonics setting is also complemented by negative epsilon Nd values suggestive of possible sediment input by subduction. In summary, the geochemical characteristics shown by the Ingaldhal volcanic rocks represent a drastic change in the tectonic setting related to plate tectonics as early as MesoArchaean.

キーワード：始生代、緑色岩、Nd 同位体、ダールワールクラトン

Keywords: Archean, greenstone, Nd isotope, Dharwar craton

南インドダールワークラトン、チトラドゥルガ片岩帯中の縞状鉄鉱層の堆積環境に関する研究  
Depositional environment of Archaean BIFs in the Chitradurga Schist Belt, Dharwar Craton,  
Southern India

鯉沼 健太郎<sup>1</sup>、\*Satish-Kumar Madhusoodhan<sup>2</sup>、三島 郁<sup>3</sup>、上野 雄一郎<sup>3</sup>、外田 智千<sup>4</sup>、豊島 剛志<sup>2</sup>  
Kentaro Koinuma<sup>1</sup>、\*Madhusoodhan Satish-Kumar<sup>2</sup>、Kaoru Mishima<sup>3</sup>、Yuichiro Ueno<sup>3</sup>、Tomokazu Hokada<sup>4</sup>、  
Tsuyoshi Toyoshima<sup>2</sup>

1.新潟大学大学院自然科学研究科、2.新潟大学理学部地質科学科、3.東京工業大学大学院地球惑星科学専攻、4.国立極地研究所

1.Graduate School of Science & Technology Niigata University, 2.Department of Geology, Faculty of Science, Niigata University, 3.Department of Earth & Planetary Sciences, Tokyo Institute of Technology, Tokyo, Japan, 4.National Institute of Polar Research, Tachikawa, Tokyo, Japan

Banded Iron Formations (BIFs) are chemically precipitated from the ocean and formed mostly in the Archaean and early Paleoproterozoic, and therefore record the information of the ancient oceanic environment. In this study we present the geochemical characteristics, in particular the REY (rare earth elements plus yttrium) and Nd isotopic composition of BIFs in the Chitradurga Schist Belt, western Dharwar craton, Southern India. The Chitradurga Schist Belt is a typical Archean greenstone belt that preserves strata roughly between 3.3 Ga and 2.5 Ga and three BIF distinct layers of different sedimentary age occur in this region. Correlating each BIFs located on different region is very difficult because Chitradurga Schist Belt is deformed highly with very complex geological structures. Therefore, we compared their geochemistry to reveal the formation history and discuss their sedimentary environment.

Chitradurga BIFs are mostly comprised of quartz and iron oxide such as hematite and magnetite, similar to many other Archean BIFs. Bulk rock geochemistry results revealed that the BIFs contain only very less amount of continental components, such as  $Al_2O_3$  (<1wt.%) and  $TiO_2$  (<0.05wt.%). Chitradurga BIFs are classified into four groups by REY pattern normalized with PAAS (Post-Archean Average Shale). Type-1 has positive Eu anomaly and Y anomaly. Type-2 has no positive Eu anomaly, large Y anomaly and some of them have negative Ce anomaly. Type-3 has extremely high positive Eu anomaly, whereas Y anomaly is absent. Type-4 has positive Eu anomaly and no Y anomaly. Most of the Chitradurga BIFs have positive Eu anomaly, which is related to hydrothermal flux, however the epsilon Nd(T) values of all types fluctuate from negative to positive values. The majority of epsilon Nd(T) values for Type-3 BIFs show only small variation between -1 and +2. Moreover, on a meso-band scale the REY patterns are similar, but epsilon Nd(T) values of neighboring layers have negative and positive values, suggesting a short lived fluctuations Nd source. However, the epsilon Nd(T) values of Type-1, 2 and 4 BIFs show a broad range between -8 and +18, deviating from the normal mantle-continent range, possibly due to secondary alteration effects.

To understand the environment of deposition, we focused on Type-3 BIFs, interbedded with pillow metabasalts and meta-pyroclastic rocks. Deposition of Type-3 BIFs is perhaps influenced of two different volcanic activities: one is enriched and the other is depleted in Nd contents. The Nd isotopic ratio of Archaean seawater was dominated by positive Nd input through active hydrothermal flux related to volcanic activity from a depleted mantle source (Alexander et al., 2009). This suggests that the depositional environment of Type-3 BIFs can be modelled by a mixing between seawater with positive epsilon Nd values and hydrothermal flux derived from enriched mantle with negative epsilon Nd values, and the variations we observe depends on the hydrothermal flux from enriched mantle. This result is consistent with the REY characteristics, large positive Eu anomaly and low Y/Ho value, and suggest a deep sea hydrothermally controlled depositional environment for

the Type-3 BIFs. Type-1 BIFs are associated with dolomite, have high Y/Ho ratio, positive Eu anomaly and broad range of Nd isotope ratio, which suggests that they were deposited in a shallow sea environment. The depositional site of Type-2 and 4 BIFs could not be constrained well due to lack of unaltered samples, but it is assumed that Type-2 and 4 BIFs were affected by less hydrothermal input, when compared with Type-3 BIFs. In summary, our results indicate that negative epsilon Nd(*T*) values does not necessarily need a source from continental crust, but can result from the mixing of Nd from an enriched mantle with Nd reservoir in the seawater.

キーワード：縞状鉄鉱層、ダールワールクラトン、Nd同位体

Keywords: Banded Iron Formation, Dahrwar craton, Nd isotope