

## The Global Earthquake Activity Rate (GEAR) forecast test

\*David D Jackson<sup>1</sup>, Peter Bird<sup>1</sup>, Yan Y Kagan<sup>1</sup>, Corne Kreemer<sup>2</sup>, Ross Stein<sup>3</sup>

1.University of California Los Angeles, 2.University of Nevada Reno, 3.Temblor.net

We present a global seismicity reference model based on uniform global datasets, transparently merging the best features of competing approaches and resulting in a testable product. In the past, two approaches to forecasting shallow seismicity have had comparable success. The tectonic method is based on maps of deforming regions and/or active faults, with some empirical calibration of seismic coupling. This captures the distribution of energy sources and may provide good forecasts for very long time windows. The smoothed seismicity approach applies optimized smoothing to cataloged earthquakes. This captures triggering, including ongoing aftershock sequences, and may work best on short to intermediate timescales. Here we combine a leading global forecast of each type using several kinds of combination: linear, log-linear, and an envelope method. We test the success of each parent and hybrid forecast in an 8-year retrospective test by information score, area skill score, and spatial-likelihood metrics that are all roughly independent of total earthquake rate. In this 2005-2012 test, the most successful hybrid model is the log-linear mixture of 60% seismicity with 40% tectonics. This hybrid outperforms both parent forecasts. The chance that this improvement results from a temporary random fluctuation is much less than 1%. We also test all models against the analog-instrumental catalog years 1918-1976; the same patterns of hybrid improvement are found. Likelihood scores are generally less than for the more recent catalog, possibly because of limitations in the older data. We compute an update of the preferred hybrid model using all modern catalog years 1977-2013, for future prospective testing. This hybrid model is named Global Earthquake Activity Rate model 1 (GEAR1) and is provided on a  $0.1^\circ \times 0.1^\circ$  global grid, for hypocentroid depths up to 70 km, with magnitude bins whose centers range from 6.0 to 9.0 in steps of 0.10. Comparing our GEAR1 forecast to the recent fault-based UCERF3 long-term forecast in California, we find that both predict the same total earthquake rates (within 4%) at each of two thresholds, but that the map patterns of the GEAR1 forecast most strongly resemble the maps of UCERF3 after spatial smoothing with characteristic distances of 25-30 km has been applied to UCERF3.

Keywords: earthquake, forecast, global, test

## Hybrid Models and Time-dependent Hazard in New Zealand

\*Matt Gerstenberger<sup>1</sup>, David Rhoades, Annemarie Christophersen, David Harte, Bill Fry

### 1.GNS Science

Recent work in earthquake forecasting in New Zealand has been at the interface of earthquake forecasting (or operational earthquake forecasting) and seismic hazard analysis. One of our aims has been to develop models that can transition from short-term forecasting to the time scales that are required by seismic hazard analysis. As part of this we have been developing hybrid forecast models that utilise alternative data sets to the earthquake catalogue. Recent results have shown significant improvements in forecast skill by including geological information and geodetic strain rate information. Another important challenge has been to develop homogeneous catalogues that allow for consistent forecasting from earthquake rates through to ground-motion prediction equations. We have also been endeavoring to understand the impact of uncertainties (e.g., from the catalogue through to model uncertainty) on the final forecasts. Finally, we will discuss some of the challenges we are currently facing in testing of earthquake forecast models.

Keywords: earthquake forecasting, statistics, seismic hazard

## 関東直下の3次元地震活動モデルについて

## 3D spatial models for seismicity beneath Kanto region

\*尾形 良彦<sup>1,2</sup>、桂 康一<sup>1</sup>、鶴岡 弘<sup>2</sup>、平田 直<sup>2</sup>\*Yoshihiko Ogata<sup>1,2</sup>, Koichi Katsura<sup>1</sup>, Hiroshi Tsuruoka<sup>2</sup>, Naoshi Hirata<sup>2</sup>

1.統計数理研究所、2.東京大学 地震研究所

1.The Institute of Statistical Mathematics, Research Organization of Information and Systems,

2.Earthquake Research Institute, University of Tokyo

経度、緯度および深さに関する3次元空間の統計モデルの開発は関東平野直下100km深までの地震活動の解析・予測のために特に必要である。関東直下ではフィリピン海プレートと太平洋プレートのスラブが重なって北米プレートに沈み込み、各プレート間やプレート内の地震活動の相互作用が複雑であり、2次元以下の時空間モデルでは詳細な解析結果や確率予測を得ることが難しいためである。

3次元空間の層的時空間ETASモデルは、任意の時刻  $t$  と位置  $(x, y, z)$  での地震活動度が過去の地震発生履歴の関数として記述されるものである。なかでも常時地震活動度  $\mu$  (background seismicity) や余震生産性  $K$  (aftershock productivity) は、地域性的変化が特に大きいので、位置の関数と考え地震活動の特徴を定量化する。さらに現在の関東直下の地震活動度は東北地方太平洋沖地震によって誘発された活動度として大森・宇津型の関数で記述する。ここで関東直下における東北沖地震からの影響度  $K_{M9}$  の地域的な違いを位置依存性として定量化する。さらに将来の大地震の確率予測のために関東直下におけるマグニチュード頻度分布が必要でGutenberg-Richter則の  $b$ -値にも3次元位置依存性を仮定する。これらの解により関東直下の地震活動を特徴づけ、大地震の長期予測や余震の確率予測に適用する。

かくて、震源データから関数  $\mu(x, y, z)$ ,  $K(x, y, z)$ ,  $K_{M9}(x, y, z)$  及び  $b(x, y, z)$  を定義し推定するために、関東直下100kmまでの直方体を震源などを頂点とする小4面体にデロネ分割し、これらの関数型を決める係数はデロネ多面体網の頂点での値であり、任意の位置  $(x, y, z)$  における関数値は、その位置を囲むデロネ多面体 (tetrahedron) 内で線形的に内挿したものである。このような局所的に線形な関数をデロネ関数と呼ぶ。しかし、それらの係数は極めて多数なので最尤法は適用不可であり、求める係数間に制約を付けて安定した解を求める。すなわちデロネ関数の変化が平滑になるように、デロネ関数の局所的な傾きにデロネ多面体の体積を乗じたものの2乗和をペナルティ関数とし、これを付加した対数尤度 (penalized log likelihood) を考える。ペナルティの強さを赤池バイズ情報量規準 (経験的バイズ法) の最小化によって決め、ペナルティ付き対数尤度を最大化する係数を最適解として得る。

キーワード：赤池バイズ情報量規準、余震生産性、常時地震活動、 $b$ -値、デロネ関数、誘発地震活動Keywords: ABIC, aftershock productivity, background seismicity rate,  $b$ -values, Delaunay function, Omori-Utsu function for induced seismicity

## Optimized physics-based earthquake forecasts for inland Japan

\*Margarita Segkou<sup>1</sup>, Jiancang Zhuang<sup>2</sup>

1.BGS, 2.ISM

We focus on the stress recovery processes after the  $M=9.0$  Tohoku mega-earthquake and how the above influences the earthquake probabilities in active faults of inland Japan. Recent studies present evidence about rapid stress recovery near the trench but as anticipated by rate-and-state friction law, returning to pre-Tohoku "normal" seismicity levels for the active faults in inland Japan is a slower process. We perform a retrospective forecast spanning 10 years (2004-2014) in the Niigata prefecture (mid-west Japan) using physics-based modeling, combining rate-and-state law and Coulomb stress changes, to study earthquake-triggering mechanisms. The key element of our innovation lies in the development of 429 forecast models, as a result of a stochastic optimization within short-time frames (10 days), in order to access the variability of fault constitutive parameters. The aforementioned optimized forecast then competes with a benchmark statistical/empirical Epidemic-Type Aftershock Sequence (ETAS) model already submitted in CSEP-Japan. The testing period starts with the  $M=6.8$  2004 Chuetsu mainshock and ends on December 2014, approximately 9 months after the  $M=6.7$  post-Tohoku mid-Niigata on April 12<sup>th</sup>, 2011. Our physics-based optimization goes hand in hand with uncertainty consideration related with the estimation of static stress changes (geometry of active faults, receiver depth, effective friction coefficient) following the  $M=6.8$  Chuetsu mainshock and important triggered events at the near source are, the 2007  $M=6.7$  Chuetsu-Oki and the  $M=6.2$  post-Tohoku event. The forecasts are evaluated for their predictability, spatial consistency and relative information gain through log-statistics by considering the statistical model as reference. We find that: (1) best-fit solutions correspond to stressing rates between 0.01 and 0.8 bar/yr immediately after the aforementioned mainshocks and (2)  $\Delta\sigma$  values vary between 0.1 and 2.0, for the few first days following the post-Tohoku and Chuetsu events.

Keywords: physics-based forecasting, stress changes, post-Tohoku recovery

## Resolving Stress Singularities: a Retrospective Japan Rate-and-State Earthquake Forecast

\*Anne Elizabeth Strader<sup>1</sup>, Yosihiko Ogata<sup>3,2</sup>, Naoshi Hirata<sup>2</sup>, Hiroshi Tsuruoka<sup>2</sup>, Danijel Schorlemmer<sup>1</sup>

1.GFZ Potsdam, 2.ERI, The University of Tokyo, 3.The Institute of Statistical Mathematics

Retrospective evaluations of rate-and-state Coulomb stress transfer have shown consistent associations between increased Coulomb stress and seismicity rates. However, stress singularities occurring at the ends of fault dislocation patches tend to provide an unrealistic calculated stress field near active faults, where most earthquakes occur. The effects of such stress calculation artefacts

may be mitigated through implementation of an inverse rate-and-state model, where seismicity rate variations are inverted to obtain Coulomb stress steps over time. The resulting stress variations, from which expected seismicity rates are derived, show potential near the Tohoku rupture plane in a short-term prospective Japan forecast due to low magnitude completeness thresholds, which allow for comprehensive delineation of the Coulomb stress field. An additional advantage of the stress inversion model is that it does not require receiver plane orientations or focal mechanisms, which add uncertainty to the calculated stress field and are not always available, limiting the reliability

of applying the forward model in a global earthquake forecast. We retrospectively test our forecast within all Japan CSEP (Collaboratory for the Study of Earthquake Predictability) testing regions, just after the 2011 Tohoku earthquake. To determine whether the observed seismicity rate variations are sufficient to define Coulomb stress field variations, we combine our forecast with ETAS forecasts currently being tested in the Japan CSEP testing center. At the 95% significance level, the rate-and-state forecast displays potential in defining the magnitude distribution of future earth-

quakes, but does not yet reliably constrain the number, or the spatial distribution of earthquakes away from the mainshock, indicating that the inverse forecast may be most effectively applied in an ensemble earthquake forecast model, where it contributes more to forecasted seismicity rates near areas with recent seismicity.

Keywords: forecast, seismology, Coulomb stress

Impact of automatic catalog quality on real-time aftershock forecasting  
Impact of automatic catalog quality on real-time aftershock forecasting

\*近江 崇宏<sup>1</sup>、尾形 良彦<sup>2,3</sup>、汐見 勝彦<sup>4</sup>、Enescu Bogdan<sup>2,5</sup>、澤崎 郁<sup>4</sup>、合原 一幸<sup>1</sup>

\*Takahiro Omi<sup>1</sup>, Yoshihiko Ogata<sup>2,3</sup>, Katsuhiko Shiomi<sup>4</sup>, Bogdan Enescu<sup>2,5</sup>, Kaoru Sawazaki<sup>4</sup>, Kazuyuki Aihara<sup>1</sup>

1.東京大学 生産技術研究所、2.統計数理研究所、3.東京大学 地震研究所、4.防災科学技術研究所、5.筑波大学 生命環境系

1.Institute of Industrial Science, the University of Tokyo, 2.The Institute of Statistical Mathematics, 3.Earthquake Research Institute, University of Tokyo, 4.National Research Institute for Earth Science and Disaster Prevention, 5.Faculty of Life and Environmental Sciences, University of Tsukuba

A hypocenter catalog is the main input for earthquake forecasting. Thus the quality of the catalog may have a considerable impact on the forecasting performance. This issue is important especially for aftershock forecasting because the short-term forecast is prepared based on the hypocenter catalog available in real-time during occurrences of aftershocks, but its quality is generally low as compared to the final version of the edited catalog. For example, many aftershocks including even moderate ones in the first few hours after a main shock are missing in real-time data. Here we are concerned with automatically determined hypocenters without any manual amendments, and we examine how the raw quality of the real-time data affects the performances of aftershock forecasting. In this study we examine the automatic hypocenter catalog of the High Sensitivity Seismograph Network (Hi-net [1]), and conduct forecast experiments of inland aftershock sequences in Japan. We compare forecasting performances between this Hi-net automatic catalog and the JMA revised catalog [2]. We also consider several automatically modified versions of the Hi-net catalog, and examine what kind of factors in the raw catalog are important for improving the forecasting performance.

[1] Okada, Y. et al. (2004), Recent progress of seismic observation networks in Japan -Hi-net, F-net, K-NET and KiK-net-. Earth, Planets and Space 56, 15-28.

[2] T. Omi, Y. Ogata, K. Shiomi, B. Enescu, K. Sawazaki, K. Aihara, "Automatic aftershock forecasting: A forecast test of aftershocks using the real-time data in Japan"  
" (in preparation)

キーワード：余震、確率予測、リアルタイムデータ

Keywords: aftershock, probabilistic forecasting, real-time data

## Universal slip statistics: from nanopillars to earthquakes

\*Dahmen A Karin<sup>1</sup>, Danijel Schorlemmer<sup>2</sup>

1.Department of Physics, University of Illinois at Urbana Champaign, 2.German Research Centre for Geosciences, Telegrafenberg, 14473 Potsdam, Germany

The deformation of many solid materials is not continuous, but discrete and jerky, with sudden, intermittent slips, similar to earthquakes. We discuss a simple model that predicts that the statistical distributions of the slips should be universal, i.e. they should be the same for many different materials, spanning a wide range of scales, from nanometer-sized crystalline pillars to earthquake faults that are a hundred kilometers long. We show a comparison of the model predictions to recent experiments on many different materials, ranging from nanocrystals, to bulk metallic glasses, to granular materials, to earthquakes and find good agreement with the model predictions. Tools from the theory of phase transition, such as the renormalization group can be used to explain the wide applicability of the simple model. The study provides intuition and a unified framework to understand the fundamental properties of shear-induced deformation in systems ranging from nanocrystals to earthquakes. It also provides many new predictions for future experiments, observations, and simulations. The results can be used for materials testing, evaluation, and hazard prevention.

### Reference:

Jonathan T. Uhl, Shivesh Pathak, Danijel Schorlemmer, Xin Liu, Ryan Swindeman, Braden A.W. Brinkman,, Michael LeBlanc, Georgios Tsekenis, Nir Friedman, Robert Behringer, Dmitry Denisov, Peter Schall, Xiaojun Gu, Wendelin J. Wright, Todd Hufnagel, Andrew Jennings, Julia R. Greer, P.K. Liaw, Thorsten Becker, Georg Dresen, and Karin A. Dahmen, Universal Quake Statistics: From Compressed Nanocrystals to Earthquakes, Scientific Reports 5, 16493 (2015)

Keywords: Earthquakes, Statistics, Universality, Seismology, Experiments, Scaling

## Evolution of earthquake rupture potential along the Pacific Plate off Japan, inferred from seismicity

Thessa Tormann<sup>1</sup>, Stefan Wiemer<sup>1</sup>, \*Bogdan Enescu<sup>2,3</sup>, Jochen Woessner<sup>4</sup>

1.ETH Zurich, Switzerland, 2.University of Tsukuba, Japan, 3.Institute of Statistical Mathematics, Tokyo, Japan, 4.Risk Management Solutions Inc., Zurich, Switzerland

One of the major unresolved questions in Seismology is the evolution in time and space of the earthquake rupture potential and thus time-dependent hazard along active faults. What happens after a major event: is the potential for further large events reduced as predicted from elastic rebound, or increased as proposed by current-state short-term clustering models? How does the rupture potential distribute in space, i.e. does it reveal imprints of stress transfer?

Based on the rich earthquake record along the Pacific Plate off Japan we investigate what information on spatial distributions and temporal changes of normalized rupture potential (*NRP*) for different magnitudes can be derived from time-varying, local statistical characteristics of well and frequently observed small-to-moderate seismicity. The *NRP* is obtained from the frequency-magnitude distribution of sampled earthquakes, specifically from the slope (*b*-value) and *y*-intercept (*a*-value) of this distribution, in a log-linear plot. The *b*-values describe the relative frequency of large versus small earthquakes, while *a*-values express the seismic activity during the observation period (in this study, the *a*-values are annualized and distance-weighted, i.e. we consider the relative earthquake-grid-point distances, with close-by events gaining higher weights than more distant events). We analyze the seismicity from 1998 ~ 2015, including the massive 2011 M9 Tohoku-oki earthquake and its aftermath.

Seismicity records show strong spatio-temporal variability in both activity rates and size distribution. We show first (Tormann et al., 2015) that the size distribution of earthquakes has significantly changed before (increased fraction of larger magnitudes -relatively small *b*-values) and after that mainshock (increased fraction of smaller magnitudes -relatively large *b*-values); these changes are particularly stronger in areas of highest Tohoku-oki coseismic slip. Remarkably, a rapid recovery of this effect is observed within only few years.

We then combine this significant temporal variability in earthquake size distributions (*b*-values) with local activity rates (*a*-values) and infer the evolution of *NRP* distributions. We study complex spatial patterns and how they evolve with time, focusing on the detailed temporal characteristics in a simplified spatial selection, i.e. inside and outside the high slip zone of the M9 earthquake. We resolve an immediate and strong *NRP* increase for large events prior to the Tohoku-oki event in the subsequent high slip patch and a very rapid decrease inside this high-stress-release area, coupled with a lasting increase of *NRP* in the immediate surroundings. Even in the center of the Tohoku rupture, the *NRP* for large magnitudes has not dropped below long-term average values and is now not significantly different from conditions a decade before the M9 event.

Keywords: Tohoku-oki earthquake, seismic hazard, subduction zone seismicity



## ISC Data for Earthquake Statistics

\*Dmitry Storchak<sup>1</sup>, Domenico Di Giacomo<sup>1</sup>, James Harris<sup>1</sup>

### 1. International Seismological Centre

The International Seismological Centre (ISC) is charged with production of the ISC Bulletin –the definitive summary of the global seismicity covering the entire period of instrumental recordings between 1904 and 2016. It is based on the reviewed seismic bulletin reports from over 130 seismic networks worldwide and includes a total of 5.8 million seismic events with 1.9 millions of them reviewed by the ISC analysts. The main purpose of this huge dataset is to include as many seismic events around the globe as possible to provide the most complete and comprehensive list of earthquakes and other seismic events in a homogeneous way. In fact, there are considerable temporal and region-to-region variations in the magnitude completeness and hypocentre location accuracy of reports from local and regional seismic networks and observatories. These variations are dictated by the distribution and quality of local seismic networks, waveform processing procedures as well as administrative and political decisions that control reporting bulletin data to the ISC. The content of local bulletin reports have also changed dramatically through the period of instrumental recordings thus affecting the content of the ISC Bulletin as a whole. Statistical studies of seismicity patterns, b-value, earthquake rate are also subject to an accurate identification of anthropogenic events or events caused by anthropogenic activities. These studies often require the removal of aftershocks series. This presentation lays out major principles of building the ISC Bulletin, describes the complexity of event information, and provides recommendations for practical use of the enclosed information.

In addition to the Bulletin as its flagship product, the ISC is maintaining a set of specialized products that both narrow down and extend the information from the ISC Bulletin to serve the purpose of specific groups of geoscientists. Thus we prepared and continue extending the ISC-GEM Global Instrumental Earthquake Catalogue widely used for estimation of global and regional earthquake hazard. The ISC-EHB dataset is maintained for users interested in less complete list of events with considerably higher accuracy of event epicenter and depth determination. It is used in global and regional studies of tectonics, surface geology and inner structure of the Earth. Other datasets include the GT bulletin, CTBTO Link, Event Bibliography and Seismological Contacts.

Keywords: earthquake, global, bulletin

## 気象庁マグニチュードによる微小地震の大きさの過小評価と地震統計への影響

Underestimate of the size of microearthquakes by the JMA magnitude scale and its influence to earthquake statistics

\*内出 崇彦<sup>1</sup>、今西 和俊<sup>1</sup>

\*Takahiko Uchide<sup>1</sup>, Kazutoshi Imanishi<sup>1</sup>

1.産業技術総合研究所 地質調査総合センター 活断層・火山研究部門

1. Research Institute of Earthquake and Volcano Geology, Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology (AIST)

Earthquake statistics needs parameterized information on earthquakes. One of such parameters is the magnitude. The local magnitude scales, such as the JMA magnitude ( $M_j$ ), based on amplitudes of seismograms are easy to estimate and therefore usually included in earthquake catalogs. The moment magnitude ( $M_w$ ) is based on the physical source parameter, seismic moment, however needs much effort for the estimation especially for microearthquakes. Though the consistency between  $M_j$  and  $M_w$  is guaranteed for the medium earthquakes, we need to check that for microearthquakes.

As for use of earthquake catalogs, we should know the completeness magnitude above which catalog is complete. A type of it is  $M_c$  defined as a magnitude where magnitude-frequency distribution starts deviating from the Gutenberg-Richter's (GR) law. Another one is based on earthquake detectability. Schorlemmer and Woessner [2008] proposed  $M_p$  based on the detectability inferred from the pick information. They showed the Californian case that  $M_p$  is smaller than  $M_c$ , which indicates the breakdown of the GR law. It is important to confirm if the breakdown really occurs. Our study investigates if the discrepancies are also seen in case of  $M_w$ .

#### $M_w$ Estimation for Microearthquakes

We stably estimate seismic moment of microearthquakes based on moment ratios to nearby small earthquakes whose seismic moments are available in the NIED MT catalog, by a multiple spectral ratio analysis [Uchide and Imanishi, under review]. Applying this method to earthquakes in Fukushima Hamadori and northern Ibaraki prefecture areas, eventually we obtained the seismic moments of a total of 19140 earthquakes ( $M_j$  0.4 - 3.8). The striking result of this study is that the change in slopes of the  $M_j$ - $M_w$  curve: 1 and 0.5 at higher and lower magnitudes, respectively (see Figure). The discrepancies between  $M_j$  and  $M_w$  are significant for microearthquakes, suggesting that  $M_j$  underestimates the sizes of microearthquakes.

#### Completeness Magnitudes and b-values

The result above must affect earthquake statistics. Here we study  $M_c$  and b-value of the GR law. Following Ogata and Katsura [1993], we assume the earthquake detectability as the cumulative normal distribution with a mean,  $\mu$ , and a standard deviation,  $\sigma$ , and estimate the GR parameters (a and b) together with  $\mu$  and  $\sigma$ . We define  $M_c = \mu + 2.33\sigma$  where the detection rate is 99%. Applying this method to the monthly seismicity data in the study area, we found that the  $M_c$  for  $M_w$  is lower than that for  $M_j$  converted into  $M_w$ , however still larger than  $M_p$  converted into  $M_w$ . This may be due to the breakdown of the GR law for microearthquakes, though another possibility is that the incompleteness of earthquake catalog overestimates the detectability, resulting the underestimate of  $M_p$ .

b-values for  $M_w$  ( $b_w$ ) are systematically larger than those for  $M_j$  ( $b_j$ ). The temporal trends for  $b_w$  and  $b_j$  are similar to each other. When  $b_j$  increases,  $b_w$  also increases. This does not affect discussions inferred from the qualitative temporal change in b-values [e.g., Nanjo et al., 2012].  $b_w$  is often larger than 1.5, indicating that the moment release is dominantly done by smaller earthquakes.


### Acknowledgement

We used the JMA Unified Earthquake Catalog, seismograms from NIED Hi-net and the NIED moment tensor catalog.

Figure: Comparison between  $M_j$  and  $M_w$  inferred from the multiple spectral ratio analyses (color image for the distribution and circles for the median  $M_w$ ) and the NIED MT solutions.

キーワード：地震統計、気象庁マグニチュード、モーメントマグニチュード、スペクトル解析、コンプリートネスマグニチュード、b値

Keywords: Earthquake statistics, JMA magnitude, Moment magnitude, Spectral analysis, Completeness magnitude, b-value



Assessment and optimization of the potential earthquake precursory information in ULF magnetic data registered at Kanto, Japan during 2000 -2010

Assessment and optimization of the potential earthquake precursory information in ULF magnetic data registered at Kanto, Japan during 2000 -2010

\*Han Peng<sup>1,2</sup>、Hattori Katsumi<sup>1</sup>、Zhuang Jiancang<sup>2</sup>

\*Peng Han<sup>1,2</sup>、Katsumi Hattori<sup>1</sup>、Jiancang Zhuang<sup>2</sup>

1.Graduate School of Science, Chiba University, Chiba, Japan、2.Institute of Statistical Mathematics, Tokyo, Japan

1.Graduate School of Science, Chiba University, Chiba, Japan, 2.Institute of Statistical Mathematics, Tokyo, Japan

In order to clarify the ULF seismo-magnetic phenomena, a sensitive geomagnetic network has been installed at Kanto, Japan since 2000. In previous study, we have verified the correlation between ULF magnetic anomalies and local sizeable earthquakes by both case and statistical studies. In this paper, we use the Molchan's error diagram to evaluate the potential earthquake precursory information in the magnetic data registered at Kanto, Japan during 2000 -2010. The results show that the earthquake predictions based on magnetic anomalies are clearly better than those based on random guess, which indicates the magnetic data contain potential useful prediction information. Further investigations suggest that the prediction efficiency depends on the distance ( $R$ ) and the size of the target earthquake events ( $E_s$ ). Finally, we introduce the probability gain ( $PG'$ ) and the probability difference ( $D'$ ) to explore the optimal prediction parameters for a given ULF magnetic station. For Seikoshi (SKS) station in Izu, optimal  $R$  and  $E_s$  are about 100 km and  $10^{8.75}$ , respectively; and for Kiyosumi (KYS) station in Boso, they are about 180 km and  $10^{8.75}$ , respectively.

キーワード : earthquake precursory information、Molchan's error diagram、statistical analysis、parameter optimization、ULF magnetic data

Keywords: earthquake precursory information, Molchan's error diagram, statistical analysis, parameter optimization, ULF magnetic data

Prospective validation of physical-based precursors and their potential for short-term earthquake forecasting. Case study for Japan 2014-2016

\*Dimitar Ouzounov<sup>1</sup>, Katsumi Hattori<sup>2</sup>

1.Center of Excellence in Earth Systems Modeling & Observations (CEESMO) , Schmid College of Science & Technology Chapman University, Orange, California, USA, 2.Department of Earth Sciences, Chiba University, Chiba, Japan

We are presenting a prospective validation of short-term pre-earthquake phenomena preceding major earthquakes. Our challenge question is: "Whether such physical-based signals are significant and could be used for early warning of large earthquakes?" To address this question we have started continuous validation of atmospheric signals in retrospective/ prospective modes over Japan. Our approach is based on multidisciplinary analysis of several physical and environmental parameters (Satellite transient infrared radiation (STIR), electron concentration in the ionosphere (GPS/TEC), radon/ion activities, air temperature and seismicity patterns) that were found to be associated with earthquakes. The science rationale for multidisciplinary analysis is based on concept Lithosphere-Atmosphere-Ionosphere Coupling (LAIC) (Pulinets and Ouzounov, 2011]), which explains the synergy of different processes and anomalous variations, usually named short-term pre-earthquake anomalies.

Our validation processes consist in two steps: (1) A continuous retrospective analysis performed over two different regions with high seismicity- Taiwan and Japan for 2003-2011 (2) Prospective testing with potential for M6.5+ events Japan for 2014-2015 period. The test results suggest appearance of physical pre-earthquakes anomalies, one to several days in advance to major events, including the largest earthquakes - M7.8 of 30 May 2015 and all other M6.5+ for that period. The false alarm ratio for the testing period has shown false positives less than 20%. Our initial prospective tests show that multi-parameter analysis could reveal short-term pre-earthquake anomalies prior to the largest earthquakes in Japan.

Keywords: earthquake precursor, forecasting, early warning system

## Prospectively Evaluating the Collaboratory for the Study of Earthquake Predictability: An Evaluation of the UCERF2 and Updated Five-Year RELM Forecasts

\*Anne Elizabeth Strader<sup>1</sup>, Max Schneider<sup>2,1</sup>, Danijel Schorlemmer<sup>1</sup>, Maria Liukis<sup>3</sup>

1.GFZ Potsdam, 2.Universität Potsdam, 3.USC

The Collaboratory for the Study of Earthquake Predictability (CSEP) was developed to rigorously test earthquake forecasts retrospectively and prospectively through reproducible, completely transparent experiments within a controlled environment (Zechar et al., 2010). During 2006-2011, thirteen five-year time-invariant prospective earthquake mainshock forecasts developed by the Regional Earthquake Likelihood Models (RELM) working group were evaluated through the CSEP testing center (Schorlemmer and Gerstenberger, 2007). The number, spatial, and magnitude components of the forecasts were compared to the respective observed seismicity components using a set of consistency tests (Schorlemmer et al., 2007, Zechar et al., 2010). In the initial experiment, all but three forecast models passed every test at the 95% significance level, with all forecasts displaying consistent log-likelihoods (L-test) and magnitude distributions (M-test) with the observed seismicity. In the ten-year RELM experiment up-date, we reevaluate these earthquake forecasts over an eight-year period from 2008-2016, to determine the consistency of previous likelihood testing results over longer time intervals. Additionally, we test the Uniform California Earthquake Rupture Forecast (UCERF2), developed by the U.S. Geological Survey (USGS), and the earthquake rate model developed by the California Geological Survey (CGS) and the USGS for the National Seismic Hazard Mapping Program (NSHMP) against the RELM forecasts. Both the UCERF2 and NSHMP forecasts pass all consistency tests, though the Helmstetter et al. (2007) and Shen et al. (2007) models exhibit greater information gain per earthquake according to the T- and W- tests (Rhoades et al., 2011). Though all but three RELM forecasts pass the spatial likelihood test (S-test), multiple forecasts fail the M-test due to overprediction of the number of earthquakes during the target period. Though there is no significant difference between the UCERF2 and NSHMP models, residual scores show that the NSHMP model is preferred in locations with earthquake occurrence, due to the lower seismicity rates forecasted by the UCERF2 model.

Keywords: forecast, likelihood, statistical seismology

## Collaboratory for the Study of Earthquake Predictability - Global and Regional Results

\*Thomas Beutin<sup>1</sup>, Danijel Schorlemmer<sup>1</sup>, Naoshi Hirata<sup>2</sup>, Hiroshi Tsuruoka<sup>2</sup>, Matt Gerstenberger<sup>3</sup>, Anne Elizabeth Strader<sup>1</sup>

1.GFZ German Research Centre for Geosciences, 14473 Potsdam, Germany, 2.Earthquake Research Institute, University of Tokyo, Tokyo 113-0032, Japan, 3.GNS Science, 1 Fairway Drive, Avalon 5010, Lower Hutt 5040, New Zealand

The Collaboratory for the Study of Earthquake Predictability (CSEP) aims to improve our understanding about the physics and predictability of earthquakes through rigorous and prospective testing of earthquake forecast models. CSEP operates four testing centers in California, New Zealand, Japan, and Europe running prospective, automated evaluations of more than 430 models. These testing centers are the technical infrastructure of CSEP and implement all procedures and protocols for rigorous testing and evaluation of earthquake prediction experiments. These experiments run in various testing regions and comprise forecast periods of 30 minutes to many years.

The CSEP software system as the basis for all CSEP testing centers is now being used for earthquake early warning systems and geodetic transient detectors. The Testing and Evaluation group of the Global Earthquake Model (GEM) project at GFZ Potsdam is expanding this system to test intensity prediction equations and ground-motion prediction equations.

We present results and the key lessons learned from all major CSEP and GEM experiments, and we give an overview of recent and ongoing developments, as well as new experiments.

Keywords: Earthquake forecasting, Seismic hazard, Statistical seismology, Earthquake statistics, Forecast testing, Software



## 体積歪による群発地震予測に関する研究

Some development on predicting earthquake swarms using volumetric strain records

\*熊澤 貴雄<sup>1</sup>、尾形 良彦<sup>1,2</sup>、木村 一洋<sup>3</sup>、前田 憲二<sup>3</sup>、小林 昭夫<sup>3</sup>

\*Takao Kumazawa<sup>1</sup>, Yosihiko Ogata<sup>1,2</sup>, Kazuhiro Kimura<sup>3</sup>, Kenji Maeda<sup>3</sup>, Akio Kobayashi<sup>3</sup>

1.統計数理研究所、2.東京大学 地震研究所、3.気象庁気象研究所

1.The Institute of Statistical Mathematics, 2.Earthquake Research Institute, 3.Meteorological Research Institute

Off the east coast of the Izu Peninsula in Japan, there is a submarine volcanic region where earthquake swarms occur caused by magma intrusions. We investigated the background seismicity rates of the swarm activity by removing the triggering effect of aftershocks. We found that such background rate changes coincide with the changes of exponentially weighted averages of volumetric strain increments at the Higashi-Izu station. We further found that such a relationship consistently depends on the distance between the strainmeter station and the location of the swarm onset. The quantitative relationships revealed here may be used to monitor magma intrusions that drive the stress changes. The models we adopted here are purely statistical, but we added some comparisons of their performances with those by physically reasonable models.

キーワード：群発地震、体積歪、常時地震活動、ETASモデル

Keywords: Earthquake swarm, Volumetric strain, Background seismicity, ETAS model

## Regional evolution of network detection completeness in Japan

\*Danijel Schorlemmer<sup>1,2</sup>, Naoshi Hirata<sup>3</sup>, Yuzo Ishigaki<sup>4</sup>, Kazuyoshi Nanjo<sup>5</sup>, Hiroshi Tsuruoka<sup>3</sup>, Thomas Beutin<sup>1</sup>, Fabian Euchner<sup>6</sup>

1.GFZ German Research Centre for Geosciences, 14473 Potsdam, Germany, 2.Department of Earth Sciences, University of Southern California, Los Angeles, USA, 3.Earthquake Research Institute, University of Tokyo, Tokyo 113-0032, Japan, 4.Seismological and Volcanological Department, Japan Meteorological Agency, Tokyo 100-8122, Japan, 5.Institute of Advanced Science, Yokohama National University, Yokohama 240-8501, Japan, 6.Institute of Geophysics, ETH Zurich, Zurich, Switzerland

An important characteristic of any seismic network is its detection completeness, which should be considered a function of space and time. Many researchers rely on robust estimates of detection completeness, especially when investigating statistical parameters of earthquake occurrence like earthquake rates. Contrary to traditional approaches, we do not estimate completeness using methods in which the completeness magnitude is defined as the deviation of the frequency-magnitude distribution from the linear Gutenberg-Richter relation. Here, we present a method based on empirical data only: phase data, station information, and the network-specific attenuation relation. For each station of the network we estimate a time-dependent distribution function describing the detection capability depending on magnitude and distance to the earthquake. For each point in time, maps of detection probabilities for certain magnitudes or overall completeness levels are compiled based on these distributions. Therefore, this method allows for inspection of station performances and their evolution as well as investigations on local detection probabilities even in regions without seismic activity.

We present a full history of network detection completeness for Japan and discuss details of this evolution, e.g. the effects of the Tohoku-oki earthquake sequence. For practical purposes we deliver completeness estimates for catalog data of selected regions and document the conservative completeness estimates researchers can use when investigating the JMA catalog in different regions over different periods. All presented results are published on the CompletenessWeb ([www.completenessweb.org](http://www.completenessweb.org)) from which the user can download completeness data from all investigated regions, software codes for reproducing the results, and publication-ready and customizable figures.

Keywords: Seismic Networks, Data Quality, Completeness, Earthquake Statistics, Statistical Seismology, Earthquake Hazard

Correcting biases in the estimates of earthquake clustering parameters caused by short-term missing of aftershocks

Correcting biases in the estimates of earthquake clustering parameters caused by short-term missing of aftershocks

\*庄 建倉<sup>1</sup>、王 婷<sup>2</sup>

\*Jiancang Zhuang<sup>1</sup>, Ting Wang<sup>2</sup>

1.統計数理研究所、2.University of Otago

1.Institute of Statistical Mathematics, 2.University of Otago

Short-term missing of aftershock in the early stage after the mainshock always biases the estimates of earthquake clustering models such as the ETAS model and the Omori-Utsu formula. For example, the parameters  $c$  and  $p$  in the Omori-Utsu formula change with the cutoff magnitude threshold. To correct the biases caused by such short-term aftershock missing, we apply a method developed by Zhuang et al (2016) to replenishing missing data. The basic idea of this method is that, if a temporal point process with time independent marks is completely observed, the whole process can be transformed into a homogeneous Poisson process on the unit square by a bivariate empirical transformation. Using this method, we can simulate the missing events and re-estimate model parameter with the replenished dataset. For example, applying this method to the aftershock sequence following the 2008 Wenchuan Ms7.9 earthquake in southwestern China, the results show that the Omori parameters  $c$  and  $p$  do not change with magnitude threshold anymore and that the missing of small events in the early stage of the aftershock sequence causes the inconsistent estimate of the earthquake clustering models.

Figure: Results from applying the replenishment algorithm to the earthquake data from Southwest China. (a) Marks (magnitudes) versus occurrence times of the earthquakes.







(b) Empirical distribution of marks (magnitudes) versus empirical distribution of occurrence times of the recorded events. (c) Rescaled marks (magnitudes) versus rescaled occurrence times of the combination of the observed events, with the rescaling based on the empirical distributions of events in the time-magnitude range with complete observation. (d) Rescaled marks (magnitudes) versus rescaled occurrence times of the observed events and replenished events. (e) Marks (magnitudes) versus occurrence times of the observed events and the replenished events. (f) Cumulative numbers of events against occurrence times. Blue polygon is corresponding to the time-magnitude range in which the missing events fall. Blue dots are the replenished events.

Reference:

Zhuang, J., T. Wang and K. Kiyosugi, (2016) Detection and replenishment of missing data in marked point processes, *in preparation*.

キーワード：ETASモデル、大森・宇津律、欠測データ、余震

Keywords: ETAS model, Omori-Utsu formula, missing data, aftershock



A new algorithm to find earthquake clusters using neighboring cell connection and tests in northern Honshu, Japan

\*Wei Peng<sup>1</sup>, Shinji Toda<sup>2</sup>

1.Graduate School of Science, Tohoku University, 2.International Research Institute of Disaster Science, Tohoku University

To study the earthquake interaction, it is important to find a group of earthquakes occurred closely in space and time objectively and quantitatively. Earthquake clusters are chosen with previous clustering techniques that characterize them as mainshock-aftershock sequences or swarm sequences with empirical laws such as Omori-Utsu law or direct assumptions about physical processes such as rate/state Coulomb stress transfer, transient stress loading, fluid migration, and structural heterogeneity. Recently several papers proposed non-parameterized techniques such as kernel-based smoothing methods (e.g., Helmstetter & Werner, 2012). The cumulative rate clustering method (CURATE, Jacobs et al., 2013) is one of the approaches without any direct assumptions. The CURATE method was applied in Central Volcanic Region of New Zealand and provided a good result for selecting the swarm sequence comparing with ETAS models. However, it is still difficult to choose a proper confined area and a proper time interval for combining sequences. To reduce the arbitrary and subjective choices of space and time parameters in the CURATE method, here we propose a new method modifying the CURATE approach. We first identify the spatial clusters by looking into the spatial distribution with time in a 2-D cell-gridded map. The spatial clusters defined as a cell size ( $S$ ) which contains earthquakes and connecting its neighborhood cells if the neighborhood cells also contain earthquake events in a time window  $T$ . From the selected spatial clusters, we then evaluate temporal clustering which is defined as the increase of the transient seismicity rate at a target event comparing to the rate from the target event to the end of the sequence. This approach gives only two free parameters,  $T$  and  $S$ , for the declustering process. We tested this method for the JMA catalog and focus on the Chuetsu region (Niigata Prefecture), with earthquakes shallower than 20 km and magnitude range from 2 to 6.9. We choose the parameter ranges from  $T = 1$  to 100 days and  $S = 0.01^\circ$  to  $0.1^\circ$ , the results show that the number of the cluster events increases with longer  $T$  and larger  $S$ . By choosing the  $T = 30$  days and  $S = 0.05^\circ$ , we successfully selected the long aftershock period sequences associated with the 2004 M6.8 Chuetsu earthquake and 2007 M6.8 Chuetse-oki earthquake, while other empirical physical models and CURATE method fail to select. It suggests that this method better finds the seismic clusters including secondary aftershocks, and thus shows better declustering performance than the others.

