

観測データに基づくアジア陸域CO₂収支変動の解析Data-driven synthesis on terrestrial CO₂ budget changes in Asia*市井 和仁^{1,2}、Patra Prabir¹、植山 雅仁³、近藤 雅征¹、三枝 信子²*Kazuhito Ichii^{1,2}, Prabir Patra¹, Masahito Ueyama³, Masayuki Kondo¹, Nobuko Saigusa²

1.海洋研究開発機構 地球表層物質循環研究分野、2.国立環境研究所 地球環境研究センター、3.大阪府立大学

1.Japan Agency for Marine-Earth Science and Technology, 2.National Institute for Environmental Studies, 3.Osaka Prefecture University

Observational data provides a good constraint on understanding terrestrial CO₂ budget in a quantitative way. In recent years, more and more observation data (both satellite and ground observation) are being available, therefore, effective use of these data is an important step to improve our understandings of terrestrial CO₂ budget. In this presentation, we will show our recent progresses on the data-driven synthesis of terrestrial CO₂ budget in Asia. Topic includes (a) data-driven terrestrial CO₂ budget comparison in Asia as a continental scale case, and (b) satellite data analysis of large fire events in Southeast Asia in 2015 as a regional scale case. The first part synthesizes results of data-driven top-down and bottom-up estimations of terrestrial CO₂ budget in Asia. We used an empirically upscaled estimation of terrestrial CO₂ budget using AsiaFlux data and remote sensing data (bottom-up approach) and GOSAT Level 4A product (top-down approach). The differences of the two estimation were explained by different definition of them in Siberia and East Asia. The empirically upscaled estimation is 'net ecosystem productivity', which is a difference of gross primary production and ecosystem respiration, and GOSAT L4A is land-atmosphere net CO₂ fluxes, which includes fire, dissolved inorganic carbon export through river, and land use changes. The Southeast Asia region shows large differences between the two estimations, implying the requirement of further research.

The second part focuses on analysis of forest fire in Southeast Asia. In 2015, an intense El Nino occurred, resulting in extremely low rainfall, anomalous fire were reported in Indonesia and countries in the tropical Asia. Using multiple satellite-based data, such as aerosol optical index, land surface temperature, active fire counts, and vegetation index, the large scale massive fire event in 2015 were analyzed. We clearly detected anomalous climate and fire occurrence in Southeast Asia during August to October from MODIS active fire counts, and aerosol optical index. These data shows strong anomalous patterns compared with the normal year. We analyzed the cause of more frequent fire events in El Nino years, and found that persistent negative anomalies in precipitation is most strongly correlated with fire frequency at interannual time scales.

Acknowledgement

This study was supported by the Environment Research and Technology Development Funds (2-1401) from the Ministry of the Environment of Japan, the JAXA Global Change Observation Mission (GCOM) project (grant No. 115), and the JSPS KAKENHI (grant No. 25281003).

キーワード：物質循環、アジア、火災、陸域生物圏

Keywords: Material Cycle, Asia, Fire, Terrestrial Biosphere