

逆推計手法を応用したNO<sub>x</sub>排出量インベントリの速報アップデートシステムの開発

Application of Inversion Technique to Quick Update of Anthropogenic NO<sub>x</sub> emission over East Asia with Satellite Observations and Chemical Transport Model

\*Yumimoto Keiya<sup>1</sup>、鵜野 伊津志<sup>2</sup>、板橋 秀一<sup>3</sup>、宮崎 和幸<sup>4</sup>

\*Keiya Yumimoto<sup>1</sup>, Itsushi Uno<sup>2</sup>, Syuichi Itahashi<sup>3</sup>, Kazuyuki Miyazaki<sup>4</sup>

1.気象研究所、2.九州大学応用力学研究所、3.電力中央研究所、4.海洋研究開発機構

1.Meteorological Research Institute, 2.Research Institute for Applied Mechanics, Kyushu University, 3.Central Research Institute of Electric Power Industry, 4.Japan Agency for Marine-Earth Science and Technology

We developed a quick update system for an emission inventory with an inversion technique, and extended NO<sub>x</sub> emission in Regional Emission inventory in the ASia version 2.1 (REAS 2.1) through 2009–May 2013 with satellite-observed tropospheric NO<sub>2</sub> vertical column densities (VCDs) and a chemical transport model. The observed NO<sub>2</sub> VCDs over the eastern Chinese region exhibited a drastic inter-annual variation over the eastern Chinese region due to the socioeconomic condition. During 2008–2009, the growing of the NO<sub>2</sub> VCD became sluggish because of pollutant controls by the 2008 Beijing Olympic game and the global depression, but revived in 2010 with a growth rate of 37.3%/year. The modeled NO<sub>2</sub> VCD with the updated emission successfully followed the inter-annual variation, and reproduced the observed seasonal cycle in which summer and winter have the seasonal bottom and peak, respectively. We estimated the updated Chinese anthropogenic NO<sub>x</sub> emissions during 2009–2012 to be 25.7, 27.3, 28.2, 28.4 Tg/year; they fell within the range of the various estimates in the literatures. An annual growth rate during 2009–2012 and 2005–2012 was estimated to be 3.5%/year (0.9 Tg/year) and 5.3%/year (1.1 Tg/year), respectively. The system has the capability of updating NO<sub>x</sub> emission in near real-time (NRT) for air quality forecasting. Figure shows annual anthropogenic NO<sub>x</sub> emission from China. Numbers in the panels represent annual Chinese anthropogenic NO<sub>x</sub> emission. Other estimates of Chinese NO<sub>x</sub> emission are also shown by symbols.

キーワード：逆推計、排出インベントリ、化学輸送モデル、衛星観測、NO<sub>x</sub>排出量

Keywords: Inversion, Emission inventory, Chemical transport model, Satellite observation, NO<sub>x</sub> emission

