

ひまわり 8 号輝度温度観測のデータ同化研究:台風 Soudelor (2015)事例

Assimilating Himawari-8 Brightness Temperature: A Case Study on Typhoon Soudelor (2015)

*本田 匠¹、Lien Guo-Yuan¹、西澤 誠也¹、吉田 龍二¹、足立 幸穂¹、寺崎 康児¹、岡本 幸三^{2,1}、富田 浩文¹、別所 康太郎³、三好 建正¹

*Takumi Honda¹, Guo-Yuan Lien¹, Seiya Nishizawa¹, Ryuji Yoshida¹, Sachiho Adachi¹, Koji Terasaki¹, Kozo Okamoto^{2,1}, Hirofumi Tomita¹, Kotaro Bessho³, Takemasa Miyoshi¹

1.理化学研究所計算科学研究機構、2.気象研究所、3.気象衛星センター

1.RIKEN Advanced Institute for Computational Science, 2.Meteorological Research Institute, 3.Meteorological Satellite Center

The Japan Meteorological Agency started full operations of the new geostationary meteorological satellite "Himawari-8" in July 2015. Himawari-8 is the first of a series of the third-generation geostationary meteorological satellites including NOAA's GOES-R (planned for launch in 2016), producing about 50 times more data with more channels and 3 times more observing frequency than the previous generation. In August 2015, Himawari-8 successfully captured rapid intensification of Typhoon Soudelor (2015), the strongest northwestern Pacific typhoon in the summer of 2015 with minimum central pressure of 900 hPa. In this study, we assimilate brightness temperature from Himawari-8 using our new ensemble data assimilation system called "SCALE-LETKF" and investigate its impact on the analyses and forecasts of Soudelor.