

JRA-55で表現された下部成層圏子午面循環の変動

Changes in the lower stratospheric residual circulation in JRA-55

*小林 ちあき¹*chiaki Kobayashi¹

1. 気象研究所

1. Meteorological Research Institute

Future projections by many climate models suggest that the Brewer-Dobson circulation (BDC) will be intensified as a result of rising greenhouse gas concentrations. However, observations show a diversity of the BDC strength changes. In this study, we investigate the changes in the BDC using JRA-55 reanalysis data compared with JRA-55-related products. In JRA-55, the annual mean tropical upwelling shows a significant increasing trend in the lower stratosphere from 1979 to 2012. JRA-55C also indicate a significant increasing trend of the upwelling, but JRA-55AMIP does not. These BDC strengths are assessed by climatological zonal mean which is removed diurnal variations.

Recently, Sakazaki et al (2015) found zonally uniform tidal signals in the tropical stratosphere. The vertical wind diurnal amplitudes in the lower stratosphere is not a negligible amount compared with climatological upwelling.

The BDC trend in JRA-55 linked with the representation of the tidal signals change related to the observing system changes in the reanalysis. Comparison of the relation among the JRA-55 family members is discussed in the presentation.

キーワード：JRA-55、ブリュワードブソン循環、下部成層圏

Keywords: JRA-55, Brewer-Dobson circulation, lower stratosphere