

赤道インド洋東部の半年周期の潜流について

On semiannual equatorial undercurrents in the eastern Indian Ocean

*名倉 元樹¹、McPhaden Michael²

*Motoki Nagura¹, Michael J. McPhaden²

1.(独) 海洋研究開発機構、2.米国大気海洋庁

1.Japan Agency for Marine-Earth Science and Technology, 2.National Oceanic and Atmospheric Administration

Unlike those in the Pacific and Atlantic, the equatorial undercurrents (EUCs) in the Indian Ocean are transient and eastward only in early boreal spring and fall. Their dynamics is investigated in this study using observations obtained from four acoustic Doppler current profilers (ADCPs) deployed along the equator in the eastern Indian Ocean (78°E, 80.5°E, 83°E and 90°E). The harmonic analysis is applied to observed zonal velocity and reveals that annual and semiannual variability contributes to zonal velocity at the depth of the EUCs (about 100 m). Whereas the annual harmonic does not show any consistent tendency of zonal phase propagation, the semiannual harmonic shows eastward propagation at the depth of the EUCs. Owing to data gaps in ADCP records, the analysis is repeatedly applied to several two-year segments, and the phase speed is estimated using results obtained from various pairs of ADCPs. The results show that eastward phase propagation of the semiannual harmonic is a statistically robust feature. For a further confirmation, zonal velocity and its divergence are calculated using ADCP records. Zonal velocity leads zonal divergence, which is another evidence for eastward phase propagation. These results suggest that the semiannual transient EUCs in the Indian Ocean are Kelvin beams radiated from the surface to the east and to the depth.

キーワード：赤道潜流、インド洋、ケルビン波のビーム

Keywords: Equatorial undercurrents, Indian Ocean, Kelvin beams