

## 現場光学観測データを用いたGCOM-Cクロロフィルa濃度プロダクトの改善

Improvement of GCOM-C chlorophyll-*a* concentration product by in-situ optical measurements\*村上 浩<sup>1</sup>、清本 容子<sup>2</sup>、佐々木 宏明<sup>2</sup>\*Hiroshi Murakami<sup>1</sup>, Yoko Kiyomoto<sup>2</sup>, Hiroaki Sasaki<sup>2</sup>

1.宇宙航空研究開発機構地球観測研究センター、2.水産総合研究センター西海区水産研究所

1.Earth Observation Research Center, Japan Aerospace Exploration Agency, 2.Seikai National Fisheries Research Institute, Fisheries Research Agency

Global Change Observation Mission for Climate (GCOM-C) which carries Second-generation Global Imager (SGLI) is planned to be launched in Japanese Fiscal Year (JFY) 2016 (from April 2016 to March 2017). SGLI has middle spatial resolution (250 m to 1000 m), wide swath (1150 km to 1400 km), 19 bands from near-UV (380 nm) to thermal infrared (12  $\mu$ m) wavelengths, and two-channel (red and near infrared) slant view polarization observations. SGLI will provide several ocean color products including normalized water-leaving radiance (*NWLR*) (or remote sensing reflectance ( $R_{rs}$ )), photosynthetically available radiation (*PAR*), chlorophyll-*a* concentration (*Chla*), colored dissolved organic matter (*CDOM*), total suspended matter concentration (*TSM*), which will contribute to coastal environment monitoring and climate researches by the SGLI 250m resolution and wide swath. *Chla* is a key parameter to know phytoplankton distribution and the ocean primary production. Traditionally, it was estimated by an empirical regression between *Chla* and blue/green ratio of  $R_{rs}$  (e.g., OC4 algorithm (O'Reilly et al., 2000)). The regression is basing on a global in-situ dataset (e.g., NASA bio-Optical Marine Algorithm Data set, NOMAD (Werdell and Bailey, 2005)). However, the relationship can be deviated due to anomalous condition of inherent optical properties (IOPs), phytoplankton absorption,  $a_{ph}$ , CDOM + detritus absorption,  $a_{dg}$ , and particle back-scattering,  $b_{bp}$ , especially in the coastal areas.

This study showed improvement of the *Chla* estimation by considering the IOP deviation through a simple IOP models (Gordon et al., 1988 and Lee et al., 2002). We tested the scheme for in-situ  $R_{rs}$  and *Chla* data observed by Seikai National Fisheries Research Institute (SNFRI) in the East China Sea, which is independent of the NOMAD dataset. Firstly, we calculated  $Chla^{1st}$  by the traditional OC4 algorithm and  $a_{ph}$  by the linear matrix inversion scheme (Hoge and Lyon, 1996, 1999) from the observed  $R_{rs}$ . Then,  $R_{rs}$  is modified by the IOP model with the estimated  $a_{ph}$ , which is assumed to be strongly related to *Chla*, and average state of  $a_{dg}$  and  $b_{bp}$  at condition of the *Chla* value. The average state of  $a_{dg}$  and  $b_{bp}$  was modeled by regression with *Chla* basing on the NOMAD dataset in advance. Finally we recalculated  $Chla^{re}$  by the OC4 algorithm applied to the modified  $R_{rs}$ . Mean absolute difference (MAD) compared to the in-situ observed *Chla* was improved from 50% ( $Chla^{1st}$ ) to 40% ( $Chla^{re}$ ).

This scheme assumed spectral shape of  $a_{ph}$ ,  $a_{dg}$  and  $b_{bp}$ , however they can change in various coastal environment. Collection of the in-situ bio-optical measurements in the various coastal areas is required to develop more robust GCOM-C algorithms and methodology to estimate coastal *Chla*.

キーワード：GCOM、GCOM-C、SGLI、海色、クロロフィルa濃度

Keywords: GCOM, GCOM-C, SGLI, ocean color, chlorophyll-a concentration