

GOSATによる太陽光誘導クロロフィル蛍光と大気CO<sub>2</sub>濃度の同時観測

Simultaneous observations of solar-induced chlorophyll fluorescence by vegetation and atmospheric CO<sub>2</sub> dynamics by GOSAT

\*野田 響<sup>1</sup>、彦坂 幸毅<sup>2</sup>、村上 和隆<sup>1</sup>、松永 恒雄<sup>1</sup>

\*Hibiki M Noda<sup>1</sup>, Kouki Hikosaka<sup>2</sup>, Kazutaka Murakami<sup>1</sup>, Tsuneo Matsunaga<sup>1</sup>

1.国立研究開発法人国立環境研究所、2.東北大学

1.NIES National Institute of Environmental Studies, 2.Tohoku University

In these decades, global warming has progressed owing to increase of greenhouse gases (GHGs) such as CO<sub>2</sub>. To deal effectively with this issue by mitigation and adaptation, it is necessary to monitor emission and sequestration of GHGs with their underlying mechanisms including biogeochemical processes and human activities. Terrestrial ecosystem, which is the large carbon sink, absorbs 123 Pg carbon per year through plant photosynthesis (IPCC 2014). Satellite remote sensing has been used to monitor the spatial and temporal dynamics of terrestrial ecosystems that are responsible for such photosynthetic CO<sub>2</sub> absorption. Such observation provides us with geographical information on the potential distribution of carbon sequestration by the aid of ecosystem models. However, as the photosynthesis of a given vegetation is quite sensitive to meteorological changes such as radiation, temperature and precipitation, we need to observe the photosynthetic 'activity' in a physiological sense, together with the atmospheric CO<sub>2</sub> concentration over continental and global scales. Joiner et al. (2011) and Frankenberg et al. (2011) have suggested that TANSO FTS on Greenhouse Gases Observing Satellite (GOSAT) could detect overlapping part of solar-induced chlorophyll fluorescence (SIF) emitted by terrestrial vegetation and Fraunhofer line. The chlorophyll fluorescence is photons of red and far-red light that emitted by chlorophylls, and in plant ecophysiology it has been a biophysical index to examine the photosynthetic responses to environmental stresses such as extreme temperatures and drought. Thus SIF remote sensing is drawn considerable attention as a new technique to observe the photosynthetic activity of the vegetation. This paper will present our on-going and future challenges by GOSAT and GOSAT-2 to observe such photosynthetic activity of terrestrial ecosystems and its possible consequences with the atmospheric CO<sub>2</sub> concentration from national, continental to global scales under climate change.

キーワード：陸域炭素循環、光合成生産

Keywords: carbon cycle of terrestrial ecosystem, photosynthetic production