

長期的な熱負荷および自然放冷過程における地下水質変化

Change in groundwater quality during the long term thermal loading and cooling

* 鈞持 将人¹、斎藤 健志¹、川本 健¹、小松 登志子¹

* Masato Kenmochi¹, Takeshi Saito¹, Ken Kawamoto¹, Toshiko Komatsu¹

1. 埼玉大学大学院理工学研究科

1. Graduate School of Science and engineering, Saitama University

Recently, the subsurface temperature increase due to global warming, heat island phenomenon, the waste heat from the underground structures and some other causes has been observed worldwide. Increase in subsurface temperature might affect groundwater quality (e.g., dissolution of toxic substances such as heavy metals in the soil, generation of metabolic substances by the change of microbial activity). But its effects are not well understood. Therefore, the purpose of this study is to investigate the effects of subsurface temperature increase on groundwater quality. In this study, the subsurface heat exchanger installed in the experimental site at Saitama University campus and water of 40 °C and 60 °C was circulated in the heat exchanger for the thermal loading test. The groundwater quality was measured at four monitoring wells with two aquifers. As a result of thermal loading, the concentration of several components (K, Na, NH₃, B, Li, Si, As) increased with subsurface temperature increase, while the concentration of a few components (Mg, Ca) decreased. These results show that the subsurface temperature increase affects groundwater quality.

キーワード：地下水質、地下温度、熱負荷

Keywords: groundwater quality, subsurface temperature, thermal loading