

深海底の凹凸地形上に形成される乱流ホットスポットの鉛直構造に関する数値的研究
Numerical Study of the Impacts of Ocean Bottom Roughness and Tidal Flow Amplitude on
Abyssal Mixing

*日比谷 紀之¹、伊地知 敬¹、ロバートソン ロビン²

*Toshiyuki Hibiya¹, Takashi Ijichi¹, Robin Robertson²

1.東京大学 大学院理学系研究科 地球惑星科学専攻、2.ニューサウスウェールズ大学 物理・環境・数理科学専攻、キャンベラ、オーストラリア

1.Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, Tokyo, Japan, 2.School of Physical, Environmental and Mathematical Sciences, The University of New South Wales, Canberra, Australia

Although an accurate representation of ocean mixing processes in global circulation models is essential for accurate climate predictions, parameterization of mixing over rough bathymetry remains uncertain. We perform here a series of eikonal calculations for a wide range of physical parameters to investigate the transfer of energy from upward propagating internal waves generated by tide-topography interactions to dissipation through nonlinear interaction with background three-dimensional Garrett-Munk-like internal waves.

Following the previous study (Mohri et al., 2010) and using a fixed N value, we assume that internal waves generated by tide-topography interactions can be classified into linear internal tides when $kU_0/\omega < 1$ and quasi-steady lee waves when $kU_0/\omega > 1$, where U_0 is the tidal flow amplitude, k the benthic bathymetric wavenumber, N the buoyancy frequency and ω the semidiurnal tidal frequency. Of special note is that the vertical group velocity C_{gz} is inversely proportional to k for linear internal tides and proportional to kU_0^2 for quasi-steady lee waves, although the resonant interaction time is roughly inversely proportional to k for both cases. As a result, the resulting mixing hotspot becomes more restricted to the ocean bottom as bottom roughness increases for $kU_0/\omega < 1$, independent of the tidal flow amplitude, but it extends upward as the tidal flow amplitude increases for $kU_0/\omega > 1$, independent of the bottom roughness. In both cases, we can find a trade-off relationship between the energy dissipation rate at the ocean bottom and its vertical extent.

The accuracy of global circulation models will be improved by reflecting these results in the parameterization of mixing over rough bathymetry.

キーワード：深海混合、パラメタリゼーション、深層海洋循環、深海凹凸地形、潮汐流

Keywords: Abyssal Mixing, Parameterization, Global Overturning Circulation, Ocean Bottom Roughness, Tidal Flow