

グリーン関数法による太平洋深層の鉛直拡散の最適化

An improved simulation of the deep Pacific Ocean using optimally estimated vertical diffusivity based on the Green's function method

*豊田 隆寛¹、杉浦 望²、佐々木 祐二、五十嵐 弘道²、石川 洋一²、畠山 隆紀²、河野 健²、川合 義美²、瀬戸 慎也²、勝俣 勝郎²、内田 裕²、土居 知将²、深澤 理郎²、淡路 敏之³

*Takahiro Toyoda¹, Nozomi Sugiura², Yuju Sasaki, Hiromichi Igarashi², Yoichi Ishikawa², Takaki Hatayama², Takeshi Kawano², Yoshimi Kawai², Shinya Kouketsu², Katsuro Katsumata², Hiroshi Uchida², Toshimasa Doi², Masao Fukasawa², Toshiyuki Awaji³

1.気象庁気象研究所、2.海洋研究開発機構、3.京都大学

1.Meteorological Research Institute, Japan Meteorological Agency, 2.Japan Agency for Marine-Earth Science and Technology, 3.Kyoto University

An improved vertical diffusivity scheme is introduced into an ocean general circulation model to better reproduce the observed features of water property distribution inherent in the deep Pacific Ocean structure. The scheme incorporates (a) a horizontally uniform background profile, (b) a parameterization depending on the local static stability, and (c) a parameterization depending on the bottom topography. Weighting factors for these parameterizations are optimally estimated based on the Green's function method. The optimized values indicate an important role of both the intense vertical diffusivity near rough topography and the background vertical diffusivity. This is consistent with recent reports that indicate the presence of significant vertical mixing associated with finite-amplitude internal wave breaking along the bottom slope and its remote effect. The robust simulation with less artificial trend of water properties in the deep Pacific Ocean illustrates that our approach offers a better modeling analysis for the deep ocean variability. This presentation is based on Toyoda et al. (2015) published in Geophysical Research Letter (42, 9916-9924, doi:10.1002/2015GL065940).

キーワード：鉛直拡散、データ同化、太平洋深層循環

Keywords: vertical diffusivity, data assimilation, Pacific Ocean