

高波長分解能赤外ヘテロダイン分光器を用いた火星大気中のメタン及び水蒸気同位体の検出  
Methane and HD0/H2O in the Martian atmosphere studied by ultra-high spectral resolution

\*中川 広務<sup>1</sup>、青木 翔平<sup>3</sup>、佐川 英夫<sup>4</sup>、高見 康介<sup>1</sup>、笠羽 康正<sup>1</sup>、村田 功<sup>1</sup>、Sonnabend Guido<sup>6</sup>、Manuela Sornig<sup>7</sup>、坂野井 健<sup>2</sup>、鍵谷 将人<sup>2</sup>、Jeffrey Kuhn<sup>8</sup>、Joseph Ritter<sup>8</sup>、岡野 章一<sup>2</sup>、田口 真<sup>5</sup>  
\*Hiromu Nakagawa<sup>1</sup>、Shohei Aoki<sup>3</sup>、Hideo Sagawa<sup>4</sup>、Kosuke Takami<sup>1</sup>、Yasumasa Kasaba<sup>1</sup>、Isao Murata<sup>1</sup>、Guido Sonnabend<sup>6</sup>、Manuela Sornig<sup>7</sup>、Takeshi Sakanoi<sup>2</sup>、Masato Kagitani<sup>2</sup>、Kuhn Jeffrey<sup>8</sup>、Ritter Joseph<sup>8</sup>、Shoichi Okano<sup>2</sup>、Makoto Taguchi<sup>5</sup>

1.東北大学 大学院理学研究科 地球物理学専攻太陽惑星空間物理学講座 惑星大気物理学分野、2.東北大学 大学院理学研究科 惑星プラズマ大気研究センター、3.Istituto di Astrofisica e Planetologia Spaziali, Istituto Nazionale di Astrofisica、4.京都産業大学理学部、5.立教大学、6.Radiometer Physics GmbH、7.German Aerospace Center、8.Institute for Astronomy, University of Hawaii, Advanced Technology Research Center

1.Planetary Atmosphere Physics Laboratory, Department of Geophysics, Graduate School of Science, Tohoku University, 2.Planetary Plasma and Atmospheric Center, Graduate School of Science, Tohoku University, 3.Istituto di Astrofisica e Planetologia Spaziali, Istituto Nazionale di Astrofisica, 4.Faculty of Science, Kyoto Sangyo University, 5.Rikkyo University, 6.Radiometer Physics GmbH, 7.German Aerospace Center, 8.Institute for Astronomy, University of Hawaii, Advanced Technology Research Center

The Mid-Infrared LAsER Heterodyne Instrument (MILAHl), which operates onboard the dedicated Tohoku 60cm telescope (T60) at the summit of Mt. Haleakala, has been designed for investigating the trace gases (Methane, HD0/H2O, etc) in the terrestrial atmospheres, such as Mars and Venus. The limitation to detect such trace gases from the ground-based is mainly due to the difficulty of correcting the atmospheric absorptions in the Earth atmosphere. High spectral resolution of MILAHl (>10E6) enables to retrieve them without any ambiguity due to the reproduction of atmospheric spectra on Earth.

In this study, we focus on the detection of methane and HD0/H2O in the Martian atmosphere. As a local oscillator (LO), newly installed quantum cascade laser (QCL) nicely covers 7.7 micron wavelength for these molecules. It is the only IR heterodyne instrument that gives access to new spectral range as compared with previous instruments of this kind.

Because the facility/instrument is just becoming to be operational in these years, the first Mars campaign will be performed on Feb.-Mar. 2016, with large Doppler shift (~15 km/s) between Mars and Earth. Prediction of the radiative transfer model indicates that the determination with two- VSMOW precision could be obtained by 15-minute integration. Upper limit 100ppb of methane will also be determined by 32-hours integration.

Further continuous observations will help to constrain (i) the possibility of biological/geological activities in the current Martian atmosphere, and (ii) water cycle and its evolution on Mars.

キーワード：メタン、HD0/H2O、ヘテロダイン

Keywords: Methane, HD0/H2O, Heterodyne