日本海東縁深海堆積物中におけるメタノールの生物地球化学的物質循環 Biogeochemical cycle of methanol in anoxic deep-sea sediments of the eastern Japan Sea

*柳川 勝紀¹、谷 篤史²、山本 直弥²、八久保 晶弘³、狩野 彰宏¹、松本 良⁴、鈴木 庸平⁵ *Katsunori Yanagawa¹, Atsushi Tani², Naoya Yamamoto², Akihiro Hachikubo³, Akihiro Kano¹, Ryo Matsumoto⁴, Yohey Suzuki⁵

 1.九州大学大学院比較社会文化研究院、2.大阪大学大学院理学研究科、3.北見工業大学 環境・エネルギー研 究推進センター、4.明治大学 ガスハイドレート研究所、5.東京大学大学院理学系研究科
1.Graduate School of Social and Cultural Studies, Kyushu University, 2.Department of Earth and Space Science, Graduate School of Science, Osaka University, 3.Environmental and Energy Resources Research Center, Kitami Institute of Technology, 4.Gas Hydrate Laboratory, Meiji University, 5.Department of Earth and Planetary Science, University of Tokyo

Methanol is one of the most important carbon and energy sources in anoxic environments. However, the biological flux and lifetime of methanol in anoxic marine sediments are largely unknown. In this study, we report quantitative methanol removal rates in subsurface sediments for the first time. Methanol concentrations in pore water from Japan Sea sediments gradually increased with depth below the sulfate-methane transition zone. Based on anaerobic incubation experiments with radiotracers, high rates of microbial methanol consumption were detected in the sediments. Our experiments also showed that the methanol oxidation to CO_2 surpassed methanol assimilation and methanogenesis from CO_2/H_2 and methanol. Nonetheless, a significant decrease in methanol was not observed after incubation, likely because of the microbial production of methanol in parallel with its consumption. This study suggests that microbial reactions play an important role in the sources and sinks of methanol in subseafloor sediments.

キーワード:メタノール、深海堆積物、海底下生命圏 Keywords: methanol, deep-sea sediment, subseafloor biosphere