深海堆積物中に記録された真核生物DNAに堆積物酸化還元状態が与える影響

Effects of sedimentary redox conditions on Eukaryotic DNA recorded in deep-sea sediments

*山口 保彦¹、幸塚 麻里子¹、山崎 俊嗣²、川村 紀子³、清家 弘治²、櫻本 晋洋²、奥津 なつみ²、鈴木 庸平¹ *Yasuhiko T. Yamaguchi¹, Mariko Kouduka¹, Toshitsugu Yamazaki², Noriko Kawamura³, Koji Seike², Yukihiro SAKURAMOTO², Natsumi Okutsu², Yohey Suzuki¹

 1.東京大学大学院理学系研究科地球惑星科学専攻、2.東京大学大気海洋研究所、3.海上保安大学校
1.Department of Earth and Planetary Science, The University of Tokyo, 2.Atmosphere and Ocean Research Institute, The University of Tokyo, 3.Japan Coast Guard Academy

Eukaryotic DNA in marine sediments can be a useful indicator of both ancient marine ecosystem in water column and living microbial eukaryotes in sediments. However, the environmental factors that influence the composition and preservation of eukaryotic DNA in marine sediments are poorly characterized. In this study, we examined effects of sedimentary redox conditions on the diversity of eukaryotic communities recorded in deep-sea surface sediment samples from 8 sites of the Japan Sea (from oxygenated abyssal sediments to sulfidic sediments in methane seeps) using a combination of various geochemical and molecular-biological tools. Sedimentary redox conditions were characterized by depth profiles of pore water (oxygen, nitrate, iron, sulfide, etc.) and bulk sedimentary organic matter (TOC, TN, δ^{13} C, δ^{15} N). The concentration and diversity of 18S rDNA in the sediment samples were investigated by qPCR and pyrosequencing. Decreases in 18S rDNA sequences also varied with the sedimentary redox conditions. These results suggest that redox conditions of surface sediments can be important factors controlling the composition and preservation of eukaryotic DNA in deep-sea sediments.

キーワード:酸化還元状態、海洋堆積物、生物多様性、DNA、真核生物 Keywords: Redox condition, Marine sediments, Biodiversity, DNA, Eukaryote