The Effect of Forest Management of Secondary Coniferous forests on User's Landscape Appreciation and Psychological Restorativeness

*Norimasa TAKAYAMA¹, Haruo SAITO², Akio FUJIWARA², Masahiro HORIUCHI³

1.Forestry and Forest Products Research Institute in Japan, 2.Fuji Iyashinomori Woodland Study Center, The University of Tokyo Forests, 3.Mount Fuji Research Institute, Yamanashi Prefectural Government

-*INTRODUCTION*: We investigated the influence of forest management on landscape appreciation and the psychological restorative effect in an on-site setting by exposing respondents to an unmanaged coniferous forest (U.F.), and a managed coniferous forest (M.F.) for a particular period. The both forests, which consisted of Japanese larch and Japanese red pine (a second-growth forest), were fairly similar in the land cover type and vegetation one another.

-METHODS: We considered the experiment in late July. We set the two experimental plots (0.25 ha) in the both forests of Fuji Iyashinomoroi Woodland Study Center as U.F. setting and M.F. setting. Here, the mean temperature, relative humidity and sound pressure were almost the same during the experiment except illuminance. The respondents were eighteen individuals (eighteen males; aged twenties to fifties) for the experiment. As for eliminating an order effect, the respondents were divided into the two groups (Group A and Group B) in every nine-person. The respondents of Group A were exposed to U.F. setting at first and then were done to M.F. setting. However, the respondents of Group B were exposed to each setting by the opposite order. They were individually exposed to the both settings while sitting for 15 min. In the both settings, the respondents were required to answer the three questionnaires to investigate the psychological restorative effect at before and after the experiment (mood; POMS, affect; PANAS, subjective restorativeness; ROS). For comparison of landscape appreciation, the respondents were required to answer other two questionnaires at after the experiment (scene appreciation (SD), a restorative property of environment (PRS)). -RESULTS: As a comparison result by the statistical test, regarding a restorative property of environment (PRS), M.F. setting had statistically higher property in "Being away" and "Coherence", "Compatibility" than U.F. setting (p< .05). About scene appreciation (SD), M.F. were appreciated statistically higher in "brightness," "openness," "comfort," "beauty," "safeness" and "healthiness" (p< .05), and "order" and "thin" (p< .01). On the other hands, by the result of two-way repeated ANOVA (difference of setting (U.F. -M.F.) xpresence of experience (before exposure -after exposure)), there were no statistical relationship with the mutual interaction between difference of setting and presence of experience in "mood" (POMS), "affect" (PANAS) and "subjective restorativeness" (ROS).

Then, as a result of having checked both the main effects, the difference of setting did not seem to raise a psychological restorativeness. Otherwise, the presence of experience could give a statistical influence negative "affect" (PANAS; p< .05) and "tension and anxiety" (POMS; p< .05). The difference of setting also reduced numerical values for them in M.F. setting. In contrast, before and after exposure could give a statistical influence and raise "vigor" in U.F. setting (POMS; p< .05).

-CONSSIDERATION: Consequently, negative affect, tension, and anxiety might come to decrease because the managed forest setting had a sufficient restorative property of the environment and the better scenic environment. Conclusively, respondents would obtain a psychological restorativeness to some extent by being exposed to M.F. setting. On the other hand, even though vigor rose in U.F. setting, we would consider the reason for it by these three hypotheses as follows;

1) all the respondents were men. 2) the sample group had a tendency toward a relatively low

neuroticism and a high extroversion by the personality traits test which we also conducted as one of the optional tests. 3) if we referred to the Kaplan's landscape preference theory, we could think of the possibility that U.F. setting would bring a sense of mystery and exploration to the respondents who had the trait mentioned above.

Keywords: Landscape appreciation, Psychological restorative effect, Forest management, Coniferous forest, Subjective restorativeness

| Semantic differential official name Semantic differential method Perceived restorativeness Scale Profile of mood states Positive and negative affect schedule Restorative outcome schedule contents scene appreciation restorative property of mood affect subjective | category | | Landscape appreciation | | Psychological restorativeness | | |
|---|---------------------------|-------------|--|---|-------------------------------|--------------------|-----------------------------|
| official name method restorativeness Scale Profile of mood states negative affect subjective schedule contents scene appreciation restorative property of environment mood affect subjective restorative property of environment number of subscales 25 5 6 2 1 timing of the measurement after exposure before and after exposure 1 wilcoxon signed rank test "beauty", "safeness", "order" and "thin" than UF. (p<01 to p< 05). "Compatibility" than U.F. (p<01 to p< 05). two-way repeated ANOVA mutual interaction main effect mutual interaction main effect n.s. n.s. n.s. n.s. | abbreviated form | | SD | PRS | POMS | PANAS | ROS |
| contents environment restorativ number of subscales 25 5 6 2 1 timing of the measurement after exposure before and after exposure before and after exposure wilcoxon signed rank test "openness", "comfort", "Coherence" and "beauty", "safeness", "Compatibility" than U.F. "healthiness", "Openness", "compatibility" than U.F. (p<.05). | official name | | | | Profile of mood states | negative affect | Restorative outcome sc |
| timing of the measurement after exposure before and after exposure M.F. was statistically M.F. was statistically M.F. was statistically wilcoxon signed rank test "openness", "comfort", "Coherence" and "Compatibility" than U.F. "healthiness", "order" (p<.05). | contents | | scene appreciation | | mood | affect | subjective restorativeni |
| M.F was statistically higher in "brightness", "openness", "compress", "Compatibility" than U.F. "healthiness", "order" and "thin" than U.F. (p<.01 to p<.05). | number of subscales | | 25 | 5 | 6 | 2 | 1 |
| higher in "brightness", "openness", "comfort", "Coherence" and "beauty", "safeness", "compatibility" than U.F. "Compatibility" than U.F. "beauty", "safeness", "order" and "thin" than U.F. "Compatibility" than U.F. (p<.01 to p<.05). | timing of the measurement | | after e | exposure | befor | e and after exposu | re |
| two-way repeated ANOVA n.s. n.s. n.s. interaction U.F.: vigor (p<.05)1 | wilcoxon signed rank test | | higher in "brightness", "openness", "comfort", "beauty", "safeness", "healthiness", "order" and "thin" than U.F. | higher in "Being away", "Coherence" and "Compatibility" than U.F. | | | |
| main effect M.F.: tension and affect (PANAS; p< n.s. | two-way repeated ANOVA | | | | n.s. | n.s. | n.s. |
| | | main effect | - | | M.F.: tension and | affect (PANAS; p | < n.s. |
| | | | | | | | |

U. F.: unmanaged forest, M. F.: managed forest, 1 (increased, 1) (decreased