

カナダ永久凍土における植物の窒素吸収様式に対する環境要因の影響

Environmental controls on patterns of plant nitrogen uptake in Canadian permafrost soils

*藤井 一至¹、稲垣 善之¹、松浦 陽次郎¹、大澤 晃²*Kazumichi Fujii¹, Yoshiyuki Inagaki¹, Matsuura Yojiro¹, Akira Osawa²

1.森林総合研究所、2.京都大学

1.Forestry and Forest Products Research Institute, 2.Kyoto University

Plants develop different strategies of nitrogen (N) uptake in response to N limitation. This may explain N supply to plants from N-limited soils and niche differentiation due to soil N levels. To analyze the relationships between N availability and patterns of plant N uptake, plant uptake rates of different N sources (¹³C, ¹⁵N-glutamic acid, ¹⁵N-urea, ¹⁵N-ammonium, ¹⁵N-nitrate) were measured in the field condition. To assess N availability, the concentrations of amino acids, ammonium, and nitrate in soil solution and the mineralization rates of glutamic acid, arginine, ornithine, and urea were studied. We compared four sites in Northwest Territories, Canada; white spruce forest (WSF) at upper slope and black spruce forest (BSS) at lower slope on foothill of glaciofluvial sands, black spruce forest (BSF) and shrub birch tundra (TND) on fluvial sediments. In soil solution, DON was abundant at all sites, but inorganic N was abundant only in TND and BSS. Urea mineralization potentials were consistently high at all sites, except for BSF. This suggests microbial N mineralization is highly limited in N-limited BSF soil. Birch and grass preferentially absorbed inorganic N, while white spruce and black spruce can utilize amino acid-N under N-limited soil to bypass N flow between soil and plants.

キーワード：土壌、永久凍土、アミノ酸

Keywords: Soil, Permafrost soil, Amino acid