

風成塵起源粒子の光ルミネッセンス年代測定に基づく過去12万年間の日本海盆および大和堆における堆積速度の変化

Optically stimulated luminescence dating back to 100ka and sedimentation rates of dark versus light layers in the Japan Basin

*杉崎 彩子¹、Murray Andrew²、Buylaert Jan-Pieter^{2,3}、多田 隆治⁴、池原 研¹、長橋 良隆⁵、里口 保文⁶、入野 智久⁷、Murray Richard⁸、Alvarez-Zarikian Carlos⁹、Expedition 346 scientists

*Saiko Sugisaki¹、Andrew Murray²、Jan-Pieter Buylaert^{2,3}、Ryuji Tada⁴、Ken Ikehara¹、Yoshitaka Nagahashi⁵、Yasufumi Satoguchi⁶、Tomohisa Irino⁷、Richard W Murray⁸、Carlos A Alvarez-Zarikian⁹、Expedition 346 scientists

1.産業技術総合研究所、2.Aarhus University、3.Technical University of Denmark、4.東京大学、5.福島大学、6.滋賀県立琵琶湖博物館、7.北海道大学、8.Boston University、9.Texas A&M University

1.Geological Survey of Japan, AIST, 2.Aarhus University, 3.Technical University of Denmark, 4.The University of Tokyo, 5.Fukushima University, 6.Lake Biwa Museum, 7.Hokkaido University, 8.Boston University, 9.Texas A&M University

Sediments in the Japan and Yamato Basins record millennial-scale changes in the East Asian Monsoon as alternations of dark and light layers. The gray-scale profile showing this millennial-scale variability has been correlated with the $d^{18}O$ record from Greenland ice cores (GISP2). However, this correlation between millennial and orbital scale changes becomes less certain during the last interglacial and the Holocene period, largely as a result of a paucity of biogenic carbonate used for ^{14}C and isotope stratigraphy; there is significant age uncertainty over this time period. We have used optically stimulated luminescence (OSL) dating of fine-grained detrital quartz (that are aeolian in origin) to provide high-resolution numerical age models for cores U1424C and U1425C drilled in the southeastern margin of the Japan Basin, and the Yamato Rise, during IODP Expedition 346.

We report our attempts to: 1) assess whether surface sediment of the cores are consistent with modern age; 2) test the reliability of OSL ages by comparison with tephra chronology and orbitally-tuned age models based on the correlation between the GRA record and the LR04 stack; 3) date cores U1424C and U1425C to construct a robust 120 ky age model with a resolution of 10 ky; and 4) conduct high-resolution dating of several pairs of dark and light layers to estimate short term changes in sedimentation rate, in particular over the last ~30 ky. We show OSL dating of fine aeolian silt in Japan Sea sediments shows great potential to establish independent numerical chronologies, and this is especially useful in cores where isotope stratigraphies and/or gray-scale signatures are poorly reconciled.

キーワード：光ルミネッセンス年代測定、IODP Expedition 346、石英

Keywords: Optically stimulated luminescence dating , IODP Expedition 346, quartz