

南太平洋環流域の遠洋性粘土に存在する大量の微小マンガングラン粒について

Abundant manganese microparticles in oxic pelagic clay of the South Pacific Gyre

*浦本 豪一郎¹、諸野 祐樹¹、富岡 尚敬¹、和穎 朗太²、上杉 健太郎³、竹内 晃久³、星野 真人³、鈴木 芳生^{3,4}、光延 聖⁵、菅 大暉⁶、宮本 千尋⁴、高橋 嘉夫⁴、稲垣 史生¹

*Uramoto Go-Ichiro¹, Yuki Morono¹, Naotaka Tomioka¹, Rota Wagai², Kentaro Uesugi³, Akihisa Takeuchi³, Masato Hoshino³, Yoshio Suzuki^{3,4}, Satoshi Mitsunobu⁵, Hiroki Suga⁶, Chihiro Miyamoto⁴, Yoshio Takahashi⁴, Fumio Inagaki¹

1.国立研究開発法人海洋研究開発機構、2.国立研究開発法人農業環境技術研究所、3.公益財団法人高輝度光科学研究センター、4.東京大学、5.愛媛大学、6.広島大学

1.Japan Agency for Marine-Earth Science and Technology, 2.National Institute for Agro-Environmental Sciences, 3.Japan Synchrotron Radiation Research Institute, 4.University of Tokyo, 5.Ehime University, 6.Hiroshima University

Manganese nodules ubiquitously occur in the abyssal plain, with varying concentrations of metals, including manganese, iron, nickel, cobalt, and copper. Typical manganese nodules observed in pelagic sediments are in centimeter to sub-millimeter size of spherical crusts and often concreted to the pavement over the seafloor. During the Integrated Ocean Drilling Program (IODP) Expedition 329 in 2010, we drilled the entire sedimentary sequence at 6 sites in the ultra-oligotrophic region of the South Pacific Gyre (SPG), where dissolved O₂ and aerobic microbial communities are present from the seafloor to the sediment-basement interface [1]. Massive manganese nodules often occur at the seafloor of these drilling sites, and the subseafloor sediments generally consist of zeolitic metalliferous clay. Using a newly developed sediment observation technique [2], we observed numerous micrometer-scale particles of (ferro-)manganese minerals in the oxic zeolitic clay from present-day to several tens of millions of years. Using a synchrotron-based X-ray microtomography and FIB-SEM-EDS, high-resolution three-dimensional micro-textures of manganese microparticles, as well as elemental compositions, were visualized, suggesting that those particles were deposited from the water column, and well preserved in the widespread area of deep-sea oxic sediments over geologic time.

[1] D'Hondt, S., Inagaki, F., Zarikian, C. A., Abrams, L. J., Dubois, N., Engelhardt, T., Evans, H., Ferdelman, T., Gribsholt, B., Harris, R. N., Hoppie, B. W., Hyun, J.-H., Kallmeyer, J., Kim, J., Lynch, J. E., McKinley, C. C., Mitsunobu, S., Morono, Y., Murray, R. W., Pockalny, R., Sauvage, J., Shimono, T., Shiraishi, F., Smith, D. C., Smith-Duque, C., Spivack, A. J., Steinsbu, B. O., Suzuki, Y., Szpak, M., Toffin, L., Uramoto, G., Yamaguchi, T. Y., Zhang, G., Zhang, X.-H., and Ziebis, W. Presence of oxygen and aerobic communities from seafloor to basement in deep-sea sediment. *Nature Geosciences*, 8(4), 299-304, 2015.

[2] Uramoto, G.-I., Morono, Y., Uematsu, K., and Inagaki, F. An improved sample preparation method for imaging microstructures of fine-grained marine sediment using microfocuss X-ray computed tomography and scanning electron microscopy. *Limnology and Oceanography: Methods*, 12, 469-483, 2014.

キーワード：マンガングラン、微小パーティクル、遠洋性粘土、深海平原、南太平洋環流域

Keywords: manganese, microparticles, pelagic clay, abyssal plain, South Pacific Gyre