

共鳴散乱ライダーで得られた中間圏下部熱圏における大気温度の時間-高度変動特性 (36°N, 140°E)
Time and height variability of temperature in Mesosphere and Lower Thermosphere region
based on resonance scattering lidar measurement at NIPR (36°N, 140°E)

*西山 尚典^{1,2}、江尻 省^{1,2}、津田 卓雄³、津野 克彦^{1,6}、阿保 真⁴、川原 琢也⁵、小川 貴代⁶、和田 智之⁶、中村 卓司^{1,2}

*Takanori Nishiyama^{1,2}, Mitsumu K. Ejiri^{1,2}, Takuo T. Tsuda³, Katsuhiko Tsuno^{1,6}, Makoto Abo⁴, Takuya Kawahara⁵, Takayo Ogawa⁶, Satoshi Wada⁶, Takuji Nakamura^{1,2}

1.国立極地研究所、2.総合研究大学院大学、3.電気通信大学、4.首都大学東京大学院システムデザイン研究科、5.信州大学工学部、6.理化学研究所

1.National Institute of Polar Research, 2.SOKENDAI (The Graduate University for Advanced Studies), 3.The University of Electro-Communications, 4.Graduate School of System Design, Tokyo Metropolitan University, 5.Faculty of Engineering, Shinshu University, 6.RIKEN

The National Institute of Polar Research (NIPR) is leading a six year prioritized project of the Antarctic research observations since 2010. One of the sub-projects is entitled 'the global environmental change revealed through the Antarctic middle and upper atmosphere'. As a part of the sub-project, a Rayleigh/Raman lidar (RR lidar) was installed at Syowa, Antarctica (69S, 39E) in January, 2011. The operation has been conducted since February 2011 and the RR lidar has kept measuring temperature profiles continuously between approximately 10 and 80 km for almost 3 years. In order to extend the height coverage to include mesosphere and lower thermosphere region, a new resonance scattering lidar system with tunable wavelengths is developed at NIPR in Tachikawa (35.7N, 139.4E).

The lidar transmitter is based on injection-seeded, pulsed alexandrite laser for 768-788 nm (fundamental wavelengths) and a second-harmonic generation (SHG) unit for 384-394 nm (second harmonic wavelengths). The laser wavelengths are tuned into the resonance wavelengths by a wavemeter that is calibrated and validated using a wavelength-stabilized He-Ne laser and a potassium vapor cell for doppler-free spectroscopy. This lidar has capabilities to measure density variations of minor constituents such as atomic iron (Fe, 386 nm), atomic potassium (K, 770 nm), calcium ion (Ca⁺, 393 nm), and nitrogen ion (N₂⁺, 390, 391 nm) and temperature profiles in the mesosphere and lower thermosphere (MLT) region. It can also estimate temperature profiles from the upper Stratosphere to the lower mesosphere using signals of Rayleigh scattering.

In this presentation, we will present time and height variability of temperature in the MLT region based on campaign observation in winter 2015-2016 focusing on Sudden Stratospheric Warming (SSW) impact on dynamics in the MLT region. In addition, the obtained temperature profiles are validated by comparisons to those obtained from satellites data such as Aura/MLS. In addition, dynamical and/or chemical response to SSW and sporadic E-layer in MLT region are discussed using neutral Fe atom density data.

キーワード：中間圏-下部熱圏、大気温度、成層圏突然昇温

Keywords: the Mesosphere and Lower Thermosphere, Temperature, Sudden Stratospheric Warming