

## TIE-GCMモデルと地上GPS-TEC観測による磁気嵐時の電離層データ同化

## Ionospheric data assimilation with TIE-GCM and GPS-TEC during geomagnetic storm period

\*陳 佳宏<sup>1</sup>、Lin Charles<sup>1</sup>、Matsuo Tomoko<sup>3,4</sup>、Chen Wei-han<sup>1</sup>、Lee I. T.<sup>2</sup>、劉 正彦<sup>5,6</sup>、林 佳廷<sup>1</sup>

\*Chia-Hung Chen<sup>1</sup>, Charles Lin<sup>1</sup>, Tomoko Matsuo<sup>3,4</sup>, Wei-han Chen<sup>1</sup>, I. T. Lee<sup>2</sup>, Jann-Yenq Liu<sup>5,6</sup>, Jia-Ting Lin<sup>1</sup>

1.台湾国立成功大学地球科学学科、2.台湾気象庁、3.アメリカコロラド大学環境科学、4.アメリカ宇宙天気センター、5.台湾国立中央大学太空科学学科、6.台湾国立中央大学リモートセンシングセンター

1.Department of Earth Sciences, National Cheng Kung University, Tainan, Taiwan, 2.Central Weather Bureau, Taipei, Taiwan, 3.Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, Colorado, USA, 4.Space Weather Prediction Center, National Oceanic and Atmospheric Administration, Boulder, Colorado, USA, 5.Institute of Space Science, National Central University, Chung-Li, Taiwan, 6.Center for Space and Remote Sensing Research, National Central University, Chung-Li, Taiwan

The main purpose of this study is to investigate the latency time for the ionosphere data assimilation during the geomagnetic storm. An Ensemble Kalman Filter (EnKF) module developed by National Center for Atmospheric Research (NCAR), called as Data Assimilation Research Testbed (DART), is applied to assimilate the ionospheric electron density into a theoretical model (Thermosphere-Ionosphere-Electrodynamics General Circulation Model, TIE-GCM) with ground-based GPS total electron content (TEC) observations during the 26 September 2011 geomagnetic storm period. Effects of various assimilation time intervals, 60-, 30-, and 10-minute, on the ionospheric forecast responses are examined by their global root-mean-square errors (RMSEs) during the entire storm period. Substantial reduction of RMSEs for 10 minutes assimilation cycle suggests the ionospheric data assimilation system greatly improve the capability of model forecast during the geomagnetic storm period. Further examination shows that the neutral state variables in the assimilation model are the important factor to change the trajectory of model forecasting. However, the assimilation model with neutral state variables still needs the shorter assimilation cycle (10-minute in this study) to restrain overfitting of neutrals and lead to higher forecast accuracy during the geomagnetic storm.

キーワード：電離層データ同化、磁気嵐

Keywords: Ionospheric data assimilation, geomagnetic storm