

昼側磁気圏界面リコネクション起源の電子スケール開放境界層におけるホイッスラーモード波動の励起

Excitation of whistler-mode waves in the electron scale open boundary layer generated by the dayside magnetopause reconnection

*内野 宏俊¹、栗田 怜²、原田 裕己³、町田 忍²

*Hiroto Uchino¹, Satoshi Kurita², Yuki Harada³, Shinobu Machida²

1.京都大学大学院理学研究科地球惑星科学専攻 地球物理学教室 太陽惑星系電磁気学講座、2.名古屋大学宇宙地球環境研究所、3.Space Sciences Laboratory, University of California, Berkeley

1.Solar-Planetary Electromagnetism Laboratory, Department of Geophysics, Faculty of Science, Kyoto University, 2.Institute for Space-Earth Environmental Research, Nagoya University, 3.Space Sciences Laboratory, University of California, Berkeley

The magnetic reconnection at the dayside magnetopause is generally because upstream physical quantities between magnetosheath and magnetosphere are quite different. Kinetic simulations of asymmetric magnetic reconnection produce an electron outflow layer mainly composed of magnetosheath electrons at the magnetosphere side of the separatrix. The simulation results suggest that this electron outflow layer corresponds to the reconnected open magnetic field closest to the magnetosphere. Based on the simulation result and data from the THEMIS probes, we show an observation of whistler mode waves in the electron outflow layer caused by asymmetric magnetic reconnection at the magnetopause. The waves propagated toward the reconnection region, and the linear growth rate of the wave was positive at the resonant velocity due to the electron temperature anisotropy. We suggest that the anisotropy can be originated from lack of the magnetospheric electrons moving anti-reconnection direction at small pitch angles since the magnetic field as a channel of the electrons connects to the magnetosheath region by the reconnection. This study quantitatively clarifies the excitation of the whistler-mode waves in the electron scale open boundary layer at the magnetopause in association with the dayside magnetopause reconnection.

キーワード：昼側磁気圏界面リコネクション、電子スケール開放境界、ホイッスラーモード波動

Keywords: Dayside magnetopause magnetic reconnection, electron scale open boundary, Whistler mode wave