

雷起源EMIC波動解析によるプラズマ圏下部の重イオン組成比推定

Cold heavy ion composition in the lower plasmasphere estimated from lightning induced EMIC waves

*松田 昇也¹、笠原 禎也¹、Kletzing Craig²

*Shoya Matsuda¹, Yoshiya Kasahara¹, Craig Kletzing²

1.金沢大学 総合メディア基盤センター、2.Department of Physics & Astronomy, University of Iowa

1.Information and Media Center, Kanazawa University, 2.Department of Physics & Astronomy, University of Iowa

Ion cyclotron whistler waves are electromagnetic ion cyclotron (EMIC) mode waves induced by lightning discharge. Propagation properties of ion cyclotron whistler waves strongly depend on the local cold heavy-ion composition. Crossover frequency is an important frequency for the ion cyclotron whistlers, which is a function of the ion composition. In this study, we examine the variation in the crossover frequency of heavy ion band ion cyclotron whistler waves observed by the Van Allen probes and the Akebono satellites. We found that the crossover frequencies of the observed events decreased with increasing altitude. This suggests the total heavy-ion composition is high at low altitudes and decreases with increasing altitude around lower plasmasphere. We can determine the composition of three species of ions by measuring two crossover frequencies. We focus on H⁺ band and He⁺ band ion cyclotron whistler waves, and estimate proton-helium ion ratio ($n(\text{He}^+)/n(\text{H}^+)$) and proton-oxygen ion ratio ($n(\text{O}^+)/n(\text{H}^+)$) in the lower plasmasphere. This wave-based approach used in this study can also be a useful means of estimating unknown cold-ion distributions in the inner magnetosphere.

キーワード：イオンサイクロトロンホイッスラ、EMIC波動、重イオン

Keywords: Ion cyclotron whistler, EMIC wave, heavy ion