

高エネルギープラズマとEMICトリガード放射の相互作用のシミュレーション

Simulation of interaction between energetic plasmas and EMIC triggered emissions

*小路 真史¹、大村 善治²

*Masafumi Shoji¹, Yoshiharu Omura²

1.名古屋大学宇宙地球環境研究所、2.京都大学生存圏研究所

1.Institute for Space-Earth Environmental Laboratory, 2.Research Institute for Sustainable
Humanosphere

Electromagnetic ion cyclotron (EMIC) rising-tone (triggered) emission is known as a fine structure of EMIC waves. Theoretical studies suggest that EMIC triggered emissions can induce significant loss of the relativistic electrons through the nonlinear trapping. We perform test particle simulations of the electrons interacting with the EMIC triggered emissions reproduced by real scale hybrid simulation. Relativistic electrons with a few MeV kinetic energy are scattered in wide pitch angle range. The lower energetic electrons are also scattered since the frequency of the triggered emissions reaches near the proton cyclotron frequency and the helium band waves are also induced close to the helium gyro frequency. Broadband EMIC rising waves generated in the small ambient magnetic field gradient model cause rapid precipitation of the relativistic electrons. Simultaneous precipitations of the energetic protons and relativistic electrons are also reproduced in the hybrid and test particle simulations, respectively.

キーワード：EMIC波動、非線形波動粒子相互作用、シミュレーション

Keywords: EMIC waves, nonlinear wave particle interaction, computer simulation