

運動論的アルヴェン波ダイナミクスのためのモーメント分離解法

Moment extracted method for solving kinetic Alfvén wave dynamics

*渡邊 智彦¹、渡邊 裕介¹、前山 伸也¹*Tomo-Hiko Watanabe¹, Yusuke Watanabe¹, Shinya Maeyama¹

1.名古屋大学大学院理学研究科

1. Graduate School of Science, Nagoya University

Kinetic Alfvén waves (KAW), which play crucial roles in a variety of phenomena in space plasmas, involve multiple space- and time-scales. For example, the wavelength along the field line may extend to the system size, while the perpendicular wave numbers are characterized by the ion gyro-radius or the electron skin depth. The characteristic time is given the wave frequency or the electron transit time. Thus, drift kinetic or gyrokinetic simulations of low-frequency plasma dynamics including the KAWs often suffer from a severe Courant condition for explicit time-integrators or a poor convergence of iteration in implicit methods.

To overcome the numerical inefficiency, we have developed a new scheme for solving the KAW dynamics including drift kinetic electrons. In the new scheme, the low-order moments of electron distribution function are calculated separately from the drift kinetic equation for electrons. It enables us to easily implement implicit time-integrators and/or the semi-Lagrangian scheme while keeping the numerical stability and the conservation property. Some applications of the moment extracted formulation will be discussed.

キーワード：アルヴェン波、シミュレーション、ジャイロ運動論

Keywords: Alfvén waves, simulation, gyrokinetics