

太陽系におけるニオブ92の初期存在量と分布

The initial abundance and distribution of ^{92}Nb in the Solar System*飯塚 毅¹、ライ イージェン²、アクラム ワヒード²、アメリン ユーリー³、シオンバクラー マリア²*Tsuayoshi Iizuka¹, Yi-Jen Lai², Waheed Akram², Yuri Amelin³, Maria Schönbaehler²

1.東京大学、2.チューリッヒ工科大学、3.オーストラリア国立大学

1.University of Tokyo, 2.ETH Zürich, 3.Australian National University

Niobium-92 is an extinct proton-rich nuclide, which decays to ^{92}Zr with a half-life of 37 Ma. Because Nb and Zr can fractionate from each other during partial melting of the mantle, mineral crystallization and metal-silicate separation, the Nb-Zr system can potentially be used to determine the timescales of silicate differentiation and core segregation for infant planets. In addition, the initial ^{92}Nb abundance in the Solar System provides constraints on the nucleosynthetic site(s) of p -nuclei (p - denotes proton-rich). These applications require the initial abundance and distribution of ^{92}Nb (expressed as $^{92}\text{Nb}/^{93}\text{Nb}$) in the Solar System to be defined. Yet previously reported initial $^{92}\text{Nb}/^{93}\text{Nb}$ values range from $\sim 10^{-5}$ to $>10^{-3}$ [1-6], and remain to be further constrained. All but one of the previous studies estimated the initial $^{92}\text{Nb}/^{93}\text{Nb}$ using Zr isotope data for single phases with fractionated Nb/Zr in meteorites such as zircons and CAIs, assuming that their source materials and bulk chondrites possessed identical initial $^{92}\text{Nb}/^{93}\text{Nb}$ and Zr isotopic compositions [1-5]. To evaluate the homogeneity of the initial ^{92}Nb abundance, however, it is desirable to define internal mineral isochrons for meteorites with known absolute ages. Although Schönbaehler et al. [6] applied the internal isochron approach to the chondrite Estacado and the mesosiderite Vaca Muerta, these meteorites include components of different origins and their formation ages are uncertain, which prohibits a precise determination of the solar initial ^{92}Nb abundance.

Here we present Nb-Zr data for mineral fractions from four unbrecciated meteorites, which originate from distinct parent bodies and whose U-Pb ages were precisely determined: the angrite NWA 4590, the eucrite Agoult and the ungrouped achondrites Ibitira. Our results show that the relative Nb-Zr isochron ages of the three meteorites are consistent with the time intervals obtained from the Pb-Pb chronometer for pyroxene and plagioclase, indicating that ^{92}Nb was homogeneously distributed among their source regions. The Nb-Zr and Pb-Pb data for NWA 4590 yield the most reliable and precise reference point for anchoring the Nb-Zr chronometer to the absolute timescale: an initial $^{92}\text{Nb}/^{93}\text{Nb}$ ratio of $(1.4 \pm 0.5) \times 10^{-5}$ at 4557.93 ± 0.36 Ma, which corresponds to a $^{92}\text{Nb}/^{93}\text{Nb}$ ratio of $(1.7 \pm 0.6) \times 10^{-5}$ at the time of the Solar System formation. On the basis of this new initial ratio, we demonstrate the capability of the Nb-Zr chronometer to date early Solar System objects including troilite and rutile, such as iron and stony-iron meteorites. Furthermore, we estimate a nucleosynthetic production ratio of ^{92}Nb to the p -nucleus ^{92}Mo between 0.0015 and 0.035. This production ratio, together with the solar abundances of other p -nuclei with similar masses, can be best explained if these light p -nuclei were primarily synthesized by photodisintegration reactions in Type Ia supernovae.

[1] Harper (1996) *ApJ* 466, 437. [2] Sanloup et al. (2000) *EPSL* 184, 75. [3] Yin et al. (2000) *ApJ* 536, L49. [4] Münker et al. (2000) *Science* 289, 1538. [5] Hirata (2001) *Chem. Geol.* 176, 323. [6] Schönbaehler et al. (2002) *Science* 295, 1705.

キーワード：エコンドライト、初期太陽系年代学、元素合成

Keywords: achondrite, early Solar System chronology, nucleosynthesis

