

In-situ X-ray diffraction studies of hydrous phases up to ~50 GPa

*Masayuki Nishi^{1,2}, Tetsuo Irifune^{1,2}, Jun Tsuchiya^{1,2}, Takeshi Arimoto¹, Sho Kakizawa¹, Yoshinori Tange³

1.Geodynamics Research Center, Ehime University, 2.Earth-Life Science Institute, Tokyo Institute of Technology, 3.Japan Synchrotron Radiation Research Institute

Recent experimental and theoretical studies suggest the high-pressure forms of hydrous minerals, phase H and delta- AlOOH , play important roles in transportation of water into the deep Earth's interior via subduction of oceanic slabs. Phase H forms a solid solution with delta- AlOOH by the substitutions of $\text{Mg}^{2+} + \text{Si}^{4+} \leftrightarrow 2\text{Al}^{3+}$ because of their similar crystal structure and volume. Since epsilon- FeOOH is also isostructural to MgSiO_4H_2 , phase H and delta- AlOOH , delta- AlOOH likely form partial solid solutions with epsilon- FeOOH . However, experimental study shows that FeOOH component in delta- AlOOH is limited (~20 mol%) because of the much larger unit-cell volume of epsilon- FeOOH , in contrast to the complete solid solution between phase H and delta- AlOOH . On the other hand, spin transition of Fe in epsilon- FeOOH at ~50 GPa likely reduce its volume significantly, which may cause the compositional change of hydrous phase in the lower mantle. Thus, the unit-cell volume of hydrous phases at high pressures is important to understand the chemical composition of hydrous phase in the mantle.

Here, we report new experimental results on the X-ray diffraction studies of Phase H, delta- AlOOH , epsilon- FeOOH , and their solid solutions up to 50 GPa. In-situ X-ray diffraction measurements were performed using a multianvil apparatus (SPEED-Mk.II) at BL04B1, SPring-8. The unit-cell volume of MgSiO_4H_2 phase H is ~3% larger than those of pure delta- AlOOH and ~15% smaller than those of epsilon- FeOOH up to 50 GPa. However, the large volume reduction of epsilon- FeOOH due to the spin transition of Fe was observed above ~50 GPa, resulting in the similar unit-cell volume with delta- AlOOH and phase H. Based on these experimental results, the chemical composition of hydrous phase in the lower mantle will be discussed.

Keywords: phase H, mantle, water