

## マントル遷移層における海洋地殻中の含水D相の安定領域

## Stability field of Phase D in hydrous basaltic system at the mantle transition zone

\*Liu Xingcheng<sup>1,2</sup>、松影 香子<sup>1</sup>、高橋 栄一<sup>1</sup>、鈴木 敏弘<sup>1</sup>

\*Xingcheng Liu<sup>1,2</sup>, Kyoko N. Matsukage<sup>1</sup>, Eiichi Takahashi<sup>1</sup>, Toshihiro Suzuki<sup>1</sup>

1.東京工業大学大学院理工学研究科地球惑星科学専攻、2.中国科学院广州地球化学研究所

1.Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan, 2.State Key Laboratory of Isotope Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences,

The distribution of water in the mantle and its effects on mantle dynamics are significant to understand the evolution of the Earth. It's believed that the water is transported from the surface into deep mantle via a variety of hydrous minerals in the subducting oceanic slabs (Ohtani, 2015). However, stability of hydrous mineral(s) in basaltic system at the pressure temperature conditions equivalent to cold slabs in the transition zone, remains unclear.

High-pressure and high-temperature experiments were conducted in a hydrous basalt similar to primitive MORB composition (JB-2 basalt plus brucite, water content=3.5wt.%) at 17-25 GPa, and 800-1200°C using multi-anvil apparatus (SAKURA in Magma Factory) at Tokyo Institute of Technology. We used Au/Pt double capsules as sample containers. In runs below 20 GPa, the sample was loaded into an Au inner capsule and the oxygen fugacity was buffered at ~NNO by NiO + Ni + Ni(OH)<sub>2</sub> powder, which stuffed in the space between Pt outer capsule and Au inner capsule. Simple Au capsule was used in higher pressure runs. The run products were analyzed by FE-EPMA (JEOL-JXA8530F, ELSI Tokyo Institute of Technology). The water contents in hydrous phases were estimated from total deficits in FE-EPMA analyses.

The phase assemblages of the run products include majorite, stishovite, Ca-Ti perovskite, hollandite, silicate melts and hydrous phase(s). The hydrous phase is Fe-Ti hydroxide at pressures lower than 18GPa or Al-bearing phase D at pressures higher than 18GPa. These two hydrous phases coexisted in one experiment (18GPa, 900°C). Our results show that the phase D (containing 46-51 wt.% SiO<sub>2</sub>, 19.1-20.6 wt.% MgO, 7.3-16.0 wt.% Al<sub>2</sub>O<sub>3</sub>, 3.8-5.1 wt.% FeO, 11-14 wt.% H<sub>2</sub>O) is stable in wide pressure ranges of the mantle transition zone at pressures of 18-20 GPa and temperatures of 800-1000°C. In our experiments, the Al<sub>2</sub>O<sub>3</sub> content and (Fe+Mg)/Si ratio of phase D increase with increasing pressure while MgO and FeO content remains constant, relatively. It is important to note that wadsleyite coexisting with Al-bearing phase D seemed to be nearly anhydrous in our experiment suggesting the importance of the subducting basaltic crust as a water carrier in mantle transition zone.

Consequently, we propose that the phase D is one of major water reservoir in cold subducting oceanic crust at the mantle transition zone. Moreover, if the subducting slabs stagnate at the bottom of the transition zone and warmed up, the water would be released from phase D in the subducting crusts, which may be the dominant way to hydrate the transition zone or to cause magmatism in back arc hot spots. Experiments at 25 GPa and 1000°C show coexisting bridgmanite, stishovite, Ca-perovskite, ferropericlasite and very Al-rich phase D indicating that subducting cold MORB crust could be the major carrier of water from transition layer to the lower mantle by the Al-bearing phase D.

キーワード：地球深部への水輸送、沈み込む海洋地殻、含水D相

Keywords: water transportation to deep Earth, subducting oceanic crust, phase D