

前弧かんらん岩の加水反応速度に対する実験的制約：有馬型熱水の湧出条件への応用
Experimental constraints on the serpentinization rate of fore-arc peridotites:
implications for the welling condition of the "Arima-type" hydrothermal fluids

*中谷 貴之¹、中村 美千彦¹

*Takayuki Nakatani¹, Michihiko Nakamura¹

1. 東北大学大学院理学研究科地学専攻地球惑星物質科学科

1.Division of Earth and Planetary Materials Science Department of Earth Science Graduate School of Science, TOHOKU Univ.

In order to place a constraint on the water circulation in subduction zones, hydration rates of peridotites have been investigated experimentally in fore-arc mantle conditions. Experiments were conducted at 400–580°C and 1.3 and 1.8 GPa, where antigorite was expected to form as a stable serpentine phase. Crushed powders of olivine + orthopyroxene and orthopyroxene + clinopyroxene were reacted with 15 wt% distilled water for 4–19 days. The synthesized serpentine was lizardite in all experimental conditions except that of 1.8 GPa and 580°C in the olivine + orthopyroxene system, in which antigorite was formed. In the olivine + orthopyroxene system, the reactions were interface-controlled except for the reaction at 400°C, which was diffusion-controlled. Corresponding reaction rates were 7.0×10^{-12} – $1.5 \times 10^{-11} \text{ m}\cdot\text{s}^{-1}$ at 500–580°C and $7.5 \times 10^{-16} \text{ m}^2\cdot\text{s}^{-1}$ at 400°C for the interface- and diffusion-controlled reactions, respectively. Based on a simple reaction-transport model with these hydration rates, we infer that leakage of the slab-derived fluid from an water-unsaturated fore-arc mantle is allowed only when focused flow occurs with a spacing larger than 77–229 km in hot subduction zones like Nankai and Cascadia, whereas the necessary spacing is just 2.3–4.6 m in intermediate-temperature subduction zones like Kyushu and Costa Rica. These calculations suggest that fluid leakage in hot subduction zones may occur after the fore-arc mantle is totally hydrated, while in intermediate-temperature subduction zones, leakage through a water-unsaturated fore-arc mantle may be facilitated.

キーワード：加水反応、スラブ流体、蛇紋石、前弧マントル

Keywords: hydration reaction, slab-fluid, serpentine, fore-arc mantle