岩手・宮城内陸地震震源域の3次元流体分布と地殻変動

Three dimensional fluid distribution and crustal deformation around the focal area of Iwate-Miyagi Nairiku Earthquake

鈴木 惇史²、*小川 康雄¹、齋藤 全史郎²、潮田 雅司⁵、市原 寛³、市來 雅啓⁴、三品 正明 Atsushi Suzuki², *Yasuo Ogawa¹, Zenshiro Saito², Masashi Ushioda⁵, Hiroshi Ichihara³, Masahiro Ichiki⁴, Masaaki Mishina

1.東京工業大学火山流体研究センター、2.東京工業大学地球惑星科学専攻、3.神戸大学、4.東北大学、5.産総 研

1.VFRC Titech, 2.EPS Titech, 3.Kobe Univ., 4.Tohoku Univ., 5.AIST

The 2008 Iwate-Miyagi Nairiku Earthquake (M 7.2) was an unusually large earthquake, which occurred near the volcanic regions. We used new data at 66 sites in a few kilometer grid intervals and existing data at 44 sites on three profiles. We inverted detailed 3d resistivity structure. In our final resistivity structures, aftershocks are distributed in high resistivity zones and avoiding low resistivity anomalies. We found a pathway of fluids from Mt. Kurikoma to the hypocentral area at 10km depth. The hypocenter is located at the rim of the conductor.

Large coseismic slip zones are located in high resistivity in our model. On the other hand, post-seismic slip zones are located in low resistivity zones and high resistivity zones. Iinuma et al.(2009) showed two different kinds of triggers of post-seismic slips. One comes from fluids and the other comes from the static stress change caused by mainshock. Our results imply the different triggers of post-seismic slips as implying in the previous work. These results would come from the difference of character like ductile or brittle and the existence of fluids.

キーワード:流体、比抵抗、地殻変動、マグネトテルリック法 Keywords: fluids, resistivity, crustal deformation, magnetotellurics