

地磁気伏角の 10^4 年スケールの永年変化

Secular variation of inclination with a timescale of tens of thousand years

*山崎 俊嗣¹、下野 貴也^{2,3}、井上 聖子²*Toshitsugu Yamazaki¹, Takaya Shimono^{2,3}, Seiko Inoue²

1.東京大学大気海洋研究所、2.筑波大学大学院生命環境科学研究科、3.明治大学ガスハイドレート研究所
1.Atmosphere and Ocean Research Institute, The University of Tokyo, 2.Graduate School of Life and Environmental Sciences, The Univeristy of Tsukuba, 3.Gas Hydrate Research Laboratory, Meiji University

Relative paleointensity records of marine sediments revealed that geomagnetic field fluctuations between polarity reversals contain variations with a timescale of tens of thousand years and longer. However, discussion on directional secular variations has been limited for timescale of tens to a few thousand years in general. This is probably because directional variations of the timescale of tens of thousand years are not easy to be detected due to the small amplitude of variations, often close to measurement errors, and difficulty in precise inter-core correlations. Exceptionally, inclination variations of the timescale of tens of thousand years were discussed using sediments from the western equatorial Pacific in terms of possible relations with persistent non-dipole components and orbital forcing (Yamazaki and Ioka, 1994; Yamazaki and Oda, 2002; Yamazaki et al., 2008). We revisited the problem of the long-term inclination secular variations using sediments from the Okhotsk Sea; three piston cores and nine gravity cores adjacent to each other were available. The sediments are of late Pleistocene age, and relative paleointensity was used for the age control. Inclination variations with the timescale of several to tens of thousand years are visible. Further accumulation of datasets for better spatial and temporal distribution is expected for elucidating geomagnetic field behavior of this timescale. For tectonic application of paleomagnetism assuming the virtual geocentric axial dipole field, a period of order of 100 kyr is required to average out secular variations to detect differences of several degrees in paleolatitudes.

キーワード：地磁気永年変化、伏角、オホーツク海

Keywords: geomagnetic secular variaiton, inclination, Okhotsk Sea