

マagmaオーシャン中の金属-ケイ酸塩間の塩素分配：地球の塩素枯渇の起源への示唆

Metal-silicate partitioning of chlorine in a magma ocean: Implications for the origin of chlorine depletion on Earth.

*桑原 秀治¹、後藤 弘匡²、小川 展弘³、山口 飛鳥³、高畑 直人³、佐野 有司³、八木 健彦⁴、杉田 精司^{1,4}

*Hideharu Kuwahara¹, Hirotada Gotou², Nobuhiro Ogawa³, Asuka Yamaguchi³, Naoto Takahata³, Yuji Sano³, Takehiko Yagi⁴, Seiji Sugita^{1,4}

1.東京大学大学院新領域創成科学研究科、2.東京大学物性研究所、3.東京大学大気海洋研究所、4.東京大学大学院理学系研究科

1.Department of Complexity Science and Engineering, The University of Tokyo, 2.The Institute for Solid State Physics, The University of Tokyo, 3.Atmosphere and Ocean Research Institute, The University of Tokyo, 4.Department of Earth and Planetary Science, The University of Tokyo

The chlorine of the Earth is highly depleted relative to other lithophile and volatile elements [1]. There are two hypotheses for terrestrial missing chlorine; Chlorine incorporation into the core and an erosion of primordial ocean. Here we experimentally investigate the former case. More specifically, the metal-silicate partitioning of chlorine in a magma ocean is experimentally investigated.

In this study, we investigated the effect of pressure and temperature on the metal-silicate partition coefficient of chlorine in order to estimate the core-mantle partitioning of chlorine. Starting materials were a mixture of high-purity oxides (SiO_2 , Al_2O_3 , CaO , MgO , FeO), metallic iron, and iron sulfide. Chemical compositions in the silicate portion match those of CI- or EH-chondrites. Chlorine was added to the mixture as FeCl_2 . The starting materials were encapsulated into either a graphite capsule or a single-crystal MgO capsule. The experiments were performed at 4 - 23 GPa and 1923 - 2673 K using multi-anvil presses at the University of Tokyo and Ehime University. The elemental compositions of recovered samples were analyzed with wavelength-dispersive electron microprobe (WDS-EPMA) and secondary ion mass spectrometry (SIMS). Our experimental results show that (1) chlorine is highly lithophile, (2) becomes more siderophile with increasing temperature, and lithophile with increasing pressure. Based on the experimental results and thermodynamic consideration, we estimated the metal-silicate partitioning coefficient of chlorine at the base of a magma ocean. The P - T conditions at the base of a magma ocean were estimated from the peridotite melting curve. Calculation results show that the metal-silicate partition coefficients of chlorine at the base of a magma ocean are much lower than the required value for explaining terrestrial missing chlorine. This result strongly suggests that Earth's core is unlikely to account for terrestrial missing chlorine. Given that the fluid-melt partition coefficient of chlorine is above the unity [e.g., 2], chlorine may have been partitioned into primordial ocean. If this is the case, terrestrial missing chlorine may require an extensive loss of primordial ocean during the planetary accretion phase.

[1] Sharp, Z. D. & Draper, D. S. (2013) *Earth Planet. Sci. Lett.* 369-370, 71-77.

[2] Metrich, N. et al. (2001) *Journal of Petrology* 42, 1471-1490.

キーワード：地球、塩素、マagmaオーシャン

Keywords: Earth, Chlorine, Magma ocean