

下部マントル鉱物の格子熱伝導率

Lattice thermal conductivity of lower mantle minerals

*太田 健二¹、奥田 善之¹、八木 貴志²、広瀬 敬³、新名 良介³

*Kenji Ohta¹, Yoshiyuki Okuda¹, Takashi Yagi², Kei Hirose³, Ryosuke Sinmyo³

1.東京工業大学 大学院理工学研究科 地球惑星科学専攻、2.産業技術総合研究所 計測標準研究部門、3.東京工業大学 地球生命研究所

1.Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2.National Metrology Institute of Japan, National Institute of Advanced Industrial Science and Technology, 3.Earth-Life Science Institute, Tokyo Institute of Technology

Heat in the Earth's interior is transported dominantly by convection in the mantle and core, and by conduction at thermal boundary layers. The thermal conductivity of the bottom thermal boundary layer of the mantle determines the magnitude of heat flux from the core, and is intimately related to the formation of mantle plumes, the long-term thermal evolution of both mantle and core, and the driving force for generation of the geomagnetic field (Lay et al. 2008). Recent technical progress both in the experiment and the theoretical calculation enables us to reveal high-pressure and high-temperature behavior of lattice thermal conductivity of lower mantle minerals, MgSiO_3 perovskite (bridgmanite) and MgO periclase. However, the effect of iron incorporation into these minerals on the lattice thermal conductivity is still controversial.

We measured the lattice component of thermal conductivities both of $(\text{Mg,Fe})(\text{Si,Al})\text{O}_3$ bridgmanite and $(\text{Mg,Fe})\text{O}$ ferropericlase at the Earth's lower mantle pressures and 300 K using a pulsed light heating thermoreflectance technique in a diamond anvil cell. We found that iron incorporation into bridgmanite shows minor effect on the thermal conductivity. On the other hand, the obtained conductivity of ferropericlase was considerably lower than that of MgO periclase. The estimated lattice thermal conductivity of bridgmanite-dominant lowermost mantle is comparable to conventionally assumed value of 10 W/m/K (Stacey, 1992). However, our results imply that local existence of $(\text{Mg,Fe})\text{O}$ ferropericlase in the lower mantle induce strong heterogeneity of thermal conductivity.

References

Lay, T. et al.: Nature Geosci. 1, 25-32 (2008).

Stacey, F.: Physics of the Earth, 3rd ed. (1992).

キーワード：下部マントル、熱伝導率、ブリッジマナイト、フェロペリクレーヌ

Keywords: lower mantle, thermal conductivity, bridgmanite, ferropericlase