

2014年チリ北部イキケ地震(Mw8.2)発生前に見られた加速的な滑り

Accelerated nucleation of the 2014 Iquique, Chile Mw 8.2 Earthquake

*加藤 愛太郎¹、福田 淳一²、熊澤 貴雄³、中川 茂樹²*Aitaro Kato¹, Jun'ichi Fukuda², Takao Kumazawa³, Shigeki Nakagawa²

1.名古屋大学大学院環境学研究科、2.東京大学地震研究所、3.統計数理研究所

1.Graduate School of Environmental Studies, Nagoya University, 2.Earthquake Research Institute, the University of Tokyo, 3.The Institute of Statistical Mathematics

The earthquake nucleation process has been vigorously investigated based on geophysical observations, laboratory experiments, and theoretical studies; however, a general consensus has yet to be achieved. Here, we detected a nucleation phase for the 2014 Iquique, Chile Mw 8.2 megathrust earthquake located within the current North Chile seismic gap, by analyzing a long-term earthquake catalog constructed from a cross-correlation detector using continuous seismic data. Accelerations in seismicity, the amount of aseismic slip, and the background seismicity, accompanied by an increasing frequency of earthquake migrations, started around 270 days before the mainshock at locations up-dip of the largest coseismic slip patch. These signals indicate that repetitive sequences of fast and slow slip took place on the plate interface at a transition zone between fully locked and creeping portions. These different sliding modes strongly interacted with each other and promoted accelerated unlocking of the plate interface during the nucleation phase.