

海底地震観測データから見る2011年東北沖地震震源域における地震発生前後のb値の時空間変化—地震後—

Spatiotemporal changes in the b-value along the plate interface before and after the 2011 Tohoku earthquake constrained by ocean bottom seismic network: Post-Tohoku

*仲谷 幸浩¹、望月 公廣¹、篠原 雅尚¹、山田 知朗¹、塩原 肇¹、日野 亮太²、東 龍介²、伊藤 喜宏³、村井 芳夫⁴、佐藤 利典⁵、植平 賢司⁶、眞保 敬⁶、八木原 寛⁷、小平 秀一⁸、町田 祐弥⁸、平田 賢治⁹、刈馬 弘晃⁹
*Yukihiro Nakatani¹, Kimihiro Mochizuki¹, Masanao Shinohara¹, Tomoaki Yamada¹, Hajime Shiobara¹, Ryota Hino², Ryosuke Azuma², Yoshihiro Ito³, Yoshio Murai⁴, Toshinori Sato⁵, Kenji Uehira⁶, Takashi Shimbo⁶, Hiroshi Yakiwara⁷, Shuichi Kodaira⁸, Yuya Machida⁸, Kenji Hirata⁹, Hiroaki Tsushima⁹

1.東京大学地震研究所、2.東北大学大学院理学研究科、3.京都大学防災研究所、4.北海道大学大学院理学研究院附属地震火山研究観測センター、5.千葉大学大学院理学研究科、6.防災科学技術研究所、7.鹿児島大学大学院理工学研究科附属南西島弧地震火山観測所、8.海洋研究開発機構、9.気象庁気象研究所

1.Earthquake Research Institute, University of Tokyo, 2.Graduate School of Science, Tohoku University, 3.Disaster Prevention Research Institute, Kyoto University, 4.Institute of Seismology and Volcanology, Faculty of Science, Hokkaido University, 5.Graduate School of Science, Chiba University, 6.National Research Institute for Earth Science and Disaster Prevention, 7.Nansei-toku Observatory for Earthquakes and Volcanoes, Faculty of Science, Kagoshima University, 8.Japan Agency for Marine-Earth Science and Technology, 9.Meteorological Research Institute, Japan Meteorological Agency

Changes in seismicity before and after the occurrence of megathrust earthquakes provide key information to better understanding the extension of its source area and the rupture propagation. In particular, it has been proposed that spatiotemporal variation of the b-value along the subduction zone megathrust reflects the interplate coupling [e.g., Ghosh et al., 2008, GRL]. In the case of the 2011 Tohoku earthquake, a target of this study, several studies have already reported the spatiotemporal changes of the b-value within its source area. However, it still remains controversial, because of problems with existing catalogs in completeness of magnitude after the first few months of the main shock and accuracy of focal depths for offshore earthquakes. Therefore, in this study, we analyze seismicity including small earthquakes along the plate interface using data from ocean bottom seismometers (OBSs) obtained during extensive seafloor aftershock observations [Shinohara et al., 2011, 2012] and recurrent OBS observations off Miyagi by Tohoku University. In order to automatically detect and locate interplate earthquakes, we applied a back projection method based on semblance analysis [Nakatani et al., 2015, GRL] to the data. In order to inspect validity of our method, we conducted three kinds of tests: synthetic tests, focal mechanism tests, and focal depth tests. We confirmed the validity of our method to evaluate seismicity along the plate interface. Also, we appropriately corrected event magnitudes determined by OBS records, which are, in general, overestimated due to large amplifications caused by seafloor sediments, by comparing with those listed in the JMA (Japan Meteorological Agency) catalog. Finally, we obtained an original earthquake catalog which shows an improvement in completeness of magnitude for interplate earthquakes. The resulted distribution of seismicity for the first three months after the main shock shows mutually complementary relationship between the active area of interplate aftershocks and the large coseismic slip zone. We also observe spatial variation of the b-value during the same observation period.

Acknowledgements: This work was supported by the MEXT of Japan, under KAKENHI and contract researches.

キーワード : b-value、 the 2011 Tohoku earthquake、 OBS data、 back projection

Keywords: b-value, the 2011 Tohoku earthquake, OBS data, back projection