

宮城県沖日本海溝浅部における海底地震計アレイ観測

Seismic observations using ocean bottom seismometer arrays off-shore Miyagi, northeast Japan

*太田 和晃¹、伊藤 喜宏¹、日野 亮太²、太田 雄策³、東 龍介³、篠原 雅尚⁴、望月 公廣⁴、佐藤 利典⁵、村井 芳夫⁶

*Kazuaki Ohta¹, Yoshihiro Ito¹, Ryota Hino², Yusaku Ohta³, Ryosuke Azuma³, Masanao Shinohara⁴, Kimihiro Mochizuki⁴, Toshinori Sato⁵, Yoshio Murai⁶

1.京都大学防災研究所、2.東北大学大学院理学研究科、3.東北大学大学院理学研究科附属地震・噴火予知研究観測センター、4.東京大学地震研究所、5.千葉大学大学院理学研究科、6.北海道大学大学院理学研究院附属地震火山研究観測センター

1.Disaster Prevention Research Institute, Kyoto University, 2.Graduate School of Science, Tohoku University, 3.Research Center for Prediction of Earthquakes and Volcanic Eruptions, Graduate School of Science, Tohoku University, 4.Earthquake Research Institute, University of Tokyo, 5.Graduate School of Science, Chiba University, 6.Institute of Seismology and Volcanology, Faculty of Science, Hokkaido University

Japan Trench is one of the most important subduction zones when discussing the occurrence condition of slow earthquakes. Although recent studies have identified some slow slip events (SSEs) and tectonic tremors in the shallow part of the Japan Trench subduction zone (Kato et al., 2012; Ito et al., 2013; Ito et al., 2015), the existing observation is still very limited because of the sparse seismic network. For the aim of examining detailed activities of shallow seismicity including tectonic tremors, we deployed ocean bottom seismometer (OBS) arrays near the trench. In this study, we analyze 6 months observed array data to show the fundamental performances of installed arrays. We installed three OBS arrays (AoA1-3) at interval of about 20km near the trench where the anticipated slip area of SSEs and the high coseismic slip area are overlapped. Each array consists of 5 stations spacing about 500m. The station at the center of array has a broad-band OBS and other 4 stations have a short-period OBS, respectively. While the observation is now going on with replacing of the OBSs, the first observation of AoA has been performed for 6 months from the 28 October 2014. Data from all stations have been successfully recovered on May 2015.

For each array data, we conduct a coherence analysis using the moving-window correlation technique to detect coherent signals and estimate their incoming directions (e.g. Fletcher et al., 2006). For every 4s time window, the optimum azimuth and apparent velocity are measured by maximizing the average cross correlation of all pairs of seismograms within an array with an assumption of plane waves.

We successfully detect many coherent signals. The number of signals detected simultaneously by all arrays is about 2,500 in the entire observation period, which includes regional and distant earthquakes or artificial signals like airgun shooting. About 1,000 signals correspond to the regional events in the earthquake catalog of the Japan Meteorological Agency (JMA). For corresponding events the azimuths estimated by two arrays (AoA2, AoA3) are almost consistent with azimuths from JMA epicenters, whereas the azimuths estimated by AoA1 are inconsistent and strongly biased to the specific direction of about 90 degrees, which is probably due to site effects. We also show detected tremor-like signals, though their origins are still uncertain at this time. Acknowledgements: This study is supported by JSPS KAKENHI (26000002).

キーワード：日本海溝、スロー地震、微動、海底地震計アレイ

Keywords: Japan Trench, slow earthquake, tremor, OBS array