

## 九州四万十帯延岡衝上断層における断層帯中軸部の変化に富んだ化学組成

Variable chemical composition of the Nobeoka thrust fault core in Shimanto Belt, Kyusyu

\*福地 里菜<sup>1,2</sup>、山口 飛鳥<sup>1</sup>、亀田 純<sup>4</sup>、木村 学<sup>3</sup>、芦 寿一郎<sup>1,2</sup>\*Rina Fukuchi<sup>1,2</sup>, Asuka Yamaguchi<sup>1</sup>, Jun Kameda<sup>4</sup>, Gaku Kimura<sup>3</sup>, Juichiro Ashi<sup>1,2</sup>

1.東京大学大気海洋研究所、2.東京大学大学院新領域創成科学研究科、3.東京大学大学院理学系研究科地球惑星科学専攻、4.北海道大学大学院理学研究院自然史科学専攻

1.Atmosphere and Ocean Research Institute, The University of Tokyo, 2.Graduate School of Frontier Sciences, The University of Tokyo, 3.Department of Earth and Planetary Science of the Graduate School of Science, The University of Tokyo, 4.Earth and Planetary System Science Department of Natural History Sciences, Graduate School of Science, Hokkaido University

The Nobeoka thrust in Kyushu is a tectonic boundary thrust in the Shimanto Belt, Cretaceous-Miocene accretionary complex in Southwest Japan. The Nobeoka thrust is presumed to be a fossilized megasplay fault which was branched from plate boundary fault (Kondo et al., 2005), and represents multiple deformations at seismogenic depths (~ 10 km below sea floor) (Kondo et al., 2005). Kondo et al. (2005) described lithology and macroscopic/microscopic structure of hanging wall, footwall, and the fault core. Fukuchi et al. (2014) showed mineralogical features across the fault zone based on X-ray Diffraction (XRD) analysis. However, Fukuchi et al. (2014) mainly focused on the illite crystallinity of the hanging wall side, and detail description on mineralogical/geochemical features of the fault core is still to be investigated. Therefore, this study was designed to determine chemical/mineralogical features of the fault core of the Nobeoka thrust. For this purpose, we performed elemental mapping on polished slab-shape samples retrieved the outcrop of the fault core of the Nobeoka thrust by using X-ray fluorescence (XRF) core scanner installed at Kochi Core Center (KCC).

Analyzed sample contains hanging wall, footwall, and ~15 cm-thick fault core. The fault core is bounded from both hanging- and footwall by ~3 mm-thick dark zones. Compared with the wall rocks, the dark boundaries and matrix of the fault core are enriched in Al, K, Ti, Mn, Fe and Mg, and depleted in Si, P, and S. My observation implies the following: (1) matrix of the fault core and the dark boundaries between the fault core and hanging-/footwall correspond with the enrichment of white mica and/or chlorite; (2) Depletion of S would reflect dissolution of pyrite and/or gypsum, suggesting the existence of oxidative fluid within the fault core.

キーワード：延岡衝上断層、XRF core scanner

Keywords: Nobeoka thrust, XRF core scanner