

Understanding Anomalous Eddy Vorticity Forcing in North Atlantic Oscillation Events

*Jie Song¹

1. LASG, Institute of Atmospheric Physics (IAP), Chinese Academy of Sciences

This study proposes an anomalous eddy vorticity forcing (EVF) decomposing procedure to investigate physical mechanisms responsible for the formation of the anomalous EVF associated with North Atlantic Oscillation (NAO) events. Utilizing the Geophysical Fluid Dynamics Laboratory (GFDL) dynamical core atmospheric model, a series of NAO initial-value short-term experiments are conducted. Applying the EVF decomposing procedure to the results of these experiments, the anomalous nonlinear EVF associated with the NAO events in the model can be decomposed into several fundamental linear eddy-eddy interaction terms and an unimportant nonlinear eddy-eddy interaction term. Compared with the NAO-free situation, synoptic-scale eddies have faster (slower) eastward phase speeds during the positive (negative) NAO events. Through a synoptic-scale eddy-eddy interaction mechanism, the behaviors of anomalous EVF components in the positive (negative) NAO events are well explained by synoptic-scale eddies with faster (slower) eastward phase speeds. Therefore, synoptic-scale eddies with faster (slower) eastward phase speeds are responsible for the development of the anomalous EVF associated with positive (negative) NAO events. Note that at the initial-stage of the NAO initial-value experiments, the faster (slower) phase speeds of the synoptic-scale eddies are specified by modifying the initial value fields, and then are amplified/maintained by the strengthening (weakening) zonal wind at the middle and high latitudes associated with the approaching positive (negative) phase NAO. Therefore, this study indicates that the properties of the synoptic-scale eddies at the initial-stage determine the upcoming NAO anomalies.

Keywords: The North Atlantic Oscillation, transient eddy forcing, dynamics

A comparison of the momentum budget in reanalysis datasets during sudden stratospheric warming events

*Patrick Martineau¹, Seok-Woo Son², Masakazu Taguchi³, Amy Butler⁴

1. UCLA, 2. Seoul National University, 3. Aichi University of Education, 4. NOAA/CIRES

The agreement between reanalysis datasets is evaluated during sudden stratospheric warming (SSW) events using the zonal-mean momentum budget. Zonal-mean variables reveal a good agreement among datasets for the fast warming and weakening of the polar vortex in the lower stratosphere. Eddy fluxes and forcings for zonal-mean zonal wind acceleration are also relatively similar in the lower atmosphere. This agreement is, however, severely degraded in the mid-to-upper-stratosphere. Discrepancies among reanalyses are particularly large during the onset of SSW events, a period characterized by unusually strong fluxes of planetary-scale waves from the troposphere to the stratosphere, and are substantially smaller after the onset. The discrepancies are also typically larger for the most intense SSW events. While the largest uncertainty in the momentum budget originates from the Coriolis torque, momentum flux convergence also presents a non-negligible spread among the reanalyses. The uncertainty of all terms of the zonal-mean momentum equation among reanalyses is reduced in the latest reanalysis products.

Does stratospheric sudden warming occur more frequently during ENSO winters than during normal winters?

*Kanghyun Song¹, Seok Woo Son¹

1. Seoul National University

Stratospheric sudden warming (SSW) events exhibit pronounced interannual variability. Based on WMO definition of SSW, it has been suggested that SSW events occur more preferably during El Niño-Southern Oscillation (ENSO) winters (both El Niño and La Niña winters) than during normal winters. This nonlinear relationship is re-examined here by considering six different definitions of SSW. For all definitions, SSW events are detected more frequently during El Niño winters than during normal winters, in consistent with an enhanced planetary-scale wave activity. However, a systematic relationship is not found during La Niña winters. While two SSW definitions, including WMO definition, show an increased SSW frequency during La Niña winters, other definitions show no change or even a reduced SSW frequency. This result is insensitive to the choice of reanalysis datasets and ENSO index, indicating that the reported ENSO-SSW relationship is not robust but dependent on the details of SSW definition. Implication of this finding to SSW-related downward coupling and surface climate variability is also discussed.

Keywords: Stratospheric sudden warming, El Niño-Southern Oscillation, Interannual variation of polar vortex, Stratospheric-tropospheric coupling

New definition of stratospheric warming events in the Northern Hemisphere based on geometry of the polar vortex

*平野 創一郎¹、佐藤 薫¹

*Soichiro Hirano¹, Kaoru Sato¹

1. 東京大学 大学院理学系研究科 地球惑星科学専攻

1. Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo

It is known that a few kinds of warming events occur in the Northern Hemisphere (NH) winter polar stratosphere. (Labitzke 1982). In early winter, the polar vortex often displaces substantially from the Pole mainly in the middle and lower stratosphere, which is called Canadian warming (CW; Manney et al. 2001) in the NH. In the middle of winter, polar stratospheric temperature sometimes increases rapidly, accompanied by collapse of the polar vortex. Such an event is called stratospheric sudden warming (SSW; Scherhag 1952). In the end of winter, a relatively sudden warming called stratospheric final warming (SFW; Black et al. 2006) occurs every year, followed by the summer state of the polar stratosphere. Traditionally, zonal mean zonal wind is used to define SSWs (Butler et al. 2015) and SFWs (Black et al. 2006). Although their definition is simple and easy to deal with, CWs, SSWs, and SFWs may be confused. Moreover, the polar vortex has notable horizontal structure during SSWs and SFWs, which cannot be captured by the definition based on zonal mean quantities. Mitchell et al. (2013) and Seviour et al. (2013) proposed new definition of SSWs based on moment diagnostics, where the polar vortex is approximated as an equivalent ellipse (Hu 1962; Waugh 1997). The moment diagnostics allow them to define displacement and splitting SSWs separately by using centroid latitude and aspect ratio of the polar vortex. However, SSWs in their definition may include CWs and SFWs. Note also that there is almost no clear definition for CWs.

In our study, CWs, SSWs, and SFWs are defined by applying moment diagnostics to geopotential height field at 3, 10, and 30 hPa in the polar region. The three kinds of warming events are clearly distinguished in our definition.

First, displacement, splitting, and disappearance events are defined at each altitude. To detect displacement events, centroid latitude is used as in the previous studies. Displacement events are categorized into major and minor ones according to how far the polar vortex displaces from the Pole. Splitting events are defined using kurtosis, which is measure of bipolarity (Matthewman et al. 2009). If geopotential height at all the lattice points in the polar region is larger than background value in the moment diagnostics, the events are defined as disappearance events.

Next, CWs, SSWs, and SFWs are defined based on the events identified above. Displacement-SFWs (D-SFWs) and splitting-SFWs (S-SFWs) are first defined if displacement and splitting events are followed by disappearance events, respectively. SSWs and CWs are then distinguished according to at which altitude the polar vortex is more perturbed. If the polar vortex displaces further from the Pole in the lower stratosphere than in the upper stratosphere, the events are defined as D-CWs. Otherwise, the events are defined as D-SSWs. If splitting events occur only in the lower stratosphere, the events are defined as S-CWs. Other splitting events are defined as S-SSWs.

We performed composite analysis in each category of each warming phenomenon. It is shown that the definition based on geometry of the polar vortex is consistent with a dynamical aspect of each warming phenomenon. We will investigate longitudinal structures of the polar vortex and material circulation during CWs, SSWs, and SFWs, using formulae of three-dimensional residual mean flows applicable to both Rossby waves and gravity waves derived by Kinoshita and Sato (2013).

キーワード : Canadian warming、成層圏突然昇温、成層圏最終昇温

Keywords: Canadian warming, stratospheric sudden warming, stratospheric final warming

Predictability of Arctic Polar-night Jet Oscillation Events and Its Impact on the Forecast Skill of Tropospheric Circulation

*野口 峻佑¹、向川 均²

*Shunsuke Noguchi¹, Hitoshi Mukougawa²

1. 気象研究所、2. 京都大学

1. Meteorological Research Institute, 2. Kyoto University

The predictability of the extratropical stratosphere and its impact on the forecast skill of tropospheric circulation in the Northern Hemisphere are examined in the framework of Polar-night Jet Oscillation (PJO). The PJO is the dominant low-frequency mode in the winter stratosphere characterized by the poleward and downward propagation of the zonal-mean zonal wind anomalies.

By using extended-range ensemble forecast datasets provided by the Japan Meteorological Agency, we have projected statistical properties of forecast results to a phase space spanned by two leading empirical orthogonal functions representing the PJO behavior. As a result, following characteristics of predictability variations during both anomalously weak and strong events of the stratospheric polar vortex (part of such events corresponds to sudden warmings and vortex intensifications) are obtained: (1) During prominent PJO conditions, regardless of weak or strong vortex events, the forecast skill of long-lasting anomalies in the lower stratosphere is significantly enhanced for forecasts starting after the onset of anomalous events. (2) The forecast skill not only in the lower stratosphere but also in the troposphere is improved after the setup of anomalous events. However, the reduction of tropospheric forecast error sometimes becomes obscure due to tropospheric internal variabilities, especially after strong vortex events. (3) In contrast to the same positive impact on the forecast skill in the lower atmosphere, the forecast uncertainty of the stratospheric condition shows different feature depending on the strength of the stratospheric polar vortex: During weak vortex events, the temporal evolution of the ensemble spread changes drastically from the exponential growth (saturates at high level) to the linear one (remains small) associated with the breakdown of the polar vortex. On the other hand, during strong vortex events, forecasts show large uncertainty throughout the event, because the westerly wind condition in several members of ensemble forecast permits intermittent upward propagation of planetary waves although the time-averaged flux from the troposphere is anomalously low.

Thus, this study provides comprehensive knowledge for the impact and uncertainty of stratosphere-troposphere coupling in a state-of-the-art ensemble prediction system. Our results and methodologies would also be particularly useful for real-time monitoring of sub-seasonal to seasonal forecasts.

キーワード：成層圏-対流圏結合、予測可能性、季節予報

Keywords: stratosphere-troposphere coupling, predictability, seasonal forecast

A three-week total ozone reduction over Rio Gallegos in Argentina in November 2009 and its relation to blocking in the troposphere

*秋吉 英治¹、門脇 正尚²、中村 東奈^{3,1}、杉田 考史¹、原田 やよい⁴、廣岡 俊彦⁵、水野 亮⁶

*Hideharu Akiyoshi¹, Masanao Kadowaki², Haruna Nakamura^{3,1}, Takafumi Sugita¹, Yayoi Harada⁴, Toshihiko Hirooka⁵, Akira Mizuno⁶

1. 国立環境研究所、2. 日本原子力研究開発機構、3. 富士通FIP、4. 気象研究所、5. 九州大学、6. 名古屋大学宇宙地球環境研究所

1. National Institute for Environmental Studies, 2. Japan Atomic Energy Agency, 3. Fujitsu FIP Corporation, 4. Meteorological Research Institute, 5. Kyushu University, 6. Institute for Space-Earth Environmental Research, Nagoya University

A three-week total ozone reduction over the southern tip of South America in November 2009 was reported by de Laat et al. (2010). Such long lasting low total ozone is unusual for this region and season. Ozone vertical profile measurements at Rio Gallegos, Argentina (51°S, 69°W), by ozone LIDAR suggest that isentropic surfaces of 675K and 475K over Rio Gallegos were inside the Antarctic polar vortex around November 13-14 and 22-23, respectively (Wolfram et al., 2012); thus, the low total ozone lasted for three weeks. Analyses of the total ozone observed by OMI, and ERA-Interim and JRA-55 reanalysis data indicate that the low total ozone event was caused by a polar vortex migration toward the South American continent at the time of the vortex breakup, and that the migration is associated with an enhanced wave flux from the troposphere to the stratosphere at around 120-150°W and 50-60°S. In November, a large positive deviation of a 500 hPa geopotential height from the zonal-mean was evident. This large positive deviation was considered to be a blocking by diagnosing the geopotential height field in accordance with the method of Mendes et al. (2012). These results suggest a relation between the long-term low total ozone event over Rio Gallegos and a blocking phenomenon in the troposphere of the Southern Hemisphere through wave propagation from the blocking region.

キーワード：オゾン、極渦、南米、ブロッキング、波動フラックス

Keywords: ozone, polar vortex, South America, blocking, wave flux

フーリエ変換型赤外分光計(FTIR)を用いた地上観測によるつくばにおけるHCl全量の再減少

Redecrease of HCl total column density observed with Fourier transform infrared spectroscopy at Tsukuba

*代 友輝¹、村田 功¹、中島 英彰²、森野 勇²、富川 喜弘³

*Yuki Dai¹, Isao Murata¹, Hideaki Nakajima², Isamu Morino², Yoshihiro Tomikawa³

1. 東北大学、2. 国立環境研究所、3. 国立極地研究所

1. Tohoku University, 2. National Institute for Environmental Studies, 3. National Institute of Polar Research

東北大学と国立環境研究所では、国立環境研究所所有の高分解能フーリエ変換型赤外分光計(FTIR)を用いて、つくばにおいて1998年12月より大気微量成分の地上観測を行っている。今回報告する塩化水素(HCl)は主に成層圏に存在する。地上から排出されたフロン類などの塩素化合物は、大気循環により成層圏に運ばれたあと光解離等の化学変化を経て、普段はHClやClONOといったリザーバー分子として貯蓄される。南極や北極の春先に低温となり極域成層圏雲が発生することでリザーバー分子は活性なClに変換され、大規模なオゾン破壊を起こすようになる。そのためHClの大気中の存在量はオゾン層破壊における潜在的な指標のひとつである。

フロン規制以前は、成層圏大気中の塩素総量は増加傾向でありHCl全量も同様に増加していたが、フロン規制により1990年代後半から世界的に減少し始めた。しかし、Mahieu et al. [2014]ではつくばを含むFTIR観測の国際的グループNetwork for the Detection of Atmospheric Composition Change/Infrared Working Groupの8地点における1997年から2011年の期間の地上観測および人工衛星観測によるGOZCARDSデータセットにより、HCl濃度が北半球下部成層圏で2007年以降再び増加していることを発見した。さらに大気モデルと観測結果との比較により、増加の原因が北半球の大気循環の数年程度の短期的な減速であることを明らかにした。

そこで本研究は、このHClの増加が2007年以降の「数年程度の短期的な」現象であったことを明らかにするべく、2001年から2016年の期間でつくば上空におけるHCl全量を解析し経年変化を調べた。解析にはロジャーズ法を用いたスペクトルフィッティングプログラムSFIT4を使用した。

その結果、HCl全量は2001年から2006年で減少傾向、2007年から2011年で増加傾向、2012年から2016年で再び減少傾向を示し、予測通り2007年以降の増加は短期的なものであったことがわかった。またヨーロッパ中期予報センターの再解析データ(ERA Interim)を使用し、つくばのある北緯36度における成層圏下部の残差鉛直速度の帯状平均の年々変化をみたところ、2007年から2012年にかけて下降流が強まることと、2012年以降に上昇流が強まることが確認された。HClのような主に成層圏に分布する成分は成層圏大循環により分子が輸送されるため、下降流で全量が増加し、逆に上昇流で全量が減少する。このことから、残差鉛直速度の傾向は観測結果と整合的である。しかし、HClの輸送を担う成層圏大循環の年々変化の全体像をとらえるためには、つくば上空のみではなく全球の残差鉛直速度の年々変化を調べる必要がある。

キーワード：フーリエ変換型分光計、オゾン破壊、塩化水素

Keywords: FTIR, Ozone depletion, Hydrogen Chloride

Latitudinal distributions of gravitational separation and mean age of the stratospheric air observed using a balloon-borne cryogenic air sampler

*石戸谷 重之¹、菅原 敏²、青木 周司³、森本 真司³、中澤 高清³、豊田 栄⁴、本田 秀之⁵

*Shigeyuki Ishidoya¹, Satoshi Sugawara², Shuji Aoki³, Shinji Morimoto³, Takakiyo Nakazawa³, Sakae Toyoda⁴, Hideyuki Honda⁵

1. 産業技術総合研究所、2. 宮城教育大学、3. 東北大学、4. 東京工業大学、5. 宇宙航空研究開発機構

1. National Institute of Advanced Industrial Science and Technology (AIST), 2. Miyagi University of Education, 3. Tohoku University, 4. Tokyo Institute of Technology, 5. Japan Aerospace Exploration Agency

We have collected the stratospheric air samples over Japan, the Arctic, Antarctica and equatorial regions at height levels from 10 to 35 km since 1985 by using two kinds of cryogenic air samplers on board a scientific balloon (Honda et al., 1996; Morimoto et al. 2009). The air samples were analyzed for atmospheric greenhouse gases and related air components, and we reported many findings such as spatiotemporal variations in the stratospheric CO₂ concentration and gravitational separation of major atmospheric components (e.g. Aoki et al., 2003; Ishidoya et al., 2013). Recently, many studies have focused on “mean age” of stratospheric air derived from clock tracers such as CO₂ and SF₆ to evaluate changes in the Brewer-Dobson circulation (BDC) responding to climate change (e.g. Engel et al., 2009; Ray et al., 2014). However, as Ray et al. (2014) reported, it is difficult to separate the competing effects on the mean age between mean circulation and mixing only from CO₂ and SF₆ ages since the mean age becomes younger and older by an enhancement of mean circulation and mixing, respectively, as the consequences of accelerating of BDC. In this regard, gravitational separation of stratospheric air, observed firstly by our observations, is expected to be an additional tool to constrain detail changes in BDC. Both the age and gravitational separation are unaffected by any chemical processes ideally, however the cause of gravitational separation, mass-dependent molecular diffusion superimposed on mass-independent atmospheric transport, is fundamentally different from the cause of age. Therefore, competing effects on gravitational separation between mean circulation and mixing are also expected to be different from those on age. In this study, we present latitudinal distributions of gravitational separation and CO₂ age and discuss the advantages of the simultaneous analyses of age and gravitational separation to the stratospheric circulation study.

References

Aoki et al. (2003) *Tellus*, 55B, 178-186.

Engel et al. (2009) *Nature Geoscience*, 2, 28-31.

Honda et al. (1996) *J. Geomagn. Geoelectr.*, 48, 1145-1155.

Ishidoya et al. (2013) *Atmos. Chem. Phys.*, 13, 8787-8796.

Morimoto et al. (2009) *J. Atmos. Oceanic Technol.*, 26, 2182-2191.

Ray et al. (2014) *J. Geophys. Res.*, 115, D21304.

キーワード：成層圏大気重力分離、成層圏大気年齢、大気球搭載型クライオジェニックサンプラー、ブリューワー・ドブソン循環

Keywords: gravitational separation of the stratospheric air, age of the stratospheric air, balloon-borne cryogenic air sampler, Brewer-Dobson circulation

航空機フラスコサンプリングに基づく日本-欧州間の上部対流圏/下部成層圏における温室効果ガスの季節変動とトレンド

Seasonal variations and trends of greenhouse gases in the upper troposphere/lowermost stratosphere by flask-based aircraft measurements between Europe and Japan

*澤 庸介¹、町田 敏暢²、松枝 秀和¹、丹羽 洋介¹、坪井 一寛¹、勝又 啓一²、梅澤 拓²、江藤 仁樹³、藤田 遼⁴、後藤 大輔⁵、森本 真司⁴、青木 周司⁴

*Yousuke Sawa¹, Toshinobu Machida², Hidekazu Matsueda¹, Yosuke Niwa¹, Kazuhiro Tsuboi¹, Keiichi Katsumata², Taku Umezawa², Hiroki Eto³, Ryo Fujita⁴, Daisuke Goto⁵, Shinji Morimoto⁴, Shuji Aoki⁴

1. 気象研究所海洋・地球化学研究部、2. 国立環境研究所、3. 日本航空株式会社、4. 東北大学、5. 国立極地研究所

1. Oceanography and Geochemistry Research Department, Meteorological Research Institute, 2. National Institute for Environmental Studies, 3. Japan Airlines, 4. Tohoku University, 5. National Institute of Polar Research

As part of the CONTRAIL project, we have conducted measurements of greenhouse gases (CO_2 , CH_4 , N_2O , and SF_6) by monthly air sampling in the upper troposphere/lowermost stratosphere (UT/LMS) onboard commercial airliners between Europe and Japan since April 2012. The observed mixing ratios showed sharp gradients around the dynamical tropopause defined by potential vorticity calculated from the meteorological reanalysis fields. In the UT north of 50 N, CH_4 and SF_6 were higher and seasonal phase of CO_2 were earlier than in the lower latitudes. In the LMS up to potential temperature of 50 K above the tropopause, CH_4 , N_2O , and SF_6 exhibited seasonal variations with maxima in November/December and minima in April/May. The remarkable seasonal variation in the LMS is explained by the subsidence of air from the deeper stratosphere in spring and by the efficient flushing of the LMS with tropospheric air in autumn. We observed persistent increasing trends of the all greenhouse gases over the past 5 years both in the UT and LMS. Our measurements constitute a unique data set in the UT/LMS useful for investigating temporal and spatial variations of these radiatively and chemically important greenhouse gases.

キーワード：温室効果ガス、上部対流圏/下部成層圏、航空機観測

Keywords: Greenhouse gases, Upper troposphere/Lower Stratosphere, Aircraft measurement

The Relationship between Boreal summer Intra-seasonal Oscillation and the Stratospheric Circulation

*原田 やよい¹

*Yayoi Harada¹

1. 気象研究所

1. Meteorological Research Institute

Boreal summer intra-seasonal oscillation (BSISO) is a phenomenon that active convective region migrate northward in the Indian Ocean and the western Pacific with a period of 30 –90 days. In this study, statistical relationships between BSISO and the stratospheric circulation are examined focusing on the variabilities of the zonal wind fields both in the troposphere and stratosphere. BSISO index based on Kikuchi et al. (2012) and the Japanese 55-year Reanalysis (JRA-55, Kobayashi et al. 2015) are used for composite analysis. To represent the zonal mean fields, the mass-weighted isentropic zonal mean (MIM) Method (Iwasaki 1989) is applied to JRA-55. The MIM method is suitable for the examination of zonal momentum budget, and enables analysis of a single hemispheric cell, such as Blewer-Dobson circulation in the stratosphere and the extra-tropospheric direct circulation. In order to extract variability synchronized with BSISO or lower (higher) frequent variability than BSISO, temporal filtering based on Duchon (1979) is also carried out to the zonal mean fields obtained by the application of the MIM method.

We found some characteristic features of the zonal wind field during the significant BSISO: Hemispheric symmetry westerly anomalies are observed from mid-troposphere up to the tropical tropopause layer (TTL) just after the northward migration of active convective region in BSISO region, and move poleward in both hemispheres in spite of meridional asymmetry of the convective activity: In the extra-tropical upper stratosphere of the Northern Hemisphere, both easterly anomalies and positive potential temperature anomalies are statistically significant during the northward migration of active convective region. Moreover, our results also reveal that the amplitude of BSISO is extraordinarily large during the strong low-frequency easterly anomalies at around the 20 hPa level associated with quasi-biennial oscillation (QBO) and suggest influence of the stratospheric circulation on the activity of BSISO in the troposphere.

キーワード：北半球夏季季節内振動、質量重み付き等温位面帯状平均、成層圏循環、成層圏準2年振動

Keywords: Boreal summer intra-seasonal oscillation, Mass-weighted isentropic zonal mean, Stratospheric circulation, Quasi-biennial oscillation

Investigating the tropical tropopause layer and the lower stratosphere using global models

*Dmitry Belikov¹, Fumio Hasebe¹

1. Section of Earth System Science, Faculty of Environmental Earth Science, Hokkaido University

The tropical tropopause layer (TTL) is the transition region from the troposphere to the stratosphere, and acts as a gateway to the stratosphere. Understanding all processes that control the TTL, and incorporating them in models, is an important prerequisite for reliable predictions of changes in the TTL in a changing climate and for predicting how these changes in turn feedback, e.g., via stratospheric ozone chemistry, on the global climate system. Over the past two decades, large efforts have been undertaken to improve data coverage in the TTL with the necessary vertical, spatial, and temporal resolution required to accurately characterize the transitional character of the TTL. However, due to a lack of global observations it is still not clear how the connection between the stratosphere and troposphere occurs and how it modulates the convective activity. The purpose of this research is to establish an integrated study of internal processes in the tropical tropopause layer and the lower stratosphere and to deepen understanding on atmospheric environmental change through systematic simulations and analysis of various tracers (i.e., ²²²Rn, SF₆, CO₂, CH₄, and others) and meteorological parameters (temperature, wind, water vapor and others). This study focuses on the age of air and gravitational separation in the stratosphere.

Keywords: the tropical tropopause layer, the lower stratosphere, the age of air

A method for obtaining high frequency, global, IR-based Convective Cloud Tops for studies of the TTL

*Leonhard Pfister¹, Rei Ueyama², Eric Jensen¹, Mark Schoeberl³

1. NASA Ames Research Center, 2. Bay Area Environmental Research Institute, 3. Science and Technology Corporation

Models of varying complexity that simulate water vapor and clouds in the Tropical Tropopause Layer (TTL) show that including convection directly is essential to properly simulating the water vapor and cloud distribution. In boreal winter, for example, simulations without convection yield a water vapor distribution that is too uniform with longitude, as well as minimal cloud distributions. Two things are important for convective simulations. First, it is important to get the convective cloud top potential temperature correctly, since unrealistically high values (reaching above the cold point tropopause too frequently) will cause excessive hydration of the stratosphere. Second, one must capture the time variation as well, since hydration by convection depends on the local relative humidity (temperature), which has substantial variation on synoptic time scales in the TTL.

This paper describes a method for obtaining high frequency (3-hourly) global convective cloud top distributions which can be used in trajectory models. The method uses rainfall thresholds, standard IR brightness temperatures, meteorological temperature analyses, and physically realistic and documented corrections IR brightness temperature corrections to derive cloud top altitudes and potential temperatures. The cloud top altitudes compare well with combined CLOUDSAT and CALIPSO data, both in time-averaged overall vertical and horizontal distributions and in individual cases (correlations of .65-.7). An important finding is that there is significant uncertainty (nearly .5 km) in evaluating the statistical distribution of convective cloud tops even using lidar. Deep convection whose tops are in regions of high relative humidity (such as much of the TTL), will cause clouds to form above the actual convection. It is often difficult to distinguish these clouds from the actual convective cloud due to the uncertainties of evaluating ice water content from lidar measurements.

Comparison with models show that calculated cloud top altitudes are generally higher than those calculated by global analyses (e.g., MERRA). Interannual variability in the distribution of convective cloud top altitudes is also investigated.

Keywords: Tropical Tropopause Layer, Convection, Hydration, Dehydration

Interannual and intraseasonal variations of clouds in the upper Tropical Tropopause Layer observed by CALIOP

Yuya Kowaka¹, *Hisahiro Takashima¹, Nawo Eguchi²

1. Fukuoka University, 2. RIAM, Kyushu University

Cloud variations in the tropical tropopause layer (TTL) during northern winter are investigated using the 10-year CALIPSO observations, in particular focusing on the cloud top level above the cold point tropopause (above approximately 18 km). The 10-year climatology of the TTL cloud shows higher occurrence frequency over South America, Africa, and the western-central Pacific. Interannual variation of the TTL clouds is strongly related with the TTL temperature variations associated with Quasi Biennial Oscillation (QBO) and El Nino and Southern Oscillation (ENSO). The TTL clouds associated with QBO appear/disappear simultaneously over South America, the equatorial Africa, and western-central Pacific. On the other hand, the TTL clouds associated with ENSO vary with the see-saw pattern between the western and central Pacific. We also investigated intraseasonal variations during December 2009 - February 2010. It is suggested that the temperature perturbation associated with the equatorial Kelvin wave and the sudden stratospheric warming (SSW) are important for the cloud formation. Interestingly, the TTL clouds occurred only over South America, Africa, and western-central Pacific along the Kelvin wave going east.

Keywords: tropical tropopause layer (TTL), CALIPSO, cloud

The Impact of Convection and Gravity Waves on Stratospheric Water and Upper Troposphere Cloud Fraction

*Mark R Schoeberl¹, Andrew Dessler, Eric J Jensen, Melody Avery, Leonhard Pfister

1. Science and Technology Corporation

Using our forward-domain-fill trajectory model we have run a series of experiments to explore the impact of convection and gravity waves on TTL cloud fraction and stratospheric water vapor. We compare results using MERRA convective fields and a satellite-based estimate of convective cloud heights. Gravity wave information comes from the Loon super pressure balloons. We compare our results to MLS stratospheric water vapor and CALIOP cloud fraction. The use of the high spatial resolution satellite-based convective cloud heights produces little change in model stratospheric water, but a nearly 50% reduction in model high cloud fraction (well below the observed cloud fraction) compared to the model results when we use the MERRA convective fields. Using the observed gravity wave temperature fields, cloud fraction increases, but the fluctuations produce too much cloud unless we attenuate the gravity wave fields below the tropopause –as is observed in radiosonde data. The end results, observed convection plus observed gravity waves, are in excellent agreement with observations. Overall, these results suggest that mid-frequency gravity waves play a more significant role in the cloudiness of the TTL region than previously recognized.

Keywords: Stratosphere, Cirrus , Dehydration, Tropics

Tropopause Layer Change on Different Time Scales

*Tao Xian¹

1. University of Science and Technology of China

Together with a comparatively small warming trend in global surface temperature, or warming hiatus, during the last 15 years, there are stagnations in the rising and thickening trends in the tropopause layer, which is sensitive to climate change. In this study, the variation of the vertical boundaries and thickness of the tropopause layer on different time scales and their contributions to the recent tropopause layer hiatus are investigated using the radiosonde observations from the Integrated Global Radiosonde Archive during 1960-2013.

The results confirm that global trends of rising tropopause layer boundary heights and thickening of the tropopause layer have stalled during recent years. The seasonal amplitude of each tropopause layer parameter become larger during the hiatus period (2002-2013) than that during the pre-hiatus period (1960-1997), except for the tropical tropopause layer top boundary. Moreover, the correlations between the tropopause height and corresponding temperature suggested by previous studies exist in all latitude bands in the period 1960-2013, with anti-correlations in the extratropical tropopause layer and positive correlation in tropical tropopause layer top boundary. In addition, the seasonal trends in the tropopause layer parameters show that significant trend difference occur during winter and spring.

Keywords: Tropopause Layer, Radiosonde

Extreme tropical lower stratospheric water vapor and ice amounts during 2015-2016 and their relation to ENSO, QBO, and convective overshooting

Karen Hepler Rosenlof¹, *Sean M Davis^{1,3}, Melody Avery², Hao Ye⁴, Andrew Dessler⁴

1. NOAA Earth System Research Laboratory, 2. NASA Langley Research Center, 3. Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado at Boulder, 4. Texas A&M University

In this presentation, we investigate the relative roles of tropical tropopause layer (TTL, ~14–19 km) temperatures and overshooting on the tropical lower stratospheric water vapor budget using satellite observations, reanalyses, and dehydration trajectory modeling applied to the highly unusual El Niño-Southern Oscillation (ENSO) and stratospheric quasi-biennial oscillation (QBO) events of 2015-2016.

To first order, the TTL temperature field regulates the amount of water vapor entering the stratosphere by controlling the amount of dehydration in the rising air. Thus, modes of climate variability such as the stratospheric QBO, variations in tropical upwelling, ENSO have the potential to impact the stratospheric entry value of water vapor via their impact on TTL temperatures. Additionally, vigorous convection that overshoots the local tropopause might also have a direct impact on stratospheric water vapor in a manner that circumvents the TTL cold trap mechanism.

The El Niño and subsequent La Niña of 2015-2016 coincided with a remarkable perturbation to the concentration of water vapor entering the stratosphere in the tropics. At the end of 2015 during the El Niño, a decadal record amount of lower stratospheric water vapor was observed in the Western Pacific, followed by a record dry anomaly that occurred after the 2016 QBO “interruption” and during the La Niña.

Coincident with the record setting amount of water vapor at the end of 2015, the TTL Western Pacific cold pool was shifted eastward from its climatological position and aligned with the center of convection over the Central Pacific. Over this region, there was an extreme decadal record amount of convective cloud ice in the lower stratosphere observed by the CALIOP satellite lidar. A trajectory-based analysis that models hydration based solely on reanalysis temperature and wind fields can account for only about half of the observed tropical lower stratospheric moistening during this event. This suggests that unresolved dynamical processes associated with convection and/or sublimation of lofted ice particles also contributed to lower stratospheric moistening. These processes could contribute to climate change-induced stratospheric water vapor increases.

Keywords: stratospheric water vapor, tropical tropopause layer, convective overshooting, ENSO, QBO

Assessment of upper tropospheric and stratospheric water vapour and ozone in reanalyses as part of S-RIP

*Sean M Davis^{1,2}, Michaela Hegglin³, Masatomo Fujiwara⁴, Rossana Dragani⁵, Yayoi Harada⁶, Chiaki Kobayashi⁷, Craig Long⁸, Gloria Manney^{9,10}, Eric Nash¹¹, Susann Tegtmeier¹², Tao Wang¹³, Krzysztof Wargan¹⁴, Jonathon Wright¹⁵

1. NOAA Earth System Research Laboratory, 2. Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado at Boulder, 3. University of Reading, 4. Hokkaido University, 5. European Centre for Medium-Range Weather Forecasts, 6. Japan Meteorological Agency, 7. Meteorological Research Institute, Japan Meteorological Agency, 8. NOAA Climate Prediction Center, 9. NorthWest Research Associates, 10. New Mexico Institute of Mining and Technology, 11. Science Systems and Applications, Inc., Lanham, 12. GEOMAR Helmholtz Centre for Ocean Research, 13. NASA Jet Propulsion Laboratory/California Institute of Technology, 14. NASA Global Modeling and Assimilation Office, 15. Tsinghua University

Reanalysis datasets are widely used to understand atmospheric processes and past variability, and are often used as “observations” for comparison with climate model output. Because of the central role of water vapour (WV) and ozone (O₃) in climate change, it is important to understand how accurately these species are represented in the existing global reanalyses, and whether or not significant differences exist amongst them. We present results from the WV and O₃ intercomparisons that were performed as part of the SPARC (Stratosphere-troposphere Processes and their Role in Climate) Reanalysis Intercomparison Project (S-RIP). Comparisons are made over a range of timescales between the different reanalyses, and between reanalyses and observational datasets.

In addition to the intercomparisons, we discuss the treatment of WV and O₃ in reanalyses to aid future research and guide the interpretation of differences between the reanalysis fields. Because total column ozone (TCO) is assimilated in the newer reanalyses, these reanalyses generally reproduce TCO well except when data coverage is lacking, such as during polar night. We find that the vertical distribution of ozone is relatively well represented in reanalyses, particularly given that for most reanalyses there are only weak constraints on the vertical profile of ozone from observations, and that most have a simplistic representation of ozone photochemical processes.

In contrast to O₃, stratospheric WV data are not currently assimilated, with humidity observations typically used only in the troposphere below a specified vertical level at or near the tropopause. Thus, the fidelity of reanalysis stratospheric WV is sensitive to how accurately the fundamental drivers of stratospheric WV such as tropical tropopause layer temperatures, methane oxidation, and the stratospheric overturning circulation are represented. Because of these issues and the known deficiencies in the representation of stratospheric transport in reanalyses, we find much poorer agreement both amongst reanalyses and between reanalyses and independent observations. For these reasons, stratospheric WV from the current generation of reanalyses should not be used in scientific studies.

Keywords: ozone, water vapor, stratosphere, reanalyses, SPARC Reanalysis Intercomparison Project

Intercomparison of atmospheric tides in global reanalyses from the stratosphere to the lower-mesosphere

*坂崎 貴俊^{1,2,3}

*Takatoshi Sakazaki^{1,2,3}

1. ハワイ大学 国際太平洋研究センター、2. 日本学術振興会海外特別研究員、3. 京都大学生存圏研究所

1. International Pacific Research Center, University of Hawai'i, 2. JSPS Overseas Research Fellow, 3. RISH, Kyoto University

Atmospheric tides in reanalyses are worth investigating because they are important lower boundary conditions of whole atmosphere model and also because they can be used for diurnal correction of satellite measurements. This study comprehensively assesses atmospheric tides in latest reanalyses (MERRA-2, MERRA, ERA-Interim, JRA55 and NCEP-CFSR), for both migrating and nonmigrating components in the region from the stratosphere to the lower-mesosphere, during the period of 2006-2012. SABER and MLS satellite measurements are used for comparison. It is found that all reanalyses reproduce realistic tides in a qualitative way, while the quantitative difference among the data sets depends on wavenumber and frequency. Particularly, there seems a systematic bias between SABER and reanalyses for diurnal migrating tide. We also analyzed long term changes in tides and found that they are artificially affected by the change in assimilated data.


Comparison of Three Retrievals of COSMIC GPS Radio Occultation Results in the Tropical Upper Troposphere and Lower Stratosphere

*Noersomadi Noersomadi^{1,2}, Toshitaka Tsuda¹

1. Research Institute for Sustainable Humanosphere (RISH), Kyoto University, Japan, 2. Center of Atmospheric Science and Technology, National Institute of Aeronautics and Space (LAPAN), Indonesia

Combining Geometrical Optics (GO) and Wave Optics (WO), COSMIC Data Analysis and Archive Center (CDAAC) retrieved two sets of the dry atmosphere temperature (T) from COSMIC GPS radio occultation (GPS-RO), which are named as atmPrf2010 and atmPrf2013. The sewing height between WO and GO varied at 10-20 km for atmPrf2010, and it was fixed at 20 km for atmPrf2013. We also derived T by applying WO throughout the troposphere and the stratosphere up to 30 km altitude, which is named as rishfsi2013. The height resolution of the atmPrf2010 varied depending on the sewing height, while rishfsi2013 provides high-resolution T profiles up to 30 km. The T profiles by atmPrf2013 are smoothed over 500 m. Among the three datasets, we compared the T variations in the upper troposphere and lower stratosphere (UTLS) over the tropics from October 1, 2011, to March 31, 2012, when radiosonde soundings were conducted as the CINDY-DYNAMO 2011 campaign. The mean T profiles were consistent between atmPrf2010 and atmPrf2013. In the other hand, the rishfsi2013 results were colder/warmer than the CDAAC retrievals below/above the tropopause. The mean T difference between atmPrf2013 and atmPrf2010 was 0.17 K at the cold point tropopause (CPT), and -0.38 K at the lapse rate tropopause (LRT), respectively. The rishfsi2013 showed the colder T at CPT by -0.77 K and -0.59 K relative to atmPrf2013 and atmPrf2010, respectively, and the warmer T by 0.60 K and 0.20 K at LRT. During CINDY-DYNAMO we found 134 radiosonde soundings which coincided with GPS-RO within ± 3 hours and collocated within 200 km from GPS-RO. The mean T difference at CPT from radiosondes was 0.32 K, 0.49 K and -0.24 K for atmPrf2010, atmPrf2013 and rishfsi2013, respectively. That is, both atmPrf2013 and atmPrf2010 had a positive bias at CPT, while rishfsi2013 had a negative bias. Similar comparisons at LRT were -0.45 K, -0.69 K, and -0.41 K, respectively, showing a positive bias for all GPS-RO retrievals. The rishfsi2013 is consistent with the retrievals at CDAAC and radiosondes, and it is useful for the studies of mesoscale T perturbations in the UTLS because of the good height resolution.

Keywords: COSMIC GPS Radio Occultation, Full Spectrum Inversion, Retrieval Algorithm, UTLS



Development of a cloud particle sensor for radiosonde sounding

*藤原 正智¹、杉立 卓治²、清水 健作²、林 真由美³、相良 一生³、稲飯 洋一⁴、柴田 隆⁵、岩崎 杉紀⁶、清水 厚⁷、野間 靖久³、川北 英明³、中川 太郎⁸、奥村 聡⁸

*Masatomo Fujiwara¹, Takuji Sugidachi², Kensaku Shimizu², Mayumi Hayashi³, Kazuo Sagara³, Yoichi Inai⁴, Takashi Shibata⁵, Suginori Iwasaki⁶, Atsushi Shimizu⁷, Yasuhisa Noma³, Hideaki Kawagita³, Taro Nakagawa⁸, Satoshi Okumura⁸

1. 北海道大学 大学院地球環境科学研究所、2. 明星電気株式会社、3. 神栄テクノロジー株式会社、4. 東北大学、5. 名古屋大学、6. 防衛大学校、7. 国立環境研究所、8. 神栄株式会社

1. Faculty of Environmental Earth Science, Hokkaido University, 2. Meisei Electric Co. Ltd, 3. Shinyei Technology Co., Ltd, 4. Tohoku University, 5. Nagoya University, 6. National Defense Academy, 7. National Institute for Environmental Studies, 8. Shinyei Kaisha

A meteorological balloon-borne cloud sensor called the cloud particle sensor (CPS) has been developed. The CPS is equipped with a diode laser at ~ 790 nm and two photodetectors, with a polarization plate in front of one of the detectors, to count the number of particles per second and to obtain the cloud-phase information (i.e. liquid, ice, or mixed). The lower detection limit for particle size was evaluated in laboratory experiments as ~ 2 μm diameter for water droplets. For the current model the output voltage often saturates for water droplets with diameter equal to or greater than ~ 80 μm . The upper limit of the directly measured particle number concentration is ~ 2 cm^{-3} (2×10^3 L^{-1}), which is determined by the volume of the detection area of the instrument. In a cloud layer with a number concentration higher than this value, particle signal overlap and multiple scattering of light occur within the detection area, resulting in a counting loss, though a partial correction may be possible using the particle signal width data. The CPS is currently interfaced with either a Meisei RS-06G radiosonde or a Meisei RS-11G radiosonde that measures vertical profiles of temperature, relative humidity, height, pressure, and horizontal winds. In the presentation, results from four flights, two in Japan and two in Indonesia, are discussed in detail.

キーワード：雲、ラジオゾンデ

Keywords: cloud, radiosonde

非保存項とバランスする3次元残差流の考察

On the three-dimensional residual mean flow balanced with nonconservative terms

*木下 武也¹、佐藤 薫²

*Takenari Kinoshita¹, Kaoru Sato²

1. 海洋研究開発機構、2. 東京大学 大学院理学系 地球惑星科学専攻

1. Japan Agency for Marine-Earth Science and Technology, 2. Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo

波と平均場の相互作用を記述する変形オイラー平均 (TEM) 系は、中層大気の力学的な子午面循環を理解する上で非常に有用な診断ツールである。また1980年代からTEM系を3次元に拡張する研究が行われ、理論上は純粋な3次元TEM系の導出に成功していると考えられている。しかし解析を行う上で、特に物質輸送 (残差流) においてはいくつかの問題がある。例えば、2次元残差流と異なり、3次元残差流の水平成分にはバランス流 (地衡流)、鉛直成分には温位面のゆがみに伴う流れが含まれること、さらに準停滞性感星波が卓越する冬半球成層圏では、惑星波そのものによる流れが無視できないこと等である。これらの流れは実質的な物質輸送を起こさない流れと考えられている。そこで、本研究では非保存項 (特に非断熱加熱率) と2次元残差流の関係に着目し、非保存項とバランスする3次元残差流の導出を新たには行う。

キーワード：子午面循環、中層大気

Keywords: meridional circulation, middle atmosphere

Impacts of mesospheric westerly-jet instability on the middle and lower atmosphere

*江口 菜穂¹、廣岡 俊彦²、小寺 邦彦³

*Nawo Eguchi¹, Toshihiko Hirooka², Kunihiko Kodera³

1. 九州大学 応用力学研究所、2. 九州大学 理学研究院 地球惑星科学部門、3. 名古屋大学 宇宙地球環境研究所
1. RIAM, Kyushu University, 2. DEPS, Kyushu University, 3. ISEER, Nagoya University

Kodera et al. [ACP, 2016] reported that an exceptional event of a strengthening of the subtropical jet (STJ) occurred in the stratosphere in association with a sudden equatorward shift of the stratospheric polar night jet (PNJ) in early December 2011. The exceptional rapid downward extension of STJ was developed from the lower mesosphere to the lower stratosphere, and the impact of this event farther penetrated into the troposphere in two regions, in the northern polar region and the tropics. The abrupt transformation of the STJ and PNJ is found to be associated with little connection to the upward propagation of planetary waves from the troposphere.

Analyses of minor constituent and wind fields derived from Aura MLS observations show that the strengthening of the PNJ and STJ were originated from the upper mesosphere and its mechanism could be explained by a wave-mean flow interaction which seems to be caused by large-scale waves enhanced through barotropic and/or baroclinic instability in mid- to high latitudes of the mesosphere. The detailed mechanism for the strengthening of both PNJ and STJ in the mesosphere and the impact on the stratosphere and troposphere will be shown in the presentation.

キーワード：亜熱帯ジェットと極夜ジェット、中間圏-成層圏-対流圏 相互作用

Keywords: subtropical jet and polar night jet, mesosphere-stratosphere-troposphere interaction