

グリーンランドアイスコア中に捕捉された粒子中のカルシウム化学種の同定：炭酸カルシウムの大気中での中和反応と関連して

Calcium speciation of particles trapped in Greenlandic ice core associated with neutralization reaction of calcite in the atmosphere

*宮本 千尋¹、飯塚 芳徳²、坂田 昂平³、高橋 嘉夫¹

*Chihiro Miyamoto¹, Yoshinori Iizuka², Kohei Sakata³, Yoshio Takahashi¹

1. 東京大学大学院理学系研究科地球惑星科学専攻、2. 北海道大学低温科学研究所、3. 広島大学大学院理学研究科地球惑星システム学専攻

1. Department of Earth and Planetary Science, Department of Science, The University of Tokyo, 2. Institute of Low Temperature Science, Hokkaido University, 3. Department of Earth and Planetary Systems Science, Graduate School of Science, Hiroshima University

Aerosol has various influences on the surface environment of the earth, and one of the influences is its global cooling effect including direct and indirect effects. The indirect cooling effect is caused by hygroscopic aerosols which act as cloud condensation nuclei (CCN) and form clouds that can reflect sunlight. However, hygroscopicity of aerosol differs depending on the chemical species that constitute the particles. Therefore, it is important to clarify the chemical species in aerosols to estimate the degree of the indirect cooling effect. Sulfate, one of the major species of aerosols, has high CCN activity, because it is considered that most of sulfate is present as ammonium sulfate ($(\text{NH}_4)_2\text{SO}_4$) with high hygroscopicity (Pilinis et al, 1989). On the other hand, it is reported that calcite (CaCO_3) in mineral particles reacts with sulfuric acid (H_2SO_4) during atmospheric transportation and forms gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) (e.g. Jones and Prospero 1996; Takahashi et al., 2009). In our previous study, sulfate species of aerosol collected in Higashi-Hiroshima, Japan were determined, which suggests that atmospheric neutralization reaction of CaCO_3 in mineral dust with H_2SO_4 causes suppression of forming $(\text{NH}_4)_2\text{SO}_4$ decreases hygroscopicity of sulfate aerosols.

Greenlandic ice sheet preserved natural and anthropogenic trace gases and particles transported from continents in the Northern hemisphere (Delmas 1992), which is one of the important samples to reconstruct various factors on climate change of the past, and to contribute to the more accurate prediction of the climate in future. Therefore, determination of amount of $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ in mineral dust, or identification of the neutralization process of CaCO_3 in ice sheet has a potential to reconstruct the variation of sulfate species in aerosols that have information of atmospheric chemical reactions in the past, which will help us to know the CCN activity of sulfate aerosols in the Northern hemisphere. However, there is no study on quantitative determination of $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ in mineral dust in ice sheet. In this study, calcium speciation experiments of particles trapped in Greenlandic ice sheet were conducted.

Ice core were drilled at southeast Greenland, SE Dome (67.2°N, 36.4°W) in 2015. The parts of the ice dated as 1971, 1978, 1987, 1995, and 2004 were sublimated in low-temperature room (-20°C) to obtain trapped particles using the method in Iizuka et al. (2009, 2012). Calcium-bearing particles in the trapped particles were identified by micro X-ray Fluorescence (μ -XRF) mapping. Subsequently, calcium species of the particles were determined by micro X-ray absorption fine structure (μ -XAFS) spectroscopy.

As a result of the calcium speciation, $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ fraction to total calcium in 1971, 1978, and 1987 were lower than CaCO_3 fraction. In contrast, $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ fraction in 1995 and 2004 were higher than CaCO_3 fraction. It is considered that chemical reaction of CaCO_3 in mineral dust with H_2SO_4 was more active in recent 20 years. On the other hand, sulfate ion (SO_4^{2-}) concentration in ice core decreased from late 1970s. The trend of SO_4^{2-} was consistent with emission record of SO_2 in industrial countries at the time,

however, emission amount of SO_2 increased in East Asia in recent 20 years, especially in China (Crippa et al., 2016), which is also an important source of mineral dust in Greenlandic ice sheet. Therefore, it was considered that calcium species in the mineral particles trapped in the ice sheet reflected chemical reactions of calcium with H_2SO_4 in China. These results suggested that suppression of $(\text{NH}_4)_2\text{SO}_4$ formation in the Northern hemisphere was associated with the neutralization reactions of CaCO_3 with H_2SO_4 in East Asia.

キーワード：エアロゾル、アイスコア、雲凝結核、中和反応、カルシウム化学種同定、X線吸収微細構造 (XAFS)

Keywords: aerosol, ice core, cloud condensation nuclei, neutralization reaction, calcium speciation, X-ray absorption fine structure (XAFS)

Greenland ice core records of biomass burning aerosol and BVOCs over the past 60 years

*Fahmida Parvin¹, Osamu Seki¹, Yoshinori Iizuka¹, Sumito Matoba¹, Ken Sawada²

1. Institute of Low Temperature Science, Hokkaido University, 2. Faculty of Science, Hokkaido University

It is vital to fully understand climate feedback mechanism within climate system in order to improve prediction of future climate by simulation model. However, biogeochemical feedback is not fully understood. Both the biomass burning aerosol and Biogenic volatile organic compounds (BVOCs) are thought to have a potential as chemistry feedback in response to climate change. Biomass burning aerosol is emitted from burning of forests and savanna for colonization and agriculture, burning of agricultural waste, and substances burned for fuel. BVOCs such as isoprene and monoterpenes are important precursors of secondary organic aerosol (SOA) come from terrestrial vegetation and marine plankton. However, substantial role of those aerosols in climate system is still uncertain. In order to better understand roles of biomass burning aerosol and BVOC in climate system, it is needed to explore link between climate and those emissions on various time scale. In this study, we reconstruct variability of biomass burning activity and emission of BVOC over the past 60 years based on novel approach, organic molecule tracers analyses in Southeastern Greenland-Dome (SE-Dome) ice core, which provides high-time resolution reconstruction of past environment. Organic molecular tracers such as biomass burning and biogenic SOA tracers are detected in SE-Dome ice core. Levoglucosan, which is produced by pyrolysis of cellulose and hemicellulose and thus is a tracer of biomass burning, showed sporadic peaks in the years of 1961, 1964, 1994, 1998 with the largest peak in 1964. In contrast, dehydroabietic acid, a specific tracer of the pyrolysis of conifer resin showed high peak at 1959, 2003 and showed a gradual increasing trend from 2009. The concentrations of isoprene SOA tracers (2-methylglyceric acid, 2-methylthreitol and 2-methylerythritol) are also dominant in SE-Dome ice core samples. Erythritol, is an analog of 2-methyl erythritol, a tracer species for isoprene SOA showed positive correlation with sugar compounds (arabitol, fructose, glucose) and with Dehydroabietic acid from the ice core with 95% confidence level. These results suggest that erythritol comes from continental sources. Monoterpene SOA traces are not found in SE-Dome ice core. Air mass backward trajectory showed that North America is the main source region of aerosols, indicating aerosols in SE-Dome region are transported mainly from North America. This study showed that general biomass burning tracers such as levoglucosan have been sporadically transported over the southeast Greenland and levoglucosan data matches with eastern USA fire history. In contrast, the ice core record of dehydroabietic acid indicated that fires of boreal conifer forest have occurred in North America during the last decades and transported to southeastern part of Greenland. The causes of historical variability of SOA tracers are complex and depend on atmospheric circulation, changes in vegetation cover and other factors such as temperature, tropospheric oxidative capacity.

Keywords: Isoprene SOA tracer, Biomass burning, levoglucosan, BVOC

A 60-year record of isotopic compositions of nitrate preserved in the high-accumulation dome ice core, South East Greenland

*服部 祥平¹、鶴田 明日香¹、飯塚 芳徳²、植村 立³、的場 澄人²、吉田 尚弘¹

*Shohei Hattori¹, Asuka Tsuruta¹, Iizuka Yoshinori², Ryu Uemura³, Sumito Matoba², Naohiro Yoshida¹

1. 東京工業大学 物質理工学院 応用化学系、2. 北海道大学 低温科学研究所、3. 琉球大学

1. Department of Chemical Science and Engineering, School of Materials and Chemical Technology, Tokyo Institute of Technology, 2. Institute of Low Temperature Science, Hokkaido University, 3. University of Ryukyus

Nitrate (NO_3^-) is one of the major anions found in snow. NO_3^- deposition results from reactions between nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$) and atmospheric oxidants. Global main sources of NO_x are fossil fuel and biomass burning, biogenic soil emissions, and lightning, and a recent increase in NO_3^- in ice cores has been associated with increasing anthropogenic emissions of NO_x . Based on the changes in NO_3^- concentration, however, it is difficult to identify specific sources of NO_x which takes into account for the changes in NO_3^- concentrations, hindering the development of mitigation policy of anthropogenic pollution and its effect on the environment.

Isotopic compositions of NO_3^- reveal changes in the nitrogen source and its formation pathways, but ice core records for NO_3^- concentrations and its isotopic compositions are problematic because of post depositional loss of NO_3^- via photolysis. In this study, we analyzed isotopic compositions of NO_3^- preserved in the high-accumulation dome ice core, South East Greenland. South East Greenland has a dome whose elevation is higher than 3000 m a.s.l. with high accumulation rate (about 1 m yr^{-1}) in water equivalent. High elevation and accumulation rate gives high-time resolution reconstruction of past environment, and provides negligible effect of the post depositional loss of nitrate (NO_3^-). In fact, the nitrogen isotopic compositions for NO_3^- are generally lower than those reported in Summit, Greenland, suggesting negligible effect of post depositional loss of NO_3^- in this site. In the presentation, we present changes in NO_3^- concentration and its isotopic composition through recent 60 years, and discuss the changes in the source and formation pathways of nitrate.

キーワード：安定同位体、硝酸

Keywords: stable isotope, nitrate

Isotopic constrains on post-depositional processing of snow nitrate in eastern Dronning Maud Land, East Antarctica.

*野呂 和嗣¹、服部 祥平²、植村 立³、福井 幸太郎⁴、平林 幹啓⁵、川村 賢二⁵、本山 秀明⁵、吉田 尚弘²、竹中 規訓¹

*Kazushi Noro¹, Shohei Hattori², Ryu Uemura³, Kotaro FUKUI⁴, Motohiro Hirabayashi⁵, Kenji Kawamura⁵, Hideaki Motoyama⁵, Naohiro Yoshida², Norimichi Takenaka¹

1. 大阪府立大学大学院工学研究科、2. 東京工業大学、3. 琉球大学、4. 立山カルデラ砂防博物館、5. 国立極地研究所
1. Graduate School of Engineering, Osaka Prefecture University, 2. Tokyo Institute of Technology, 3. University of Ryukyu, 4. Tateyama Caldera Sabo Museum, 5. National Institute of Polar Research

Stable isotopic ratio of nitrate in deep ice core would be valuable information for paleo-atmospheric analysis. However, post-depositional change should have an effect on the information. In previous study, this process is pointed out as physical release or photochemical reaction of nitrate on snowpack surface. Furthermore, the reactions are rare source of nitrogen oxides(NO_x) for clean atmosphere in Antarctica. To determine the mechanism of the process, isotopic composition of N(d15N) is used as an indicator for nitrate photolysis. Nitrate photolysis as post-depositional change enriches d15N for remaining nitrate on snowpack. Thus, the isotope composition(d15N) enable us to estimate how post-depositional process work in the Antarctic Plateau. In order to test spatial variation of isotopic compositions, between coastal site and inland site, here we present latitudinal variation of d15N value of NO₃⁻ in surface snow in eastern Dronning Maud Land, East Antarctica. Snow samples were collected from the surface to depths of 80 cm and 30 cm at low- and high-elevation sites during the 54th Japanese Antarctic Research(JARE), and the surface to depth 50 cm at coastal sites during 57th Japanese Antarctic Research, respectively. The d 15N of nitrate in snow were considerably increased from coastal to inland based on sample collected in 54th JARE. For the snow pit analysis in 57th JARE from surface to 180 cm, d15N ranged from -8.5 permil to +30.5 permil, and there no appreciably change. In contrast, spatial variability was observed in coastal sites samples within 100 km in JARE57. It suggests that the nitrate source would be different in the small area. The NO₃⁻ mass fraction *f* of pit samples on each depths were evaluated using measured d15N values, assuming initial d15N value(-10 permil), NO₃⁻ concentrations, and atmosphere-snow fractionation constant ¹⁵ε (-60 permil) for δ 15N based on a Rayleigh-type process (Frey et al., 2009; Berhanu et al., 2014). From the calculation, *f* in the pit samples ranged 0.51 to 0.97. This value would be corresponding to nitrogen oxides flux value at the same site in JARE57. In detail, high concentration NO and HONO were emitted from snow surface. NO₂, however, wasn't detected from snow surface and in the atmosphere.

キーワード：同位体、硝酸

Keywords: isotope, Nitrate

南極ドームふじアイスコアの解析から得られた過去72万年間における気候の不安定性と平均状態の関係

State dependence of climatic instability over the past 720,000 years from Antarctic ice cores and climate modelling

*川村 賢二^{1,2,3}、阿部 彩子⁴、本山 秀明^{1,2}、ドームふじ アイスコアプロジェクト

*Kenji Kawamura^{1,2,3}, Ayako Abe-Ouchi⁴, Hideaki Motoyama^{1,2}, Dome Fuji Ice Core Project

1. 情報・システム研究機構 国立極地研究所、2. 総合研究大学院大学、3. 海洋研究開発機構、4. 東京大学

1. National Institute of Polar Research, Research Organization of Information and Systems, 2. SOKENDAI, 3. JAMSTEC, 4. University of Tokyo

Climatic variabilities on millennial and longer timescales with a bipolar seesaw pattern have been documented in palaeoclimatic records, but their frequencies, relationships with mean climatic state, and mechanisms remain unclear. Understanding the processes and sensitivities that underlie such changes will underpin better understanding of the climate system and projections of its future change. We investigate the long-term characteristics of climatic variability using a new ice-core record from Dome Fuji, East Antarctica, combined with an existing long record from the Dome C ice core. Antarctic warming events over the past 720,000 years are most frequent when the Antarctic temperature is slightly below average on orbital time scales, equivalent to an intermediate climate during glacial periods, whereas interglacial and fully glaciated climates are unfavourable for a millennial-scale bipolar seesaw. Numerical experiments using a fully coupled atmosphere–ocean general circulation model (AOGCM) with freshwater hosing in the northern North Atlantic showed that climate becomes most unstable in intermediate glacial conditions associated with large changes in sea ice and the Atlantic Meridional Overturning Circulation (AMOC). Model sensitivity experiments suggest that the prerequisite for the most frequent climate instability with bipolar seesaw pattern during the late Pleistocene is associated with reduced atmospheric CO₂ concentration via global cooling and sea ice formation in the North Atlantic, in addition to extended Northern Hemisphere ice sheets.

キーワード：ドームふじ氷床コア、古気候、二酸化炭素、突然の気候変動、大西洋子午面循環

Keywords: Dome Fuji ice core, Paleoclimate, CO₂, Abrupt Climate change, Atlantic meridional overturning circulation

Stability of AMOC and bipolar seesaw under different background climatic condition

*阿部 彩子^{1,2,3}、陳 永利¹、川村 賢二^{3,2}、大垣内 るみ²、吉森 正和⁴、岡 顕¹、シェリフ多田野 サム¹、小長谷 貴志¹

*Ayako Abe-Ouchi^{1,2,3}, Wing-Le Chan¹, Kenji Kawamura^{3,2}, Rumi Ohgaito², Masakazu Yoshimori⁴, Akira Oka¹, Sam Sherriff-Tadano¹, Takashi Obase¹

1. 東京大学大気海洋研究所、2. 独立行政法人 海洋研究開発機構、3. 国立極地研究所、4. 北海道大学大学院地球環境科学研究院

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. JAMSTEC, 3. National Institute of Polar Research, 4. Hokkaido Univeresity

Millennial climate changes known as D-O cycles and AIM recorded in ice cores in both Hemispheres show a higher amplitude in the middle-level of a glacial cycle than in the interglacial state or severe glacial state. Here we investigate the stability of AMOC and climate by analyzing several sensitivity experiments using a coupled atmosphere and ocean GCM (MIROC4m). The stability under different climates are compared; modern climate state with the pre-industrial condition, middle glacial climate state and full glacial condition, mainly differing in the ice sheet configuration and the amount of Greenhouse Gases. The results under middle glacial condition show the largest cooling/warming response in North Atlantic and a reasonable bipolar warming/cooling signal, which are consistent to ice core data and deep-sea data. We show the stability diagram of AMOC in the model under different background conditions and discuss the implication on the mechanism of abrupt climate changes in the past.

キーワード：気候、気候モデル、古気候

Keywords: climate, climate model, paleoclimate

Importance of the sea ice-surface wind feedback on simulating the LGM AMOC

*シェリフ多田野 サム¹、阿部 彩子¹

*Sam Sherriff-Tadano¹, Ayako Abe-Ouchi¹

1. 東京大学大気海洋研究所

1. Atmosphere and Ocean Research Institute, University of Tokyo

Simulating and understanding the dynamics of the glacial Atlantic meridional overturning circulation (AMOC) is one of the main challenges in the paleoclimate community. However, most climate models in the Paleoclimate Model Intercomparison Project cannot reproduce a weak AMOC in their simulations of the last glacial maximum (LGM). Previous studies show that a stronger glacial surface wind over the northern North Atlantic induced by the Laurentide ice sheet is causing this model-data discrepancy. They suggest that a weaker wind over the northern North Atlantic may be important in simulating the weak LGM AMOC. Dome F members (2017) recently succeed in simulating a weak LGM AMOC in their simulations using a coupled model MIROC, and show that the surface wind over the northern North Atlantic is substantially reduced compare to previous LGM simulations. Differences in the sea ice extent over the northern North Atlantic are suggested to be the cause of the weaker surface wind, though the impact of the sea ice extent on the surface wind remains elusive. In addition, it remains unclear that whether the weakening of the surface wind plays a role in simulating the weak LGM AMOC. Therefore, in this study, we first investigate the role of sea ice expansion on the weakening of the surface wind, and then we explore the impact of the weakening of the surface wind on the LGM AMOC. For this purpose, simulations results from MIROC4m are utilized (Dome F members, 2017). To clarify the impact of changes in sea ice and associated changes in diabatic heating on the surface wind, sensitivity experiments are conducted with an atmospheric general circulation model and a linear baroclinic model. Additional experiments are conducted with the coupled model MIROC, which we modified the surface wind stress over the northern North Atlantic to assess the impact of weakening of the surface wind on the LGM AMOC. Results show that expansion of sea ice substantially weakens the surface wind over the northern North Atlantic. Analysis and experiments with linear baroclinic model shows that two processes are crucial in reducing the surface wind; the suppression of atmosphere-ocean heat exchange and the increase in the stability of the boundary layer over the sea ice due to an intense surface cooling. Additional experiments with MIROC show that the weakening of the surface wind due to the expansion of sea ice plays a role in maintaining the weak AMOC in the LGM simulation.

キーワード : LGM、AMOC、地表風

Keywords: LGM, AMOC, surface wind

グリーンランドNEEM氷床コアと南極ドームふじ氷床コアによる完新世のメタン濃度の復元

Atmospheric CH₄ concentration during the Holocene reconstructed from the NEEM (Greenland) and Dome Fuji (East Antarctica) ice cores

*大藪 幾美¹、川村 賢二^{1,2,3}、東 久美子^{1,2}、北村 享太郎¹、中澤 高清⁴、青木 周司⁴、Brook Edward⁵、Blunier Thomas⁶

*Ikumi Oyabu¹, Kenji Kawamura^{1,2,3}, Kumiko Goto-Azuma^{1,2}, Kyotaro Kitamura¹, Takakiyo Nakazawa⁴, Shuji Aoki⁴, Edward Brook⁵, Thomas Blunier⁶

1. 情報・システム研究機構 国立極地研究所、2. 総合研究大学院大学、3. 海洋研究開発機構、4. 東北大学、5. オレゴン州立大学、6. コペンハーゲン大学

1. National Institute of Polar Research, Research Organization of Information and Systems, 2. SOKENDAI (The Graduate University of Advanced Studies), 3. Japan Agency for Marine Science and Technology, 4. Tohoku University, 5. Oregon State University, 6. University of Copenhagen

Methane (CH₄) is an important greenhouse gas, whose atmospheric concentration has been increasing due to human activities for the last few centuries. Orbital-scale variations of atmospheric CH₄ correlate with climatic precession, because the size of wetlands and their CH₄ production rate respond to Northern Hemisphere (NH) summer insolation, through the variations in temperature and rainfall on NH landmasses. The correlation between CH₄ concentration and NH summer insolation held for the last three interglacial periods, but the relationship collapsed during the latter half of the Holocene. NH summer insolation kept decreasing whereas the atmospheric CH₄ concentration increased since ~5 kyr BP. Several explanations have been proposed for this trend, such as peat growth in circum-Arctic region¹, emission from tropical wetlands due to increasing rainfall in the Southern Hemisphere (SH)^{2,3} and agricultural activity⁴, and the exact mechanisms have been under continuing debate.

Inter polar difference (IPD) of CH₄ concentrations have provided important constraint on the evolution of CH₄ source distribution and its relationship with climate^{2,5,6,7}. However, time resolution and analytical precision of previous studies have not always been adequate to investigate precise IPD. In addition, reconstruction of accurate CH₄ variation is difficult during most of the Holocene from Greenland ice cores, because the depths for this time period often corresponds to poor quality ice (brittle zone). To reconstruct the CH₄ IPD during the Holocene, we have been measuring CH₄ concentrations in the NEEM (Greenland) and Dome Fuji (DF) (Antarctica) ice cores. Accurate CH₄ reconstruction from the Holocene NEEM ice core is challenging because of the brittle zone. We indeed found high CH₄ spikes in the brittle zone, thus we investigated them by measuring additional 3–5 samples from the neighboring depths (within ~50 cm, ~5 years) and checking the reproducibility, and then rejected the data which is more than 15 ppb higher than their means. Reproducibility after removing the outliers are ±2.5 and ±1.7 ppb for the NEEM and DF ice cores, respectively.

We investigate the integrity of our Holocene CH₄ data. The CH₄ variations of the NEEM core, including centennial to millennial variations in the brittle zone, agree well with the GISP2 data (recent high-precision data by the Oregon State University group) (ref.7 and unpublished data). For Antarctica, the variations of CH₄ concentration of the DF core also agree well with those of the WAIS divide core from West Antarctica (ref.7 & 8 and unpublished data), after considering centennial-scale smoothing effect on the DF record caused by slow gas trapping. These comparisons suggest that our new records, as well as

the most recent records by other groups, provide reliable reconstruction of the past atmospheric CH₄ variations over the entire Holocene.

IPD are deduced from two different combinations of cores: the NEEM and DF cores, and the NEEM and WAIS Divide cores. For this analyses, the gas time scales of the NEEM and DF cores are placed on the WAIS Divide ice core chronology by pattern matching of the CH₄ records. IPD from both NEEM/DF combination and NEEM/WAIS combination increased from the early Holocene to mid Holocene, and then decreased toward the late Holocene. We employ a simple 3-box model^{2,6} to deduce CH₄ emissions from different latitudinal bands at 1000-yr intervals. The model calculates the emissions in the low-latitude box (30°S-30°N) and northern box (30-90°N), while small emission from southern box (90-30°S) is kept constant. The model results show that northern emission decreased, while low-latitude emission increased during the last half of the Holocene. This suggests significant contribution from the low-latitude sources to the atmospheric CH₄ increase since ~5 ka. A recent model study suggests that CH₄ emission from the SH tropics may have increased due to SH summer insolation rise⁷. Several terrestrial proxies suggest increased rainfall in the tropical regions in South America during the latter half of the Holocene⁹, probably in response to the increase in SH summer insolation. Although we cannot reject the anthropogenic hypothesis at this stage and more investigations are needed, our results are consistent with the hypothesis that tropical SH emission was responsible for the CH₄ rise during the latter half of the Holocene.

1)Blunier et al.,1995. 2)Chappellaz et al., 1997. 3)Singarayer et al., 2011. 4)Ruddiman & Thomson., 2001. 5)Nakazawa et al., 1993. 6)Brook et al., 1996. 7)Mitchell et al., 2013. 8)WAIS Divide Project Members, 2014. 9)Prado et al., 2013.

キーワード : アイスコア、メタン、完新世、グリーンランド、南極
Keywords: Ice core, Methane, Holocene, Greenland, Antarctica

Increasing insolation and greenhouse gas concentration trigger Bølling-Allerød warming

*小長谷 貴志¹、阿部 彩子^{1,2}

*Takashi Obase¹, Ayako Abe-Ouchi^{1,2}

1. 東京大学理学系研究科地球惑星科学専攻、2. 海洋研究開発機構

1. The University of Tokyo, Department of Earth Planetary Science, 2. Japan Agency for Marine-Earth Science and Technology

During the last deglaciation, a major global warming was punctuated by several abrupt climate changes, likely related to Atlantic Meridional Overturning Circulation (AMOC) (Clark et al. 2012). Transient deglaciation experiments from the Last Glacial Maximum have been conducted by applying time-dependent insolation, greenhouse gas concentrations, and glacial meltwater forcing (Liu et al. 2009). They have showed that reduction in glacial meltwater discharge rate into North Atlantic induces abrupt recovery of AMOC, warming of Greenland and cooling of Antarctica (bipolar response) during the period of Bølling-Allerød (BA, ~14.6 ka).

We conduct a transient simulation from the Last Glacial Maximum to BA using an atmosphere-ocean coupled general circulation model (AOGCM) MIROC 4m (an IPCC-class Japanese community model). The model is initialized with the 21ka, and we change insolation, greenhouse gas concentrations and meltwater fluxes following the protocol of PMIP4 (Ivanovic et al. 2016). Glacial meltwater is derived from ice sheet reconstruction (ICE6g, Peltier et al. 2015). We assume the glacial meltwater due to ice sheet loss is uniformly applied to the area of 50-70N North Atlantic Ocean. We conduct additional experiments branched from 16 ka, where 50-80% of ICE6g meltwater fluxes are applied without reducing the meltwater fluxes before the BA.

The model results show that abrupt resumption of AMOC and warming of Greenland occurred at around the period of BA even under hosing of 0.06 Sv. Transition from cold stadial mode to warm interstadial mode occurs in about 100 years, which is consistent with reconstructions (Buizert et al. 2014). The result implies that increasing summer insolation and greenhouse gas concentration trigger abrupt AMOC recovery and warming in the Northern Hemisphere, and large fluctuation of meltwater due to ice sheet melting may not be necessary.

更新世の長周期気候変動の解明に向けて:氷床モデルによる更新世初期の4万年変動の再現

Toward understanding the climate change in the Pleistocene: Reproduction of the dominant 40-kyr periodicity in the early Pleistocene using an ice-sheet model

*渡辺 泰士¹、阿部 彩子¹、齋藤 冬樹²

*Yasuto Watanabe¹, Ayako Abe-Ouchi¹, Fuyuki SAITO²

1. 東京大学、2. 海洋研究開発機構

1. The University of Tokyo, 2. JAMSTEC

The climate change in the Pleistocene is characterized by glacial-interglacial cycles that have a dominant periodicity at tens of thousands of years. Milankovitch theory suggests that variation of Earth's orbital parameters changes the way the sunlight enters the Earth and the northern hemisphere high-latitude summer insolation that have a dominant periodicity at 20-kyr causes the glacial-interglacial cycles. In contrast to this periodicity, however, the dominant periodicity of the climate change in the early Pleistocene was 40-kyr. Various hypotheses are proposed but it has not yet been fully understood. One reason is that the 40-kyr cycles have not been reproduced using a realistic 3-D model with realistic input. Our aim is to reproduce the 40-kyr cycles using a 3-D ice-sheet model with realistic input and to reveal the role of each orbital parameter by comparing the result with records of proxy. We used an ice-sheet model for Integrated Earth system Studies (ICES; Abe-Ouchi et al., 2013), coupled with climate parameterization according to the results of a global climate model MIROC. The input is variability of insolation and atmospheric CO₂ concentration and the output is time evolution of ice-sheet distribution over the northern hemisphere.

We conducted experiments for two 40-kyr cycles. One is from MIS-49 to 47 and the other is MIS-45 to 43. These cycles are chosen as a representative of 40-kyr cycles that has a long and stable interglacial and that has a relatively short interglacial and long glacial respectively.

As a result, 40-kyr cycles are reproduced and the shape of the variation is similar to the proxy record. This is because the surface temperature exceeds a threshold that the North American ice-sheet starts deglaciating once in a 40-kyr cycle. Phase analysis of this result suggests that the difference in the shape of these cycles is explained by lead-lag relationship between obliquity and precession. For a long interglacial period, the peak of climatic precession precedes that of obliquity, and vice versa for a short interglacial period.

In summary, climatic precession decides a timing of a deglaciation because of the large influence on insolation and obliquity has a role as a pacemaker of 40-kyr cycles because of an existence of a threshold of an ice-sheet deglaciation. These are the role of the orbital parameters in the early Pleistocene. We would also analyse several sensitivity experiments under different basal conditions, and those coupled with the temperature anomaly calculated by using atmosphere-ocean-vegetation GCM MIROC-LPJ (Oishi and Abe-Ouchi, 2011).

キーワード : 更新世、氷期間氷期サイクル

Keywords: Pleistocene, glacial-interglacial cycles

High glacial dust amount worked to warm the polar regions at the Last Glacial Maximum: a modelling study using MIROC-ESM

High glacial dust amount worked to warm the polar regions at the Last Glacial Maximum: a modelling study using MIROC-ESM

*大垣内 るみ¹、阿部 彩子²、大石 龍太²、竹村 俊彦³、渡邊 真吾¹、河宮 未知生¹

*Rumi Ohgaito¹, Ayako Abe-Ouchi², Ryouta O'ishi², Toshihiko Takemura³, Shingo Watanabe¹, Michio Kawamiya¹

1. 海洋研究開発機構、2. 東京大学大気海洋研究所、3. 九州大学応用力学研究所

1. Japan Agency for Marine-Earth Science and Technology, 2. Atmosphere-Ocean Research Institute, Univ. of Tokyo, 3. Research Institute for Applied Mechanics, Kyushu University

Aerosol and its impacts on climate is one of uncertain factors on future climate projection. One of the aerosol species, mineral dust aerosol (dust) can be investigated with the past climate changes. Especially, need of more investigation of the feedback of dust and vegetation under the different climatic condition compared to present day is repeatedly mentioned in the paleoclimate chapter of Intergovernmental Panel on Climate Change 5th Assessment Report. Last Glacial Maximum (LGM, c.a. 21,000 years before present) is known with the enhancement of dust, globally but especially over the high latitude regions from the ice core and sediment core proxy data (Winkler et al. 2008, Lamy et al. 2014). It has been difficult to reproduce dust amount over the high latitudes with General Circulation Models (GCMs) at LGM and the effect of dust on high latitudes has not been cleared. The previous works used atmospheric part of GCM or Earth System Model (ESM) (Mahowald et al. 2006, Takemura et al. 2009, Albani et al. 2012, Hopcroft et al. 2015). For the first time, using an ESM, MIROC-ESM (Watanabe et al. 2011), we tested the impact of enhanced dust on LGM climate with adding glaciogenic dust (Mahowald et al. 2006) (hereafter, called LGMglac) on a standard LGM simulation following the Paleoclimate Modelling Intercomparison Project phase 3 (PMIP3) protocol (Sueyoshi et al. 2013). The resulting deposition distribution of dust in LGMglac matched better to the latest global dust data archives (Kohfeld et al. 2013, Albani et al. 2014). The experiment LGM is the identical with the one for PMIP3. LGMglac deviated from the LGM in a spin-up stage and the corresponding period with the LGM experiment is taken for the analyses. We have found that the high LGM dust amount warms the northern high latitudes and the surrounding of the Antarctica. Sensitivity experiments using atmospheric part of MIROC-ESM suggested that both of radiative forcing and the aging of snow and ice albedo by dust are important for the LGMglac-LGM warming in the northern hemisphere. On the other hand, over the Antarctica, the positive radiative forcing at surface plays a role for the warming with additional dust but the effect on the surface temperature of the high glacial dust amount is a little at the borehole sites over the high plateau of the Antarctica.

キーワード：ダスト、最終氷期最大期

Keywords: Dust, LGM

東南極氷床「ドームふじ」頂部位置の氷期・間氷期の変遷に応じた移動について

On the migration of Dome Fuji summit of East Antarctica over glacial - interglacial periods

*藤田 秀二^{1,2}、福井 幸太郎³、本山 秀明^{1,2}、中澤 文男^{1,2}

*Shuji Fujita^{1,2}, Kotaro Fukui³, Hideaki Motoyama^{1,2}, Fumio Nakazawa^{1,2}

1. 大学共同利用機関法人 情報・システム研究機構 国立極地研究所、2. 総合研究大学院大学 複合科学研究科 極域科学専攻、3. 立山砂防カルデラ博物館

1. National Institute of Polar Research, Research Organization of Information and Systems(ROIS), 2. Department of Polar Science, The Graduate University for Advanced Studies (SOKENDAI), 3. Tateyama Caldera Sabo Museum

南極ドームふじ近傍は、Oldest Ice（100万年を上回る年代をもつ最古のアイスコア）の掘削候補地として着目されている地域である。日本の南極観測では、1980年代からこの近傍のアイスコア研究と雪氷環境研究にターゲットをしばった観測研究をおこない、1990年代と2000年代にわたり2本の深層アイスコアを取得した。将来に掘削を計画するOldest Iceの掘削候補地の同定には、氷床表面環境、内部環境、底面環境の入念な観測とデータの検討が必要である。私達は、これまでこの地域で日本が実施してきた観測データをまとめ、氷床内部層の3次元構造の分析をすすめている。その結果、氷床内部に蓄積された歪みのパターンは、現在のドームふじの南方に過去のドームがあることを示唆した。さらに、最終氷期のLGM以降に堆積した雪の量は、強い南北勾配をもち、北方すなわち海側ほど堆積が多いことがわかった。最終氷期の当年代面を氷床表面とした地形図を作成すると、氷期の氷床ドームは現在のドームふじの南方約60kmの、氷下に山塊のある地域に出現した。研究発表ではデータ分析の最新状況を示す。

キーワード：南極、氷床、アイスコア

Keywords: Antarctica, ice sheet, ice core

「最古の氷」に対する $^{26}\text{Al}/^{10}\text{Be}$ 放射年代決定の可能性：ドームふじ第二期深層コアからの洞察

On the possibility of radiometric dating on the “oldest ice” using cosmogenic $^{26}\text{Al}/^{10}\text{Be}$ ratio: Insights from the Dome Fuji second deep (DF2) ice core

*堀内 一穂¹、内田 智子²、菅原 愛¹、清野 祥子¹、松田 紗耶¹、松崎 浩之³、本山 秀明⁴

*Kazuho Horiuchi¹, Tomoko Uchida², Ai Sugawara¹, Shoko Seino¹, Saya Matsuda¹, Hiroyuki Matsuzaki³, Hideaki Motoyama⁴

1. 弘前大学、2. 東北大学、3. 東京大学、4. 極地研

1. Graduate School of Science and Technology, Hirosaki University, 2. Institute of Geology and Paleontology, Tohoku University, 3. MALT, The University of Tokyo, 4. National Institute of Polar Research

Deep ice core records have been playing a crucial role in paleoclimatology. However, there are no records that have reached over ca. 800 ka. To overcome this problem, several research groups are planning to recover the “oldest ice” (~1.5 Ma) from near the bases of ice-sheets overlying inland Antarctica (see e.g. Schiermeier, 2016). Nevertheless, it may be not an easy task to obtain an accurate chronology from such deepest parts of ice cores, because layer inclination and/or folding may prevent us to construct accurate stable-isotope chronologies, which usually rely on normal stratigraphy.

Cosmogenic ^{26}Al and ^{10}Be are produced by interactions of cosmic rays with specific elements in the atmosphere. Because the atmospheric production is similar between ^{26}Al and ^{10}Be , an exponential decrease of the $^{26}\text{Al}/^{10}\text{Be}$ ratio with time, in ordinary cases, should represent the difference of the decay constants of the nuclides ($T_{1/2}$ of the $^{26}\text{Al}/^{10}\text{Be}$ ratio is 1.45 Myr). In this presentation, we show the profiles of the $^{26}\text{Al}/^{10}\text{Be}$ ratios of certain stratigraphic intervals of the Dome Fuji second deep (DF2) ice core, plotted against the latest age model for this core (Dome Fuji ice core project members, 2017). By investigating these data, we discuss about the possibility of the radiometric dating on the “oldest ice” of an age of up to 1.5 Ma.

連続融解分析（CFA）システムによる南極ドームふじ深層コアの高時間分解能分析

High resolution analyses of the Dome Fuji deep ice-core using a Continuous Flow Analysis (CFA) System

*東 久美子^{1,2}、平林 幹啓¹、尾形 純¹、北村 享太郎¹、川村 賢二^{1,2}、塚川 佳美¹、Dallmayr Remi¹、中澤 文男^{1,2}、福田 かおり¹、荒井 美穂³、藤田 秀二^{1,2}、本山 秀明^{1,2}

*Kumiko Goto-Azuma^{1,2}, Motohiro Hirabayashi¹, Jun Ogata¹, Kyotaro Kitamura¹, Kenji Kawamura^{1,2}, Yoshimi Ogawa-Tsukagawa¹, Dallymayr Remi¹, Fumio Nakazawa^{1,2}, Kaori Fukuda¹, Miho Arai³, Shuji Fujita^{1,2}, Hideaki Motoyama^{1,2}

1. 国立極地研究所、2. 総合研究大学院大学、3. 山形大学

1. National Institute of Polar Research, 2. SOKENDAI, 3. Yamagata University

国立極地研究所ではアイスコアを高時間分解能で連続分析するため、連続融解分析（Continuous Flow Analysis、略してCFA）システムを開発した。本システムはアイスコアの融解ユニットと分析ユニットから構成される。分析ユニットは大きく分けて、アイスコアの融解水を分析する装置と、アイスコアに含まれる空気成分を分析する装置で構成される。現在、前者として電気伝導度計、レーザー式水同位体比アナライザー、ICP質量分析計、固体微粒子分析計、ブラックカーボン分析装置を用いており、後者としてはレーザー式メタンガス分析装置を用いている。また、融解水の一部はフラクションコレクターにより、自動的にサンプル瓶に注入される。国立極地研では、このCFAシステムを用いて南極ドームふじ深層コアの分析を開始した。本発表では、融解水を用いた分析について報告する。まず、本システムの性能を評価するために実施した様々なテストについて紹介し、次に、初期的な分析結果について報告する。

キーワード：CFA分析、深層氷床コア、南極ドームふじ

Keywords: CFA analyses, Deep ice core, Dome Fuji, Antarctica

連続融解システムによるドームふじ氷床コア中のメタン濃度の高分解能測定

High-resolution measurements of methane concentration in the Dome Fuji ice core using Continuous Flow Analysis (CFA) system

*荒井 美穂¹、川村 賢二^{2,3,4}、北村 享太郎²、東 久美子^{2,3}、尾形 純²、Dallmayr Remi²、中澤 文男^{2,3}、平林 幹啓²、中澤 高清⁵、青木 周司⁵、鈴木 利孝¹

*Miho Arai¹, Kenji Kawamura^{2,3,4}, Kyotaro Kitamura², Kumiko Goto-Azuma^{2,3}, Jun Ogata², Remi Dallmayr², Fumio Nakazawa^{2,3}, Motohiro Hirabayashi², Takakiyo Nakazawa⁵, Shuji Aoki⁵, Toshitaka Suzuki¹

1. 山形大学大学院理工学研究科、2. 国立極地研究所、3. 総研大、4. 海洋研究開発機構、5. 東北大学理学研究科 大気海洋変動観測研究センター

1. Graduate School of Science and Engineering, Yamagata University, 2. National Institute of Polar Research, 3. SOKENDAI, 4. JAMSTEC, 5. Center for Atmospheric and Oceanic Studies, Graduate School of Science, Tohoku University

国立極地研究所では、氷床コアの連続融解分析（CFA）システムが開発され、ドームふじ氷床コアの測定が開始された。CFAシステムでは連続的に氷床コアを融解し同時に様々な測定をするため、多くの情報を得ることが可能となる。融解水に含まれる成分（水同位体、化学成分、ダスト、ブラックカーボン等）の測定だけでなく、氷床コア中に保存された過去の大気を抽出しメタンの濃度を測定することができる。

CFAシステムを用いるメリットは、従来の測定に比べて圧倒的に高い分解能のデータを得られることにある。メタンの詳細な変動データによって、これまでの手法では見ることのできなかった新たな古環境変動が見出される可能性がある。また、複数のコアの年代をより正確に合わせて比較することが可能となる。例えば、Buizertら（2015）は、CFAによって得られた南極WAIS Divide氷床コアのメタン濃度データとグリーンランドNGRIPコアの酸素同位体比データ（気温の指標）を対比させることにより、両コアの気温変動のタイミングを正確に比較した結果、南極の気温の変曲点（極大、極小）がグリーンランドの急激な気温変化に約200年遅れることを見出した。ドームふじコアのような低涵養量の地点のコアでは、空気の年代決定に不可欠となる氷と空気の年代差の誤差が大きいが、これが他のコアとの比較によって正確に推定できるようになる可能性がある。

CFAにおいて、氷の融解水からの気体の抽出には、通常脱気装置に用いられるような薄膜を使用する。しかし、溶存気体の抽出は完全には行えないため、そのことによる測定対象成分の分別効果を補正する必要がある。その量を、超純水に濃度既知の標準ガスを混入させて実際のアイスコア試料と同様に抽出する実験によって見積もった結果、メタン濃度の場合には真の値に比べて数%低めに測定されると推定した。

第2期ドームふじコアのCFA分析は、最終的には表面から2400m（約30万年前）にかけて行われる予定であり、現在、主に最終退氷期を対象とした分析を開始したところである。予稿投稿時点では、300～314m（ガス年代：約7500–8000年前）の測定が行われた。気体抽出時の分別を補正したCFA分析値と、氷床コアを個別に分析する従来の測定方法による値との差は約1%以下であり、補正が正しく行われていることが判明した。当日は、CFAの気体分析部と標準ガスを用いたキャリブレーションの詳細に加え、ドームふじコアからのメタンのCFAによる復元結果を報告する。

キーワード：アイスコア、温室効果ガス、古気候学、南極

Keywords: icecore, greenhouse gas, paleoclimatology, Antarctica

2016年中央アジア・パミール山域レーニン峰アイスコア掘削報告 Preliminary report on the ice core drilling in Pamir-Arai Mountains, Central Asia in 2016

*竹内 望¹、藤田 耕史²、川村 賢二³、對馬 あかね⁴、宮内 謙史郎¹、堀 耀一朗¹、Vladimir Aizen⁵
、Azamat Osmonov⁶

*Nozomu Takeuchi¹, Koji Fujita², Kenji Kawamura³, Akane Tsushima⁴, Kenshiro Miyachi¹,
YOICHIRO HORI¹, Vladimir Aizen⁵, Azamat Osmonov⁶

1. 千葉大学、2. 名古屋大学、3. 国立極地研究所、4. 総合地球環境学研究所、5. アイダホ大学、6. 中央アジア地球物理研究所

1. Chiba University, 2. Nagoya University, 3. National Institute of Polar Research, 4. Research Institute for Humanity and Nature, 5. University of Idaho, 6. Central Asian Institute of Applied Geo-Science

中央アジア山岳域の西端に位置するパミール高原は、アジアモンスーンよりも偏西風の影響を強く受ける中央アジアの中でも特別な気候条件をもつ地域である。パミール高原には数多く氷河が存在し、その融解水は下流部乾燥域の人間社会を支え、最終的にアラル海に注ぐ。この地域の水資源変動の将来予測のためにも、パミール山域の過去の気候変動の復元は重要である。2016年8月に中央アジア・キルギス共和国パミールアライ山域でアイスコアを掘削した。掘削した氷河は、レーニン峰（7134m）から西へ約20km離れた標高約5300mの氷帽頂上である。3本のアイスコアを掘削し、1本目は約7mで水が現れ、2本目は約12mでクレバスに達し、3本目は約37mで底部岩盤に達した。底部に達した3本目のコアは、平均密度は847 kg m⁻³で、15m以深はほぼ再凍結氷であった。掘削高温度の測定の結果は、底部で-6.3°Cであった。以上の結果は、パミール山域では5300mの標高でも氷河表面は融解再凍結が起こること、また掘削したアイスコアは過去環境復元の目的としては限定的な試料であることを示している。アイスコアは今後、同位体、化学分析等が進められる予定である。

キーワード：中央アジア、アイスコア、パミール

Keywords: central asia, ice core, Pamir